

Rookery Bay Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 27 January, 2025

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	3
Water Quality	4
Dissolved Oxygen - Discrete	4
Dissolved Oxygen - Continuous	5
Dissolved Oxygen Saturation - Discrete	6
Dissolved Oxygen Saturation - Continuous	7
Salinity - Discrete	8
Salinity - Continuous	9
Water Temperature - Discrete	10
Water Temperature - Continuous	11
pH - Discrete	12
pH - Continuous	13
Water Clarity	14
Turbidity - Discrete	14
Turbidity - Continuous	15
Total Suspended Solids - Discrete	16
Chlorophyll a, Uncorrected for Pheophytin - Discrete	17
Chlorophyll a, Corrected for Pheophytin - Discrete	18
Secchi Depth - Discrete	19
Colored Dissolved Organic Matter - Discrete	20

Indicators

Nutrients

Total Nitrogen - Discrete




Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	2763	29	1989 - 2024		0.26	0.1757	0.21458	0.00577 0.0002

Total Phosphorus - Discrete




Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P	
Lab	No significant trend	2383	26	1999 - 2024		0.034	0.0277	0.03786	0.00008	0.4884

Water Quality

Dissolved Oxygen - Discrete




Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	4538	33	1989 - 2024	5.83	-0.0701	6.0986	-0.00976	0.0815

Dissolved Oxygen - Continuous




Table 4: Seasonal Kendall-Tau Results for All Stations - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10441	2	1997 - 1998	7.2	-	-	-	NA
rkbllhwq	No significant trend	570691	21	2001 - 2024	5.5	0	5.48	0	0.8577

Dissolved Oxygen Saturation - Discrete




Table 5: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	791	24	1998 - 2024	80.9	0.0693	79.25336	0.18388	0.2483

Dissolved Oxygen Saturation - Continuous




Table 6: Seasonal Kendall-Tau Results for All Stations - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10441	2	1997 - 1998	102.4	-	-	-	NA
rkbllhwq	No significant trend	583009	21	2001 - 2024	81.7	0.02	80.58	0.02	0.5540

Salinity - Discrete




Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	5272	45	1954 - 2024	34.1	0.0144	34.04711	0.0035	0.6726

Salinity - Continuous




Table 8: Seasonal Kendall-Tau Results for All Stations - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12256	2	1997 - 1998	34.5	-	-	-	NA
rkbllhwq	Significantly decreasing trend	657842	24	2001 - 2024	33.1	-0.16	34.68	-0.1	0.0001

Water Temperature - Discrete




Table 9: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	5225	44	1954 - 2024	26.4	0.0768	25.08853	0.01003	0.0273

Water Temperature - Continuous




Table 10: Seasonal Kendall-Tau Results for All Stations - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12610	2	1997 - 1998	23.8	-	-	-	NA
rkbllhwq	Significantly increasing trend	688994	24	2001 - 2024	27.0	0.25	26.12	0.05	0.0000

pH - Discrete




Table 11: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	2450	37	1955 - 2024	7.9	-0.1389	8.03758	-0.00383	0.0004

pH - Continuous




Table 12: Seasonal Kendall-Tau Results for All Stations - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12610	2	1997 - 1998	8.0	-	-	-	NA
rkbllhwq	Significantly increasing trend	629829	24	2001 - 2024	7.9	0.14	7.83	0	0.0010

Water Clarity

Turbidity - Discrete




Table 13: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	1843	31	1989 - 2024	4.995	0.2463	3.33593	0.12308	0.0000

Turbidity - Continuous




Table 14: Seasonal Kendall-Tau Results for All Stations - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10654	2	1997 - 1998		11	-	-	NA
rkbllhwq	Significantly decreasing trend	605017	24	2001 - 2024		8	-0.27	12.4	-0.2

Total Suspended Solids - Discrete




Table 15: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	245	23	1989 - 2024		8	0.1144	3.26636	0.09091 0.0803

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Table 16: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	3141	26	1999 - 2024	5	0.1899	4.55801	0.06768	0.0000

Chlorophyll a, Corrected for Pheophytin - Discrete




Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1546	15	2002 - 2024	4.2	0.1077	3.31555	0.06083	0.0885

Secchi Depth - Discrete




Table 18: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P	
Field	No significant trend	436	21	1998 - 2024		1.2	-0.1218	1.21954	-0.01134	0.1146

Colored Dissolved Organic Matter - Discrete




Table 19: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	183	11	2001 - 2021		7	0.5051	4.7	0.325 0.0515