

Rocky Bayou State Park Aquatic Preserve

SEACAR Habitat Analyses

Last compiled on 03 September, 2024

Contents

Threshold Filtering	2
Value Qualifiers	3
Water Column	5
Seasonal Kendall-Tau Analysis	5
Water Quality - Discrete	5
Chlorophyll a, Corrected for Pheophytin - Discrete Water Quality	6
Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality	9
Colored Dissolved Organic Matter - Discrete Water Quality	11
Dissolved Oxygen - Discrete Water Quality	14
Dissolved Oxygen Saturation - Discrete Water Quality	17
pH - Discrete Water Quality	19
Salinity - Discrete Water Quality	21
Secchi Depth - Discrete Water Quality	23
Total Nitrogen - Discrete Water Quality	25
Total Phosphorus - Discrete Water Quality	28
Turbidity - Discrete Water Quality	30
Water Temperature - Discrete Water Quality	33

Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection's (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR_QAQCFlagCode* and *SEACAR_QAQC_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>	<i>Sensor Type</i>
Dissolved Oxygen	mg/L	0	50	YSI EXOs
Dissolved Oxygen	mg/L	0	50	Analysis Only - 2022-04-04
Dissolved Oxygen	mg/L	0	50	6600 Series
Salinity	ppt	0	70	6600 Series
Salinity	ppt	0	70	YSI EXOs
Salinity	ppt	0	70	Analysis Only - 2022-04-04
Water Temperature	Degrees C	-5	45	YSI EXOs
Water Temperature	Degrees C	-5	45	Analysis Only - 2022-04-04
Water Temperature	Degrees C	-5	45	6600 Series
pH	pH	2	14	Analysis Only - 2022-04-04
pH	pH	2	14	6600 Series
pH	pH	2	14	YSI EXOs
Dissolved Oxygen Saturation	%	0	500	YSI EXOs
Dissolved Oxygen Saturation	%	0	500	6600 Series
Dissolved Oxygen Saturation	%	0	500	Analysis Only - 2022-04-04
Specific Conductivity	mS/cm	0	100	6600 Series
Specific Conductivity	mS/cm	0	200	YSI EXOs
Turbidity	NTU	0	4000	YSI EXOs
Turbidity	NTU	0	1000	6600 Series
Turbidity	NTU	0	4000	Analysis Only - 2022-04-04

Table 2: Discrete Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	0.000001	22
Salinity	ppt	0	70
Water Temperature	Degrees C	3	40
pH		2	13
Dissolved Oxygen Saturation	%	0.000001	310
Specific Conductivity	mS/cm	0.005000001	100
Turbidity	NTU	0	-
Total Suspended Solids (TSS)	mg/L	0	-
Chlorophyll a uncorrected for pheophytin	ug/L	0	-
Chlorophyll a corrected for pheophytin	ug/L	0	-
Secchi Depth	m	0.000001	50
Light Extinction Coefficient	m^{-1}	0	-
Colored dissolved organic matter, CDOM	PCU	0	-
Fluorescent dissolved organic matter, FDOM	QSE	0	-
Total Nitrogen	mg/L	0	-
Total Kjeldahl Nitrogen TKN	mg/L	0	-
NO ₂ +3 Filtered	mg/L	0	-
NH ₄ Filtered	mg/L	0	-
Total Phosphorus	mg/L	0	-

Parameter Name	Units	Low Threshold	High Threshold
PO4 Filtered	mg/L	0	-
Ammonia- Un-ionized (NH3)	mg/L	0	-
Nitrate (N)	mg/L	0	-
Nitrite (N)	mg/L	0	-
Nitrogen, organic	mg/L	0	-

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 & 2

SEACAR QAQC Description	Include	SEACAR QAQCFlagCode
Exceeds Maximum threshold. Not verified in raw data	No	2Q
Exceeds Maximum threshold. Verified in raw data	No	3Q
Below Minimum threshold. Not verified in raw data	No	4Q
Below Minimum threshold. Verified in raw data	No	5Q
Within threshold tolerance	Yes	6Q
No defined thresholds for this parameter	Yes	7Q

Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

Qualifier Source	Value Qualifier	Include	MDL	Description
STORET-WIN	H	No	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	No	0	Estimated value
STORET-WIN	V	No	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	No	0	Lab analysis from an improperly preserved sample; data may be inaccurate

Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

H - Value based on field kit determination; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

I - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

Q - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

S - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

U - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported.

Systemwide Monitoring Program (SWMP) value qualifier codes

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	Yes	Optional parameter not collected
SWMP	-2	No	Missing data
SWMP	-3	No	Data rejected due to QA/QC
SWMP	-4	No	Outside low sensor range
SWMP	-5	No	Outside high sensor range
SWMP	0	Yes	Passed initial QA/QC checks
SWMP	1	No	Suspect data
SWMP	2	Yes	Reserved for future use
SWMP	3	Yes	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	Yes	Historical: Pre-auto QA/QC
SWMP	5	Yes	Corrected data

Water Column

The water column habitat extends from the surface of all water bodies to the bottom sediments and encompasses the different features found in the water at different depths (National Oceanographic Center, 2016). The water column habitat must be viewed in relation to its interconnectedness with other habitats. A healthy water column is an integral component in ensuring a healthy marine and coastal ecosystem. Having a flourishing marine and coastal ecosystem in Florida is necessary to support a strong economy. The health of the water column is dependent upon factors as diverse as land use (e.g., agriculture, mining, forestry practices); human population growth; emissions, (e.g., power plants, automobiles, wastewater); climate (e.g., rainfall, temperature, winds and currents); and decadal trends (e.g., El Niño/La Niña, Atlantic Multidecadal Oscillation, climate change).

The water column is composed of various physical, chemical and biological features, and only a small number of them are adequately monitored. Features of the water column that are monitored are used as indicators of the water column health and help assess the status of other habitats. These indicators include nutrient concentrations (nitrogen and phosphorus); water quality (dissolved oxygen, temperature, salinity and pH); water clarity (Secchi depth, turbidity, chlorophyll-a and colored dissolved organic matter); and nekton (fish, macroinvertebrates and megafauna).

Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as “insufficient data to conduct analysis”. Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use_In_Analysis* = **TRUE**. Any that fail either test will be excluded from the analyses and labeled as *Use_In_Analysis* = **FALSE**.

Water Quality - Discrete


The following files were used in the discrete analysis:

- *Combined_WQ_WC_NUT_Chlorophyll_a_corrected_for_pheophytin-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Chlorophyll_a_uncorrected_for_pheophytin-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Colored_dissolved_organic_matter_CDOM-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen_Saturation-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_pH-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Salinity-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Secchi_Depth-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Total_Nitrogen-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Total_Phosphorus-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Total_Suspended_Solids_TSS-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Turbidity-2024-Jul-11.txt*
- *Combined_WQ_WC_NUT_Water_Temperature-2024-Jul-11.txt*

Chlorophyll a, Corrected for Pheophytin - Discrete Water Quality

Chlorophyll-a is monitored as a measure of microalgae growing in the water. Algae are a natural part of coastal and aquatic ecosystems but in excess can cause poor water quality and clarity, and decreased levels of dissolved oxygen.


Seasonal Kendall-Tau Trend Analysis



$p < 0.00005$ appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Chlorophyll a, Corrected for Pheophytin



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 6: Programs contributing data for Chlorophyll a, Corrected for Pheophytin

ProgramID	N_Data	YearMin	YearMax
5002	59	2012	2016
537	35	2018	2023
514	19	2020	2022

Program names:

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

514 - Florida LAKEWATCH Program

Value Qualifiers

- N_{Total} is total amount of data for a given year
- $N_{_}$ is the total amount of values flagged with the respective value qualifier in a given year
- $perc_{_}$ is the percent of data flagged with the respective value qualifier as a proportion of N_{Total}

Table 7: Value Qualifiers for Chlorophyll a, Corrected for Pheophytin

<i>Year</i>	<i>N_Total</i>	<i>N_I</i>	<i>perc_I</i>	<i>N_Q</i>	<i>perc_Q</i>	<i>N_U</i>	<i>perc_U</i>
2015	15					2	13.3
2016	17	4	23.5			1	5.9
2018	1					1	100.0
2019	8	2	25.0	2	25		
2021	17	6	35.3			3	17.6
2022	15	1	6.7			2	13.3

Note: ¹**I** - Reported value is greater than or equal to lab method detection limit, but less than quantitation limit ²**Q**
 - Sample held beyond the accepted holding time ³**U** - Compound was analyzed for but not detected

Programs containing Value Qualified data:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring


514 - Florida LAKEWATCH Program

5002 - Florida STORET / WIN

Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality

Seasonal Kendall-Tau Trend Analysis

Chlorophyll a, Uncorrected for Pheophytin, Lab, All Depths
Rocky Bayou State Park Aquatic Preserve



RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	462	23	3.5	TRUE	-0.0142	0.8100	0	3.536806	8.9441	0.627	0

$p < 0.00005$ appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Chlorophyll a, Uncorrected for Pheophytin




Table 8: Programs contributing data for Chlorophyll a, Uncorrected for Pheophytin

ProgramID	N_Data	YearMin	YearMax
514	223	2001	2022
5002	182	2001	2016
537	44	2018	2023
103	13	2020	2021

Program names:

514 - Florida LAKEWATCH Program

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

103 - EPA STOrage and RETrieval Data Warehouse (STORET)

Value Qualifiers

- N_{Total} is total amount of data for a given year
- $N_{}$ is the total amount of values flagged with the respective value qualifier in a given year
- $perc_{}$ is the percent of data flagged with the respective value qualifier as a proportion of N_{Total}

Table 9: Value Qualifiers for Chlorophyll a, Uncorrected for Pheophytin

Year	N_{Total}	N_I	$perc_I$	N_Q	$perc_Q$	N_U	$perc_U$
2015	25					1	4.0
2016	29	2	6.9			1	3.5
2018	13	1	7.7				
2019	20	1	5.0	2	10		

Note: ¹**I** - Reported value is greater than or equal to lab method detection limit, but less than quantitation limit ²**Q** - Sample held beyond the accepted holding time ³**U** - Compound was analyzed for but not detected

Programs containing Value Qualified data:

5002 - Florida STORET / WIN


537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

Colored Dissolved Organic Matter - Discrete Water Quality

Colored Dissolved Organic Matter (CDOM) occurs naturally in every water body. It is made up of mainly plant material, algae and bacteria. The composition is determined by its source; plants, soil, algae, and wastewater are common sources.

Seasonal Kendall-Tau Trend Analysis


Colored Dissolved Organic Matter, Lab, All Depths
Rocky Bayou State Park Aquatic Preserve



p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Colored Dissolved Organic Matter



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 10: Programs contributing data for Colored Dissolved Organic Matter

ProgramID	N_Data	YearMin	YearMax
514	112	2001	2022
537	17	2019	2023

Program names:

514 - Florida LAKEWATCH Program

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

Value Qualifiers

- N_{Total} is total amount of data for a given year
- $N_{_}$ is the total amount of values flagged with the respective value qualifier in a given year

- $perc_Q$ is the percent of data flagged with the respective value qualifier as a proportion of N_Total

Table 11: Value Qualifiers for Colored Dissolved Organic Matter

<i>Year</i>	<i>N_Total</i>	<i>N_Q</i>	<i>perc_Q</i>
2019	7	4	57.1
2020	6	6	100.0
2021	5	5	100.0
2022	4	4	100.0
2023	2	2	100.0

Note: ¹**Q** - Sample held beyond the accepted holding time

Programs containing Value Qualified data:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring


514 - Florida LAKEWATCH Program

Dissolved Oxygen - Discrete Water Quality

Dissolved Oxygen (DO) is a key indicator of water quality. Oxygen enters surface waters by air-sea gas exchange, by wind action, or as a byproduct of aquatic plant photosynthesis. The actual quantity of DO in aquatic environments is dependent on the above processes as well as water temperature and salinity.

Seasonal Kendall-Tau Trend Analysis

Dissolved Oxygen, Field, All Depths
Rocky Bayou State Park Aquatic Preserve




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	671	24	6.56	TRUE	0.136	0.0032	0.02714286	5.596685	12.6725	0.3153	1

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Dissolved Oxygen



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 12: Programs contributing data for Dissolved Oxygen

ProgramID	N_Data	YearMin	YearMax
5002	345	2001	2016
537	290	2008	2023
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

Program names:

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring


69 - Fisheries-Independent Monitoring (FIM) Program

103 - EPA STOrage and RETrieval Data Warehouse (STORET)

There are no qualifying Value Qualifiers for Dissolved Oxygen in Rocky Bayou State Park Aquatic Preserve

Dissolved Oxygen Saturation - Discrete Water Quality


Seasonal Kendall-Tau Trend Analysis



$p < 0.00005$ appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Dissolved Oxygen Saturation



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 13: Programs contributing data for Dissolved Oxygen Saturation

ProgramID	N_Data	YearMin	YearMax
5002	346	2000	2016
537	288	2008	2023

Program names:

5002 - Florida STORET / WIN


537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

There are no qualifying Value Qualifiers for Dissolved Oxygen Saturation in Rocky Bayou State Park Aquatic Preserve

pH - Discrete Water Quality

The **pH** of water is the measure of how acidic or basic the water body is on a scale of 0-14, with lower readings indicating acidic and higher readings indicating basic, and a pH of 7 being neutral. Florida's natural waters fall between 6.5 and 8.5 on this scale. A water body's pH can change due to precipitation, geology, vegetation, water pollution and air pollution.


Seasonal Kendall-Tau Trend Analysis



$p < 0.00005$ appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for pH



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 14: Programs contributing data for pH

ProgramID	N_Data	YearMin	YearMax
5002	345	2000	2016
537	290	2008	2023
69	20	1994	2022
103	12	2020	2021

Program names:

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

69 - Fisheries-Independent Monitoring (FIM) Program


103 - EPA STOrage and RETrieval Data Warehouse (STORET)

There are no qualifying Value Qualifiers for pH in Rocky Bayou State Park Aquatic Preserve

Salinity - Discrete Water Quality

Salinity is a measure of the amount of salt in the water. In estuarine ecosystems, salinity is influenced by precipitation, evaporation, surface-water inputs, and exchange with coastal waters.

Seasonal Kendall-Tau Trend Analysis




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	658	25	21.94	TRUE	0.0248	0.7064	0.01623932	20.89179	11.8113	0.378	0

$p < 0.00005$ appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Salinity



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 15: Programs contributing data for Salinity

ProgramID	N_Data	YearMin	YearMax
5002	346	2000	2016
537	288	2008	2023
69	20	1994	2022
102	4	1994	1994

Program names:

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

69 - Fisheries-Independent Monitoring (FIM) Program


102 - National Status and Trends Mussel Watch

There are no qualifying Value Qualifiers for Salinity in Rocky Bayou State Park Aquatic Preserve

Secchi Depth - Discrete Water Quality

Secchi depth is a measure of the transparency or clarity of the water by a device called a Secchi disk. A Secchi disk is a black and white disk that is lowered into the water on a cord. The Secchi depth is the depth at which the disk can no longer be seen. The deeper the Secchi depth, the greater the water clarity.

Seasonal Kendall-Tau Trend Analysis




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
Surface	474	24	2.4384	TRUE	-0.1713	0.0003	-0.02665347	2.795615	9.3193	0.5924	-1

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Secchi Depth



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 16: Programs contributing data for Secchi Depth

ProgramID	N_Data	YearMin	YearMax
514	227	2001	2022
537	220	2001	2023
5002	15	2013	2016
69	12	1995	2022

Program names:

514 - Florida LAKEWATCH Program

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

5002 - Florida STORET / WIN

69 - Fisheries-Independent Monitoring (FIM) Program

There are no qualifying Value Qualifiers for Secchi Depth in Rocky Bayou State Park Aquatic Preserve

Total Nitrogen - Discrete Water Quality

Nitrogen and **Phosphorous** are key nutrients that provide nourishment essential for the growth and maintenance of aquatic plants and animals; however, excess nutrients can cause harmful algal blooms and other water quality concerns. Nutrients enter water bodies several ways, including runoff from rain events and atmospheric deposition from natural and industrial sources.

Total Nitrogen Calculation:


The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1) $TN = TKN + NO_3O_2;$
- 2) $TN = TKN + NO_3 + NO_2;$
- 3) $TN = ORGN + NH_4 + NO_3O_2;$
- 4) $TN = ORGN + NH_4 + NO_2 + NO_3;$
- 5) $TN = TKN + NO_3;$
- 6) $TN = ORGN + NH_4 + NO_3;$

Additional Information:

- Rules for use of sample fraction:
 - FDEP report that if both “Total” and “Dissolved” are reported, only “Total” is used. If the total is not reported, they do use dissolved as a best available replacement.
 - An analysis of all SEACAR data shows that 90% of all possible TN calculations can be done using nitrogen components with the same sample fraction, rather than use nitrogen components with mixed total/dissolved sample fractions. In other words, TN can be calculated when TKN and NO_3O_2 are both total sample fraction, or when both are dissolved sample fraction. This is important, because then the calculated TN value is not based on components with mixed sample fractions.
- Values inserted into data:
 - ParameterName = “Total Nitrogen”
 - SEACAR_QAQCFlagCode = “1Q”
 - SEACAR_QAQC_Description = “SEACAR Calculated”

Seasonal Kendall-Tau Trend Analysis




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	642	23	0.2	TRUE	0.125	0.0044	0.001907143	0.1456745	6.5856	0.8316	1

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Total Nitrogen



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 17: Programs contributing data for Total Nitrogen

ProgramID	N_Data	YearMin	YearMax
537	234	2001	2023
514	223	2001	2022
5002	185	2001	2016

Program names:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

514 - Florida LAKEWATCH Program

5002 - Florida STORET / WIN

Value Qualifiers

- N_{Total} is total amount of data for a given year
- $N_{\text{Qualifier}}$ is the total amount of values flagged with the respective value qualifier in a given year
- $\text{perc}_{\text{Qualifier}}$ is the percent of data flagged with the respective value qualifier as a proportion of N_{Total}

Table 18: Value Qualifiers for Total Nitrogen

Year	N_Total	N_Q	perc_Q
2014	35	3	8.6
2020	16	3	18.8

Note: ¹Q - Sample held beyond the accepted holding time

Programs containing Value Qualified data:


537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

5002 - Florida STORET / WIN

514 - Florida LAKEWATCH Program

Total Phosphorus - Discrete Water Quality

Seasonal Kendall-Tau Trend Analysis




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	472	23	0.007	TRUE	0.1913	0.0000	0.0001402202	0.004218804	8.4735	0.6704	1

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Total Phosphorus



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 19: Programs contributing data for Total Phosphorus

ProgramID	N_Data	YearMin	YearMax
537	234	2001	2023
514	225	2001	2022
5002	16	2013	2016
103	8	2020	2021

Program names:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

514 - Florida LAKEWATCH Program

5002 - Florida STORET / WIN

103 - EPA STOrage and RETrieval Data Warehouse (STORET)

Value Qualifiers

- N_{Total} is total amount of data for a given year
- $N_{}$ is the total amount of values flagged with the respective value qualifier in a given year
- $perc_{}$ is the percent of data flagged with the respective value qualifier as a proportion of N_{Total}

Table 20: Value Qualifiers for Total Phosphorus

<i>Year</i>	<i>N_Total</i>	<i>N_I</i>	<i>perc_I</i>	<i>N_Q</i>	<i>perc_Q</i>
2013	29	2	6.9		
2015	30	4	13.3		
2016	29	5	17.2		
2018	14	1	7.1		
2019	21	2	9.5		
2020	18	7	38.9	2	11.1
2021	29	10	34.5	2	6.9
2022	14	7	50.0		
2023	7	5	71.4		

Note: ¹**I** - Reported value is greater than or equal to lab method detection limit, but less than quantitation limit ²**Q**
 - Sample held beyond the accepted holding time

Programs containing Value Qualified data:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

5002 - Florida STORET / WIN


514 - Florida LAKEWATCH Program

Turbidity - Discrete Water Quality

Turbidity results from suspended solids in the water, including silts, clays, tannins, industrial wastes, sewage and plankton, which are all factors that contribute to how clouded or murky a water column is. Turbidity is caused by soil erosion, excess nutrients, pollutants, and physical forces such as winds, currents and bottom feeders.

Seasonal Kendall-Tau Trend Analysis

Turbidity, Lab and Field Combined, All Depths
Rocky Bayou State Park Aquatic Preserve




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	597	22	1.1	TRUE	-0.0674	0.2204	-0.02222222	1.897656	8.353	0.6814	0

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Turbidity



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 21: Programs contributing data for Turbidity

ProgramID	N_Data	YearMin	YearMax
5002	303	2001	2016
537	282	2008	2023
103	12	2020	2021

Program names:

5002 - Florida STORET / WIN

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring


103 - EPA STOrage and RETrieval Data Warehouse (STORET)

There are no qualifying Value Qualifiers for Turbidity in Rocky Bayou State Park Aquatic Preserve

Water Temperature - Discrete Water Quality

Temperature determines the capacity of water to hold oxygen. Cooler water can hold more dissolved oxygen because water molecules are more tightly packed, making it harder for oxygen to escape. Additionally, as water temperature increases, fish and other aquatic organisms become more active and consume oxygen at a faster rate.

Seasonal Kendall-Tau Trend Analysis




RelativeDepth	N_Data	N_Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	363	17	24.05556	TRUE	0.1022	0.0877	0.04635312	22.49018	18.0087	0.0814	0

p < 0.00005 appear as 0 due to rounding.

SennIntercept is intercept value at beginning of record for monitoring location

Map showing location of Discrete sampling sites for Water Temperature



The bubble size on the above plots reflects the amount of data available at each sampling site

Table 22: Programs contributing data for Water Temperature

ProgramID	N_Data	YearMin	YearMax
537	290	2008	2023
5002	37	2013	2016
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

Program names:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring

5002 - Florida STORET / WIN

69 - Fisheries-Independent Monitoring (FIM) Program

103 - EPA STOrage and RETrieval Data Warehouse (STORET)

102 - National Status and Trends Mussel Watch

There are no qualifying Value Qualifiers for Water Temperature in Rocky Bayou State Park Aquatic Preserve