

Rocky Bayou State Park Aquatic Preserve

SEACAR Habitat Analyses

Last compiled on 22 May, 2025

Contents

Funding & Acknowledgements	2
Threshold Filtering	2
Value Qualifiers	3
Water Column	5
Seasonal Kendall-Tau Analysis	5
Water Quality - Discrete	5
Chlorophyll a, Corrected for Pheophytin - Discrete	6
Chlorophyll a, Uncorrected for Pheophytin - Discrete	7
Colored Dissolved Organic Matter - Discrete	9
Dissolved Oxygen - Discrete	11
Dissolved Oxygen Saturation - Discrete	14
pH - Discrete	15
Salinity - Discrete	17
Secchi Depth - Discrete	19
Total Nitrogen - Discrete	21
Total Phosphorus - Discrete	24
Turbidity - Discrete	26
Water Temperature - Discrete	28
References	31

Funding & Acknowledgements

The data used in this analysis is from the Export Standardized Tables in the SEACAR Data Discovery Interface (DDI). Documents and information available through the SEACAR DDI are owned by the data provider(s) and users are expected to provide appropriate credit following accepted citation formats. Users are encouraged to access data to maximize utilization of gained knowledge, reducing redundant research and facilitating partnerships and scientific innovation.

With respect to documents and information available from SEACAR DDI, neither the State of Florida nor the Florida Department of Environmental Protection makes any warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose arising out of the use or inability to use the data, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

This report was funded in part, through a grant agreement from the Florida Department of Environmental Protection, Florida Coastal Management Program, by a grant provided by the Office for Coastal Management under the Coastal Zone Management Act of 1972, as amended, National Oceanic and Atmospheric Administration. The views, statements, findings, conclusions and recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of the State of Florida, NOAA or any of their sub agencies.

Published: 2025-05-22



Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection's (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR_QAQCFlagCode* and *SEACAR_QAQC_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

Parameter Name	Units	Low Threshold	High Threshold
Dissolved Oxygen	mg/L	-0.000001	50
Dissolved Oxygen Saturation	%	-0.000001	500
Salinity	ppt	-0.000001	70
Turbidity	NTU	-0.000001	4000
Water Temperature	Degrees C	-5.000000	45
pH	None	2.000000	14

Table 2: Discrete Water Quality threshold values

Parameter Name	Units	Low Threshold	High Threshold
Ammonia, Un-ionized (NH3)	mg/L	-	-
Ammonium, Filtered (NH4)	mg/L	-	-
Chlorophyll a, Corrected for Pheophytin	ug/L	-	-
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Colored Dissolved Organic Matter	PCU	-	-

Parameter Name	Units	Low Threshold	High Threshold
Dissolved Oxygen	mg/L	-0.000001	25
Dissolved Oxygen Saturation	%	-0.000001	310
Fluorescent Dissolved Organic Matter	QSE	-	-
Light Extinction Coefficient	m^-1	-	-
NO2+3, Filtered	mg/L	-	-
Nitrate (NO3)	mg/L	-	-
Nitrite (NO2)	mg/L	-	-
Nitrogen, organic	mg/L	-	-
Phosphate, Filtered (PO4)	mg/L	-	-
Salinity	ppt	-0.000001	70
Secchi Depth	m	0.000001	50
Specific Conductivity	mS/cm	0.005000	100
Total Kjeldahl Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Phosphorus	mg/L	-	-
Total Suspended Solids	mg/L	-	-
Turbidity	NTU	-	-
Water Temperature	Degrees C	3.000000	40
pH	None	2.000000	13

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 and 2

SEACAR QAQC Description	Include	SEACAR QAQCFlagCode
Exceeds maximum threshold	0	2Q
Below minimum threshold	0	4Q
Within threshold tolerance	1	6Q
No defined thresholds for this parameter	1	7Q

Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

Qualifier Source	Value Qualifier	Include	MDL	Description
STORET-WIN	H	0	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	0	0	Estimated value
STORET-WIN	V	0	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	0	0	Lab analysis from an improperly preserved sample; data may be inaccurate

Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

H - Value based on field kit determiniation; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

I - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

Q - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

S - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

U - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported

Systemwide Monitoring Program (SWMP) value qualifier codes

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	1	Optional parameter not collected
SWMP	-2	0	Missing data
SWMP	-3	0	Data rejected due to QA/QC
SWMP	-4	0	Outside low sensor range
SWMP	-5	0	Outside high sensor range
SWMP	0	1	Passed initial QA/QC checks
SWMP	1	0	Suspect data
SWMP	2	1	Reserved for future use
SWMP	3	1	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	1	Historical: Pre-auto QA/QC
SWMP	5	1	Corrected data

Water Column

The water column habitat extends from the water's surface to the bottom sediments, and it's where fish, dolphins, crabs and people swim! So much life makes its home in the water column that the health of marine and coastal ecosystems, as well as human economies, depend on the condition of this vulnerable habitat. Local patterns of rainfall, temperature, winds and currents can rapidly change the condition of the water column, while global influences such as [El Niño/La Niña](#), large-scale fluctuation in sea temperatures and climate change can have long-term effects. Inputs from the prosperity of our day-to-day lives including farming, mining and forestry, and emissions from power generation, automobiles and water treatment can also alter the health of the water column. Acting alone or together, each input can have complex and lasting effects on habitats and ecosystems.

SEACAR evaluates water column health with several essential parameters. These include nutrient surveys of nitrogen and phosphorus, and water quality assessments of salinity, dissolved oxygen, pH, and water temperature. Water clarity is evaluated with Secchi depth, turbidity, levels of chlorophyll a, total suspended solids, and colored dissolved organic matter. Additionally, the richness of nekton is indicated by the abundance of free-swimming fishes and macroinvertebrates like crabs and shrimps.

Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as "insufficient data to conduct analysis". Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use_In_Analysis = TRUE*. Any that fail either test will be excluded from the analyses and labeled as *Use_In_Analysis = FALSE*. The points for all Water Column plots displayed in this section are monthly averages. Trend significance will be denoted as "Significant Trend" (when $p < 0.05$), or "Non-significant Trend" (when $p \geq 0.05$). Any parameters with insufficient data to perform Seasonal Kendall-Tau test will have their monthly averages plotted without a corresponding trend line.

Water Quality - Discrete

The following files were used in the discrete analysis:

- *Combined_WQ_WC_NUT_Chlorophyll_a_corrected_for_pheophytin-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Chlorophyll_a_uncorrected_for_pheophytin-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Colored_dissolved_organic_matter_CDOM-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen_Saturation-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_pH-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Salinity-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Secchi_Depth-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Nitrogen-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Phosphorus-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Suspended_Solids_TSS-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Turbidity-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Water_Temperature-2025-Mar-06.txt*

Chlorophyll a, Corrected for Pheophytin - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 1: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 6: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	No significant trend	103	13	2012 - 2024	3	-0.1845	3.2357	-0.025	0.0504

Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2012 and 2024.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 7: Programs contributing data for Chlorophyll a, Corrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	60	2012	2017
537	43	2018	2024

Program names:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
 5002 - Florida STORET / WIN²

Chlorophyll a, Uncorrected for Pheophytin - Discrete Seasonal Kendall-Tau Trend Analysis




Figure 3: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 8: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	No significant trend	269	24	2001 - 2024	4	-0.0018	3.7917	0	0.9007

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 2001 and 2024.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 9: Programs contributing data for Chlorophyll a, Uncorrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	183	2001	2017
537	52	2018	2024
514	21	2018	2019
103	13	2020	2021

Program names:

- 103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³
- 514 - Florida LAKEWATCH Program⁴
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
- 5002 - Florida STORET / WIN²

Colored Dissolved Organic Matter - Discrete Seasonal Kendall-Tau Trend Analysis




Figure 5: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 10: Seasonal Kendall-Tau Trend Analysis for Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	No significant trend	128	24	2001 - 2024	13.5	0.0916	9.6216	0.1818	0.3155

Colored dissolved organic matter showed no detectable trend between 2001 and 2024.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 11: Programs contributing data for Colored Dissolved Organic Matter

ProgramID	N_Data	YearMin	YearMax
514	108	2001	2019
537	20	2019	2024

Program names:

514 - Florida LAKEWATCH Program⁴

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

Dissolved Oxygen - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	Significantly increasing trend	822	26	1994 - 2024	6.625	0.1142	5.5936	0.0225	0.0088

Monthly average dissolved oxygen increased by 0.02 mg/L per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 13: Programs contributing data for Dissolved Oxygen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	395	2001	2018
537	391	2008	2024
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

Program names:

- 69 - Fisheries-Independent Monitoring (FIM) Program⁵
- 102 - National Status and Trends Mussel Watch⁶
- 103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
- 5002 - Florida STORET / WIN²

Dissolved Oxygen Saturation - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 14: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	Significantly increasing trend	735	25	2000 - 2024	89.3	0.1529	70.1717	0.4611	0.0008

Monthly average dissolved oxygen saturation increased by 0.46% per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 15: Programs contributing data for Dissolved Oxygen Saturation

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	387	2008	2024
5002	348	2000	2017

Program names:

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
 5002 - Florida STORET / WIN²

pH - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 11: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 16: Seasonal Kendall-Tau Trend Analysis for pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	No significant trend	841	27	1994 - 2024	7.89	0.0319	7.8603	0.0019	0.4326

pH showed no detectable trend between 1994 and 2024.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 17: Programs contributing data for pH

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	418	2000	2019
537	391	2008	2024
69	20	1994	2022
103	12	2020	2021

Program names:

69 - Fisheries-Independent Monitoring (FIM) Program⁵

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

5002 - Florida STORET / WIN²

Salinity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 18: Seasonal Kendall-Tau Trend Analysis for Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
All	No significant trend	832	27	1994 - 2024	21.535	0.0488	21.1466	0.039	0.3053

Salinity showed no detectable trend between 1994 and 2024.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 19: Programs contributing data for Salinity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	420	2000	2019
537	388	2008	2024
69	20	1994	2022
102	4	1994	1994

Program names:

69 - Fisheries-Independent Monitoring (FIM) Program⁵

102 - National Status and Trends Mussel Watch⁶

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

5002 - Florida STORET / WIN²

Secchi Depth - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 15: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 20: Seasonal Kendall-Tau Trend Analysis for Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	Significantly increasing trend	458	25	1995 - 2024	-2.4384	0.1678	-2.7703	0.0254	0.0003

Monthly average Secchi depth became shallower by 0.03 m per year, indicating a decrease in water clarity.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 21: Programs contributing data for Secchi Depth

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	229	2001	2024
514	201	2001	2019
5002	16	2013	2017
69	12	1995	2022

Program names:

69 - Fisheries-Independent Monitoring (FIM) Program⁵

514 - Florida LAKEWATCH Program⁴

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

5002 - Florida STORET / WIN²

Total Nitrogen - Discrete

Total Nitrogen Calculation:

The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1) $TN = TKN + NO_3O_2;$
- 2) $TN = TKN + NO_3 + NO_2;$
- 3) $TN = ORGN + NH_4 + NO_3O_2;$
- 4) $TN = ORGN + NH_4 + NO_2 + NO_3;$
- 5) $TN = TKN + NO_3;$
- 6) $TN = ORGN + NH_4 + NO_3;$

Additional Information:

- Rules for use of sample fraction:
 - Florida Department of Environmental Protection (FDEP) report that if both “Total” and “Dissolved” components are reported, only “Total” is used. If the total is not reported, then the dissolved components are used as a best available replacement.
 - Total nitrogen calculations are done using nitrogen components with the same sample fraction, nitrogen components with mixed total/dissolved sample fractions are not used. In other words, total nitrogen can be calculated when TKN and NO₃O₂ are both total sample fractions, or when both are dissolved sample fractions. *Future calculations of total nitrogen values may be based on components with mixed sample fractions.*
- Values inserted into data:
 - ParameterName = “Total Nitrogen”
 - SEACAR_QAQCFlagCode = “1Q”
 - SEACAR_QAQC_Description = “SEACAR Calculated”

Seasonal Kendall-Tau Trend Analysis




Figure 17: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 22: Seasonal Kendall-Tau Trend Analysis for Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly increasing trend	635	24	2001 - 2024	0.2	0.1611	0.1462	0.0022	0.0002

Monthly average total nitrogen increased by less than 0.01 mg/L per year.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 23: Programs contributing data for Total Nitrogen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	246	2001	2024
514	203	2001	2019
5002	186	2001	2017

Program names:

514 - Florida LAKEWATCH Program⁴

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

5002 - Florida STORET / WIN²

Total Phosphorus - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 19: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 24: Seasonal Kendall-Tau Trend Analysis for Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly increasing trend	465	24	2001 - 2024	0.007	0.2622	0.0035	0.0002	0

Monthly average total phosphorus increased by less than 0.01 mg/L per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 25: Programs contributing data for Total Phosphorus

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	246	2001	2024
514	203	2001	2019
5002	17	2013	2017
103	8	2020	2021

Program names:

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³

514 - Florida LAKEWATCH Program⁴

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹

5002 - Florida STORET / WIN²

Turbidity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 21: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 26: Seasonal Kendall-Tau Trend Analysis for Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly decreasing trend	222	11	2008 - 2018	1	-0.2377	3.4417	-0.2134	0.0034

Monthly average turbidity decreased by 0.21 NTU per year, indicating an increase in water clarity.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 27: Programs contributing data for Turbidity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	379	2008	2024
5002	304	2001	2017
103	12	2020	2021

Program names:

- 103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
- 5002 - Florida STORET / WIN²

Water Temperature - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 23: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 28: Seasonal Kendall-Tau Trend Analysis for Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	Significantly increasing trend	652	19	1994 - 2024	24.7778	0.1627	22.8548	0.0727	0.0027

Monthly average water temperature increased by 0.07°C per year.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 29: Programs contributing data for Water Temperature

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	505	2008	2024
5002	111	2013	2019
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

Program names:

- 69 - Fisheries-Independent Monitoring (FIM) Program⁵
- 102 - National Status and Trends Mussel Watch⁶
- 103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX³
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring¹
- 5002 - Florida STORET / WIN²

References

1. Choctawhatchee Basin Alliance. [Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring](#). (2024).
2. Florida Department of Environmental Protection (DEP). [Florida STORET / WIN](#). (2024).
3. U.S. Environmental Protection Agency (EPA). [EPA STOrage and RETrieval Data Warehouse \(STORET\)/WQX](#). (2023).
4. University of Florida (UF); Institute of Food and Agricultural Sciences. [Florida LAKEWATCH Program](#). (2024).
5. Florida Fish and Wildlife Conservation Commission (FWC). [Fisheries-Independent Monitoring \(FIM\) Program](#). (2022).
6. National Oceanic and Atmospheric Administration (NOAA); Center for Coastal Monitoring and Assessment. [National Status and Trends Mussel Watch](#). (2000).