

Gasparilla Sound-Charlotte Harbor Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	7145	33	1993 - 2025	0.708	0.11573	0.61993	0.00362	0.0022

Monthly average total nitrogen increased by less than 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	6166	33	1993 - 2025	0.101	0.05058	0.11913	0.00045	0.1195

Total phosphorus showed no detectable trend between 1993 and 2025.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
CHEW1	Insufficient data to calculate trend	51073	2	2024 - 2025	6.3	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	60923	38	1971 - 2025	6.5	-0.15246	6.9639	-0.01177	0

Monthly average dissolved oxygen decreased by 0.01 mg/L per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
CHEW1	Insufficient data to calculate trend	51071	2	2024 - 2025	89.3	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	957	16	1992 - 2025	87.7	0.47686	17.45312	2.38521	0

Monthly average dissolved oxygen saturation increased by 2.39% per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	Significantly decreasing trend	76169	50	1954 - 2025	26.69	-0.2622	32.93476	-0.10558	0

Monthly average salinity decreased by 0.11 ppt per year.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293252	Insufficient data to calculate trend	1002	4	2011 - 2014	30.0	-	-	-	-
CHEW1	Insufficient data to calculate trend	64435	2	2024 - 2025	27.3	-	-	-	-

There was insufficient data to fit a model for two locations.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293252	Insufficient data to calculate trend	1003	4	2011 - 2014	26.4	-	-	-	-
CHEW1	Insufficient data to calculate trend	65195	2	2024 - 2025	27.1	-	-	-	-

There was insufficient data to fit a model for two locations.




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	62002	50	1954 - 2025	26.1	0.14694	24.70554	0.01637	0

Monthly average water temperature increased by 0.02°C per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
CHEW1	Insufficient data to calculate trend	58569	2	2024 - 2025	7.9	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	57843	38	1955 - 2025	7.9	0.02342	7.87425	0.0003	0.5201

pH showed no detectable trend between 1955 and 2025.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Table 13: Seasonal Kendall-Tau Results - Specific Conductivity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
CHEW1	Insufficient data to calculate trend	64435	2	2024 - 2025	42.52	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 25: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
CHEW1	Insufficient data to calculate trend	47402	2	2024 - 2025	2	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 28: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	9707	31	1995 - 2025	1.8	-0.19001	2.73838	-0.02676	0

Monthly average turbidity decreased by 0.03 NTU per year, indicating an increase in water clarity.




Figure 29: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 30: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	5278	30	1996 - 2025	9	0.10305	10.18973	0.11142	0.007

Monthly average total suspended solids increased by 0.11 mg/L per year, indicating a decrease in water clarity.




Figure 31: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 32: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	2482	29	1997 - 2025	4.86	-0.13253	6.77022	-0.06211	0.0036

Monthly average chlorophyll a, uncorrected for pheophytin, decreased by 0.06 µg/L per year, indicating an increase in water clarity.




Figure 33: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 34: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	3013	21	2001 - 2025	3.1	-0.23488	6.40635	-0.15174	0

Monthly average chlorophyll a, corrected for pheophytin, decreased by 0.15 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 35: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 36: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	28461	32	1994 - 2025	-0.9	-0.08062	-1.13239	-0.0035	0.0299

Secchi depth showed no detectable trend between 1994 and 2025.




Figure 37: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 38: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	1203	9	2017 - 2025	11	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 39: Map showing location of discrete water quality sampling locations within the boundaries of *Gasparilla Sound-Charlotte Harbor Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.