

Kristin Jacobs Coral Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Discrete	6
Dissolved Oxygen Saturation - Discrete	8
Salinity - Discrete	10
Water Temperature - Continuous	11
Water Temperature - Discrete	16
pH - Discrete	18
Water Clarity	20
Turbidity - Discrete	20
Total Suspended Solids - Discrete	22
Chlorophyll a, Uncorrected for Pheophytin - Discrete	24
Chlorophyll a, Corrected for Pheophytin - Discrete	26
Secchi Depth - Discrete	28
Colored Dissolved Organic Matter - Discrete	30

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	871	7	2009 - 2023	0.07	-	-	-	-

There was insufficient data to fit a model for total nitrogen.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	17904	15	2009 - 2025	0.014	-0.33625	0.02424	-0.00091	1e-04

Monthly average total phosphorus decreased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	715	18	1972 - 2024	6.5	0.10593	5.81879	0.01349	0.3679

Dissolved oxygen showed no detectable trend between 1972 and 2024.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	1418	14	1996 - 2025	101.5	0.37642	47.31868	1.46683	1e-04

Monthly average dissolved oxygen saturation increased by 1.47% per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 9: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	Significantly increasing trend	1895	29	1972 - 2025	36	0.26277	30.19797	0.08842	3e-04

Monthly average salinity increased by 0.09 ppt per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous

National Data Buoy Center - 5




Figure 11: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 6: Seasonal Kendall-Tau Results for All Stations - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
LKWF1	Significantly increasing trend	1378722	37	1984 - 2025	26.5	0.43	25.67	0.03	0

At twenty-one program locations, monthly average water temperature increased between 0.03 and 0.09°C per year. No detectable change in monthly average water temperature was observed at one location.




Figure 12: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature on Coral Reefs in the Florida Keys - 986




Figure 13: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 7: Seasonal Kendall-Tau Results for All Stations - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
1	Significantly increasing trend	73700	11	2013 - 2023	26.50	0.23	26.62	0.06	0.0018
2	Significantly increasing trend	68589	11	2013 - 2023	26.74	0.32	26.65	0.06	0
3	Significantly increasing trend	69054	11	2013 - 2023	26.69	0.32	26.51	0.07	0
4	Significantly increasing trend	77960	11	2013 - 2023	26.69	0.28	26.36	0.06	1e-04
5	Significantly increasing trend	60729	11	2013 - 2023	26.69	0.28	26.33	0.06	1e-04
84	Significantly increasing trend	117729	17	2007 - 2023	26.37	0.40	25.91	0.07	0
85	Significantly increasing trend	118414	17	2007 - 2023	26.29	0.38	26.03	0.07	0
86	Significantly increasing trend	113127	17	2007 - 2023	26.20	0.39	25.80	0.07	0
87	Significantly increasing trend	117324	17	2007 - 2023	26.50	0.37	25.81	0.07	0
88	Significantly increasing trend	123687	17	2007 - 2023	26.42	0.38	25.97	0.07	0
89	Significantly increasing trend	120476	17	2007 - 2023	26.32	0.38	26.06	0.07	0
90	Significantly increasing trend	103477	17	2007 - 2023	26.48	0.39	25.91	0.07	0
91	Significantly increasing trend	111048	17	2007 - 2023	26.60	0.38	25.93	0.08	0
92	Significantly increasing trend	120189	17	2007 - 2023	26.50	0.40	25.90	0.08	0
93	Significantly increasing trend	115688	17	2007 - 2023	26.56	0.38	26.04	0.06	0
94	Significantly increasing trend	98419	17	2007 - 2023	25.62	0.34	24.82	0.08	0
95	Significantly increasing trend	109401	17	2007 - 2023	25.62	0.36	24.85	0.08	0
96	No significant trend	25550	6	2007 - 2012	24.87	-0.25	24.98	-0.23	0.0801
97	Significantly increasing trend	105832	14	2010 - 2023	26.50	0.42	26.21	0.08	0
98	Significantly increasing trend	94894	14	2010 - 2023	26.45	0.42	26.18	0.09	0
6	Significantly increasing trend	69753	11	2013 - 2023	26.84	0.19	26.41	0.05	0.0078

At twenty-one program locations, monthly average water temperature increased between 0.03 and 0.09°C per year. No detectable change in monthly average water temperature was observed at one location.




Figure 14: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 15: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 8: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	1928	29	1972 - 2025	27	0.28685	24.55026	0.08125	0

Monthly average water temperature increased by 0.08°C per year.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 17: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	376	12	1972 - 2025	7.99	0.06508	7.63924	0	1

pH showed no detectable trend between 1972 and 2025.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Discrete




Figure 19: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	18027	10	2011 - 2025	0.3	-0.11378	0.5434	-0.0102	0.2184

Turbidity showed no detectable trend between 2011 and 2025.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 21: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 11: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	17471	9	2017 - 2025	3.47	-	-	-	-

There was insufficient data to fit a model for total suspended solids.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 23: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 12: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	328	5	2009 - 2023	0.3863	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, uncorrected for pheophytin.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 25: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	12209	8	2013 - 2025	0.47	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, corrected for pheophytin.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 27: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 14: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	9089	16	1997 - 2025	-6.1	-0.21356	-1.47039	-0.22398	0.0014

Monthly average Secchi depth became deeper by 0.22 m per year, indicating an increase in water clarity.




Figure 28: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 29: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	1	1	2023 - 2023	7.5	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 30: Map showing location of discrete water quality sampling locations within the boundaries of *Kristin Jacobs Coral Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.