

Lemon Bay Aquatic Preserve
SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Discrete	6
Dissolved Oxygen Saturation - Discrete	8
Salinity - Discrete	10
Water Temperature - Discrete	12
pH - Discrete	14
Water Clarity	16
Turbidity - Discrete	16
Total Suspended Solids - Discrete	18
Chlorophyll a, Uncorrected for Pheophytin - Discrete	20
Chlorophyll a, Corrected for Pheophytin - Discrete	22
Secchi Depth - Discrete	24
Colored Dissolved Organic Matter - Discrete	26

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	2795	30	1995 - 2025	0.565	0.05136	0.56186	0.00111	0.1753

Total nitrogen showed no detectable trend between 1995 and 2025.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	2844	30	1995 - 2025	0.092	-0.52344	0.19036	-0.00482	0

Monthly average total phosphorus decreased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	12522	38	1971 - 2025	6.1	-0.04961	6.27021	-0.00616	0.1375

Dissolved oxygen showed no detectable trend between 1971 and 2025.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	4703	28	1998 - 2025	87	-0.08763	86.81334	-0.16667	0.0286

Monthly average dissolved oxygen saturation decreased by 0.17% per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 9: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	Significantly decreasing trend	11490	58	1954 - 2025	34	-0.08886	34.64345	-0.01673	0.0025

Monthly average salinity decreased by 0.02 ppt per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 11: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	13715	59	1954 - 2025	26.4	0.12076	24.56355	0.01521	1e-04

Monthly average water temperature increased by 0.02°C per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 13: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 7: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	10950	38	1955 - 2025	8	-0.11971	8.11393	-0.0024	7e-04

Monthly average pH decreased by less than 0.01 pH units per year.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Discrete




Figure 15: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 8: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	5742	31	1995 - 2025	2.6	0.081	2.85169	0.01599	0.0269

Monthly average turbidity increased by 0.02 NTU per year, indicating a decrease in water clarity.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 17: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	936	22	1995 - 2025	11	0.38274	3.86373	0.5182	0

Monthly average total suspended solids increased by 0.52 mg/L per year, indicating a decrease in water clarity.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 19: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	795	20	1998 - 2025	4.1	-0.09302	4.81915	-0.04028	0.1066

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 1998 and 2025.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 21: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 11: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	1102	23	1998 - 2025	4.635	-0.17256	6.78864	-0.0857	2e-04

Monthly average chlorophyll a, corrected for pheophytin, decreased by 0.09 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 23: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 12: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	1645	31	1995 - 2025	-1.3	-0.33775	-0.92498	-0.027	0

Monthly average Secchi depth became deeper by 0.02 m per year, indicating an increase in water clarity.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 25: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	573	9	2017 - 2025	10	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *Lemon Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.