

Nature Coast Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	3153	12	2005 - 2025	0.39	0.12674	0.34664	0.00226	0.169

Total nitrogen showed no detectable trend between 2005 and 2025.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	3150	12	2005 - 2025	0.009	-0.18859	0.01589	-0.00016	0.0712

Total phosphorus showed no detectable trend between 2005 and 2025.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
BBSHS	Insufficient data to calculate trend	6555	1	2016 - 2016	5.7	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	6422	36	1982 - 2025	7.33	-0.04857	7.8147	-0.0063	0.2084

Dissolved oxygen showed no detectable trend between 1982 and 2025.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
BBSHS	Insufficient data to calculate trend	6555	1	2016 - 2016	82.6	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Insufficient data to calculate trend	3260	9	2014 - 2025	100.4	-	-	-	-

There was insufficient data to fit a model for dissolved oxygen saturation.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	Significantly increasing trend	11150	41	1975 - 2025	24.6	0.08582	22.14096	0.05275	0.0205

Monthly average salinity increased by 0.05 ppt per year.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
BBSHS	Significantly decreasing trend	235670	12	2004 - 2016	18.4	-0.22	21.33	-0.38	0.005

At one program location, monthly average salinity decreased by 0.38 ppt per year.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02310674	Significantly decreasing trend	9951	21	2005 - 2025	25.5	-0.17	25.22	-0.05	4e-04
02310712	Significantly decreasing trend	7075	16	2006 - 2025	25.4	-0.12	25.03	-0.03	0.0445
283925082441201	Insufficient data to calculate trend	793	3	2024 - 2026	21.6	-	-	-	-
284506082435801	Insufficient data to calculate trend	848	3	2024 - 2026	22.1	-	-	-	-
BBSHS	No significant trend	244110	12	2004 - 2016	24.5	0.05	23.77	0.03	0.5239

At two program locations, monthly average water temperature decreased by 0.03°C per year at one site and by 0.05°C per year at the other. No detectable change in monthly average water temperature was observed at one location. There was insufficient data to fit a model for two locations.




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	6513	41	1974 - 2025	23.9	0.12115	22.14171	0.02832	0.0019

Monthly average water temperature increased by 0.03°C per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
BBSHS	Insufficient data to calculate trend	6555	1	2016 - 2016	8	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	5677	35	1991 - 2025	8.15	-0.03259	8.13897	-0.00084	0.3881

pH showed no detectable trend between 1991 and 2025.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Table 13: Seasonal Kendall-Tau Results - Specific Conductivity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02310674	Significantly decreasing trend	12487	20	2006 - 2025	23.80	-0.37	28.46	-0.57	0
02310712	Significantly decreasing trend	10064	16	2006 - 2025	32.60	-0.38	36.1	-0.53	0
283925082441201	Insufficient data to calculate trend	1391	3	2024 - 2026	44.70	-	-	-	-
284506082435801	Insufficient data to calculate trend	1414	3	2024 - 2026	42.30	-	-	-	-
BBSHS	No significant trend	244107	12	2004 - 2016	29.73	-0.13	31.53	-0.33	0.0716

At two program locations, monthly average specific conductivity decreased by 0.53 ms/cm per year at one site and by 0.57 ms/cm per year at the other. No detectable change in monthly average specific conductivity was observed at one location. There was insufficient data to fit a model for two locations.




Figure 25: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
BBSHS	Insufficient data to calculate trend	6368	1	2016 - 2016	2	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 28: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1683	28	1995 - 2024	0.85	0.03268	0.81866	0.00286	0.4051

Turbidity showed no detectable trend between 1995 and 2024.




Figure 29: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 30: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1162	10	2015 - 2024	3.41	0.2236	3.70853	0.15299	0.0986

Total suspended solids showed no detectable trend between 2015 and 2024.




Figure 31: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 32: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	2144	12	2005 - 2025	1	-0.21712	1.73813	-0.02759	0.0346

Monthly average chlorophyll a, uncorrected for pheophytin, decreased by 0.03 µg/L per year, indicating an increase in water clarity.




Figure 33: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 34: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	3093	11	2015 - 2025	1	-0.21302	1.77462	-0.02565	0.0115

Monthly average chlorophyll a, corrected for pheophytin, decreased by 0.03 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 35: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 36: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	3814	18	2003 - 2025	-1.6	-0.34931	-1.02368	-0.03728	0

Monthly average Secchi depth became deeper by 0.03 m per year, indicating an increase in water clarity.




Figure 37: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 38: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	4387	10	2016 - 2025	10.543	-0.08039	12.96262	-0.10402	0.3114

Colored dissolved organic matter showed no detectable trend between 2016 and 2025.




Figure 39: Map showing location of discrete water quality sampling locations within the boundaries of *Nature Coast Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.