

Pine Island Sound Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Discrete	10
Salinity - Discrete	12
Salinity - Continuous	14
Water Temperature - Continuous	16
Water Temperature - Discrete	18
pH - Continuous	20
pH - Discrete	22
Water Clarity	24
Turbidity - Continuous	24
Turbidity - Discrete	26
Total Suspended Solids - Discrete	28
Chlorophyll a, Uncorrected for Pheophytin - Discrete	30
Chlorophyll a, Corrected for Pheophytin - Discrete	32
Secchi Depth - Discrete	34
Colored Dissolved Organic Matter - Discrete	36

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	6243	31	1995 - 2025	0.622	0.37732	0.37067	0.01533	0

Monthly average total nitrogen increased by 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	6064	34	1987 - 2025	0.03583	-0.19509	0.04934	-0.00058	0

Monthly average total phosphorus decreased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293249	Insufficient data to calculate trend	1302	4	2010 - 2013	8.8	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	36250	39	1985 - 2025	6.9	-0.01859	7.24679	-0.00192	0.568

Dissolved oxygen showed no detectable trend between 1985 and 2025.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	3650	12	2014 - 2025	91.08864	-0.04453	90.09168	-0.0873	0.5424

Dissolved oxygen saturation showed no detectable trend between 2014 and 2025.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 11: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	35254	64	1954 - 2025	32.8	-0.05259	33.33206	-0.01031	0.0909

Salinity showed no detectable trend between 1954 and 2025.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 13: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 7: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293249	Insufficient data to calculate trend	1871	4	2010 - 2013	32	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 14: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 15: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293249	Insufficient data to calculate trend	2445	4	2010 - 2013	28.1	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 16: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 17: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	38296	62	1954 - 2025	26.4	0.15614	24.23034	0.01608	0

Monthly average water temperature increased by 0.02°C per year.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 19: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 10: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293249	Insufficient data to calculate trend	1164	4	2010 - 2013	8.3	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 20: Map showing location of pH continuous water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 21: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 11: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	33845	39	1955 - 2025	8.1	-0.00848	8.11732	-0.00008	0.8038

pH showed no detectable trend between 1955 and 2025.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 23: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 12: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
02293249	Insufficient data to calculate trend	1174	4	2010 - 2013	5.4	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 24: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 25: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	6711	31	1995 - 2025	2	-0.02912	2.56474	-0.0068	0.4367

Turbidity showed no detectable trend between 1995 and 2025.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 27: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 14: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	2835	29	1987 - 2025	11.9	-0.0792	19.05064	-0.13045	0.0674

Total suspended solids showed no detectable trend between 1987 and 2025.




Figure 28: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 29: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1254	22	1998 - 2025	4.3	0.01359	4.68341	0.00556	0.8137

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 1998 and 2025.




Figure 30: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 31: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	4034	22	2001 - 2025	3.9	-0.09516	6.05682	-0.05014	0.0491

Monthly average chlorophyll a, corrected for pheophytin, decreased by 0.05 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 32: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 33: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 17: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	17340	32	1994 - 2025	-1	-0.12597	-1.11327	-0.00419	6e-04

Monthly average Secchi depth became deeper by less than 0.01 m per year, indicating an increase in water clarity.




Figure 34: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 35: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	1525	9	2017 - 2025	7.44	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 36: Map showing location of discrete water quality sampling locations within the boundaries of *Pine Island Sound Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.