

Rocky Bayou State Park Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Discrete	6
Dissolved Oxygen Saturation - Discrete	8
Salinity - Discrete	10
Water Temperature - Discrete	12
pH - Discrete	14
Water Clarity	16
Turbidity - Discrete	16
Total Suspended Solids - Discrete	18
Chlorophyll a, Uncorrected for Pheophytin - Discrete	20
Chlorophyll a, Corrected for Pheophytin - Discrete	22
Secchi Depth - Discrete	24
Colored Dissolved Organic Matter - Discrete	26

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	641	24	2001 - 2024	0.2	0.17718	0.14513	0.0023	0

Monthly average total nitrogen increased by less than 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	464	25	2001 - 2025	0.007	0.28558	0.00336	0.00022	0

Monthly average total phosphorus increased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	746	27	1994 - 2025	6.65	0.16132	5.47208	0.042	2e-04

Monthly average dissolved oxygen increased by 0.04 mg/L per year.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	653	26	2000 - 2025	89.2	0.10999	71.25729	0.375	0.0216

Monthly average dissolved oxygen saturation increased by 0.38% per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 9: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	758	28	1994 - 2025	21.59	-0.01919	22.90342	-0.0275	0.6273

Salinity showed no detectable trend between 1994 and 2025.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 11: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	512	20	1994 - 2025	25.0278	0.16244	22.90553	0.07135	0.0029

Monthly average water temperature increased by 0.07°C per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 13: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 7: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	770	28	1994 - 2025	7.9	0.03835	7.84565	0.0025	0.3813

pH showed no detectable trend between 1994 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Discrete




Figure 15: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 8: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	222	11	2008 - 2018	1	-0.2377	3.44167	-0.21339	0.0034

Monthly average turbidity decreased by 0.21 NTU per year, indicating an increase in water clarity.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 17: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	17	4	2013 - 2017	3	-	-	-	-

There was insufficient data to fit a model for total suspended solids.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 19: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	267	25	2001 - 2025	4	-0.00571	3.79167	0	1

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 2001 and 2025.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 21: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 11: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	115	14	2012 - 2025	3	-0.17758	3.21845	-0.01429	0.033

Monthly average chlorophyll a, corrected for pheophytin, decreased by 0.01 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 23: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 12: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	413	24	1995 - 2025	-2.4384	0.11864	-2.7188	0.01793	0.0257

Monthly average Secchi depth became shallower by 0.02 m per year, indicating a decrease in water clarity.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 25: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	21	7	2019 - 2025	13	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *Rocky Bayou State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.