

Terra Ceia Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	2444	31	1995 - 2025	0.3765	-0.24302	0.57194	-0.00655	0

Monthly average total nitrogen decreased by 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	3277	31	1995 - 2025	0.106	-0.10809	0.15264	-0.00106	0.0037

Monthly average total phosphorus decreased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data to calculate trend	8153	2	2008 - 2009	4.2	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	25110	37	1989 - 2025	6.52	-0.0277	6.649	-0.00282	0.4223

Dissolved oxygen showed no detectable trend between 1989 and 2025.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data to calculate trend	8153	2	2008 - 2009	59.5	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	5224	33	1993 - 2025	90.3	0.09554	86.30112	0.09583	0.1081

Dissolved oxygen saturation showed no detectable trend between 1993 and 2025.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	25524	49	1966 - 2025	30.62	-0.04698	30.98007	-0.01303	0.1548

Salinity showed no detectable trend between 1966 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data to calculate trend	645	2	2006 - 2007	12.0	-	-	-	-
02300084	Insufficient data to calculate trend	696	2	2006 - 2007	11.0	-	-	-	-
023000842	Insufficient data to calculate trend	578	2	2006 - 2007	28.0	-	-	-	-
TCBH	Insufficient data to calculate trend	8304	2	2008 - 2009	32.6	-	-	-	-

There was insufficient data to fit a model for four locations.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data to calculate trend	659	2	2006 - 2007	26.6	-	-	-	-
02300084	Insufficient data to calculate trend	717	2	2006 - 2007	27.5	-	-	-	-
023000842	Insufficient data to calculate trend	571	2	2006 - 2007	27.7	-	-	-	-
TCBH	Insufficient data to calculate trend	8305	2	2008 - 2009	20.0	-	-	-	-

There was insufficient data to fit a model for four locations.




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	26272	51	1966 - 2025	26.7	0.0907	24.99941	0.01344	0.0077

Monthly average water temperature increased by 0.01°C per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data to calculate trend	8306	2	2008 - 2009	8.1	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	23347	37	1989 - 2025	8	-0.09489	8.0431	-0.00161	0.0085

Monthly average pH decreased by less than 0.01 pH units per year.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Table 13: Seasonal Kendall-Tau Results - Specific Conductivity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data to calculate trend	710	2	2006 - 2007	18.60	-	-	-	-
02300084	Insufficient data to calculate trend	775	2	2006 - 2007	17.00	-	-	-	-
023000842	Insufficient data to calculate trend	669	2	2006 - 2007	43.10	-	-	-	-
TCBH	Insufficient data to calculate trend	8305	2	2008 - 2009	49.71	-	-	-	-

There was insufficient data to fit a model for four locations.




Figure 25: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data to calculate trend	8263	2	2008 - 2009	2	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 28: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	8131	31	1995 - 2025	1.9	0.00276	2.15566	0.00047	0.9288

Turbidity showed no detectable trend between 1995 and 2025.




Figure 29: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 30: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	1723	31	1995 - 2025	10.7	-0.4038	19.08939	-0.58483	0

Monthly average total suspended solids decreased by 0.58 mg/L per year, indicating an increase in water clarity.




Figure 31: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 32: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	1093	25	1999 - 2025	3.2	-0.09334	3.46386	-0.02291	0.029

Monthly average chlorophyll a, uncorrected for pheophytin, decreased by 0.02 $\mu\text{g/L}$ per year, indicating an increase in water clarity.




Figure 33: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 34: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	54	6	2000 - 2022	5.1	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, corrected for pheophytin.




Figure 35: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 36: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	8222	31	1995 - 2025	-0.9	-0.32258	-0.84727	-0.01727	0

Monthly average Secchi depth became deeper by 0.02 m per year, indicating an increase in water clarity.




Figure 37: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 38: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	679	8	2018 - 2025	5.8	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 39: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.