

Tomoka Marsh Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 05 May, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	417	22	1997 - 2025	0.7879	-0.21496	1.03206	-0.00988	3e-04

Monthly average total nitrogen decreased by 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	775	26	1997 - 2025	0.0962	0.0507	0.0979	0.00015	0.7126

Total phosphorus showed no detectable trend between 1997 and 2025.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	40492	2	2021 - 2022	6.5	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	759	26	1997 - 2025	5.41	0.02125	4.97338	-0.00176	0.7954

Dissolved oxygen showed no detectable trend between 1997 and 2025.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	40492	2	2021 - 2022	88.4	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	464	12	2009 - 2025	71.4	0.15469	68.52066	0.68764	0.0231

Monthly average dissolved oxygen saturation increased by 0.69% per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	752	25	1998 - 2025	15.915	-0.00747	14.54189	0.00278	1

Salinity showed no detectable trend between 1998 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	23.3	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	25.5	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	775	26	1997 - 2025	25	0.14464	24.44885	0.04286	0.0092

Monthly average water temperature increased by 0.04°C per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	7.8	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	748	26	1997 - 2025	7.54	0.10928	7.40872	0.00417	0.1062

pH showed no detectable trend between 1997 and 2025.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Table 13: Seasonal Kendall-Tau Results - Specific Conductivity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	36.84	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 25: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41169	2	2021 - 2022	7	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 28: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	533	26	1997 - 2025	7.12	0.04555	7.19532	0.03382	0.3347

Turbidity showed no detectable trend between 1997 and 2025.




Figure 29: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 30: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	486	26	1997 - 2025	14	-0.07451	14.99022	-0.20833	0.0328

Monthly average total suspended solids decreased by 0.21 mg/L per year, indicating an increase in water clarity.




Figure 31: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 32: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	351	9	2000 - 2025	7.94587	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, uncorrected for pheophytin.




Figure 33: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 34: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	324	9	2002 - 2025	6.31455	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, corrected for pheophytin.




Figure 35: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 36: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Insufficient data to calculate trend	85	9	2000 - 2023	-0.60961	-	-	-	-

Secchi depth showed no detectable trend between 2000 and 2025.




Figure 37: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 38: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	293	6	2020 - 2025	60.5168	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 39: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.