

Fort Clinch State Park Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 09 April, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Discrete	6
Dissolved Oxygen Saturation - Discrete	8
Salinity - Discrete	10
Water Temperature - Discrete	12
pH - Discrete	14
Water Clarity	16
Turbidity - Discrete	16
Total Suspended Solids - Discrete	18
Chlorophyll a, Uncorrected for Pheophytin - Discrete	20
Chlorophyll a, Corrected for Pheophytin - Discrete	22
Secchi Depth - Discrete	24

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	56	13	2000 - 2025	0.3545	-0.29922	0.50547	-0.00743	0.0562

Total nitrogen showed no detectable trend between 2000 and 2025.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	58	13	2000 - 2025	0.035	-0.13333	0.05592	-0.00058	0.4608

Total phosphorus showed no detectable trend between 2000 and 2025.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	1719	27	1994 - 2025	7.1	-0.23003	7.80625	-0.0375	0

Monthly average dissolved oxygen decreased by 0.04 mg/L per year.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Insufficient data to calculate trend	48	5	2000 - 2014	85.45	-	-	-	-

There was insufficient data to fit a model for dissolved oxygen saturation.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 9: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	1725	27	1994 - 2024	33.5	-0.00903	33.32809	-0.0056	0.7904

Salinity showed no detectable trend between 1994 and 2024.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 11: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	1741	27	1994 - 2025	22.9	0.11667	21.48022	0.04439	0.0079

Monthly average water temperature increased by 0.04°C per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 13: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 7: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	1720	27	1994 - 2025	7.8	0.0731	7.79625	0.00248	0.1211

pH showed no detectable trend between 1994 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Discrete




Figure 15: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 8: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	52	11	2000 - 2025	5.2	-0.08943	10.71538	-0.06637	0.9216

Turbidity showed no detectable trend between 2000 and 2025.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 17: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	28	4	2004 - 2014	21.15	-	-	-	-

There was insufficient data to fit a model for total suspended solids.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 19: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	34	9	2004 - 2025	4.55	-	-	-	-

There was insufficient data to fit a model for chlorophyll a, uncorrected for pheophytin.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 21: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 11: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	52	10	2004 - 2025	4.15	-0.01667	3.99773	0.05	0.8428

Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2004 and 2025.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 23: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 12: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	1639	25	2000 - 2024	-0.9	0.05219	-0.84562	0.00444	0.2346

Secchi depth showed no detectable trend between 2000 and 2024.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Fort Clinch State Park Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.