

Rookery Bay Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 09 April, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	2640	30	1989 - 2025	0.26	0.1689	0.20291	0.00472	0.0002

Monthly average total nitrogen increased by less than 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	2270	33	1989 - 2025	0.034	-0.03641	0.04007	-0.00013	0.2966

Total phosphorus showed no detectable trend between 1989 and 2025.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10441	2	1997 - 1998	7.2	-	-	-	-
rkblhwq	No significant trend	613216	23	2001 - 2026	5.6	0.03	5.46	0	0.5182

No detectable change in monthly average dissolved oxygen was observed at one location. There was insufficient data to fit a model for one location.




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	3909	34	1989 - 2025	5.9	-0.09659	6.17588	-0.01511	0.0161

Monthly average dissolved oxygen decreased by 0.02 mg/L per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10441	2	1997 - 1998	102.4	-	-	-	-
rkblhwq	No significant trend	625534	23	2001 - 2026	82.1	0.07	80.86	0.07	0.1302

No detectable change in monthly average dissolved oxygen saturation was observed at one location. There was insufficient data to fit a model for one location.




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	197	19	1998 - 2025	86.1	-0.07319	95.01341	-0.30873	0.4526

Dissolved oxygen saturation showed no detectable trend between 1998 and 2025.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	5327	46	1954 - 2025	34.14	0.02259	34.05806	0.00467	0.5218

Salinity showed no detectable trend between 1954 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12256	2	1997 - 1998	34.5	-	-	-	-
rkblhwq	Significantly decreasing trend	700367	26	2001 - 2026	33.2	-0.13	34.36	-0.08	0.0023

At one program location, monthly average salinity decreased by 0.08 ppt per year. There was insufficient data to fit a model for one location.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12610	2	1997 - 1998	23.8	-	-	-	-
rkblhwq	Significantly increasing trend	731519	26	2001 - 2026	26.9	0.24	26.16	0.05	0

At one program location, monthly average water temperature increased by 0.05°C per year. There was insufficient data to fit a model for one location.




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	4571	45	1954 - 2025	26.22	0.0837	25.15326	0.01114	0.0137

Monthly average water temperature increased by 0.01°C per year.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12610	2	1997 - 1998	8.0	-	-	-	-
rkblhwq	Significantly increasing trend	672354	26	2001 - 2026	7.9	0.15	7.83	0	0.0003

At one program location, monthly average pH increased by less than 0.01 pH units per year. There was insufficient data to fit a model for one location.




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	1887	38	1955 - 2025	7.9	-0.13597	8.06668	-0.00354	0.0004

Monthly average pH decreased by less than 0.01 pH units per year.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Table 13: Seasonal Kendall-Tau Results - Specific Conductivity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	12256	2	1997 - 1998	52.41	-	-	-	-
rkblhwq	Significantly decreasing trend	700390	26	2001 - 2026	50.59	-0.12	52.11	-0.1	0.003




Figure 25: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbbmwq	Insufficient data to calculate trend	10654	2	1997 - 1998	11	-	-	-	-
rkblhwq	Significantly decreasing trend	647527	26	2001 - 2026	8	-0.28	12.22	-0.19	0

At one program location, monthly average turbidity decreased by 0.19 NTU per year. There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 28: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	1879	32	1989 - 2025	5	0.23872	3.58473	0.10936	0

Monthly average turbidity increased by 0.11 NTU per year, indicating a decrease in water clarity.




Figure 29: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 30: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	260	24	1989 - 2025	8	0.13598	2.6125	0.12	0.0239

Monthly average total suspended solids increased by 0.12 mg/L per year, indicating a decrease in water clarity.




Figure 31: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 32: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	3290	27	1999 - 2025	5	0.17531	4.37536	0.05756	0.0001

Monthly average chlorophyll a, uncorrected for pheophytin, increased by 0.06 $\mu\text{g/L}$ per year, indicating a decrease in water clarity.




Figure 33: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 34: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1713	16	2002 - 2025	4.3	0.09735	3.12217	0.04914	0.1062

Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2002 and 2025.




Figure 35: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 36: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	434	22	1998 - 2025	-1.2	0.10906	-1.16309	0.01134	0.1121

Secchi depth showed no detectable trend between 1998 and 2025.




Figure 37: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 38: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	1	1	2021 - 2021	110	-	-	-	-

There was insufficient data to fit a model for colored dissolved organic matter.




Figure 39: Map showing location of discrete water quality sampling locations within the boundaries of *Rookery Bay Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.