

Tomoka Marsh Aquatic Preserve

SEACAR Habitat Analyses

Last compiled on 10 June, 2026

Contents

Funding & Acknowledgements	2
Threshold Filtering	2
Value Qualifiers	3
Water Column	5
Seasonal Kendall-Tau Analysis	5
Water Quality - Discrete	5
Dissolved Oxygen - Discrete	6
Dissolved Oxygen Saturation - Discrete	7
pH - Discrete	9
Salinity - Discrete	11
Total Nitrogen - Discrete	13
Total Phosphorus - Discrete	16
Turbidity - Discrete	18
Water Temperature - Discrete	20
Water Quality - Continuous	23
pH - Continuous	25
Dissolved Oxygen Saturation - Continuous	26
Dissolved Oxygen - Continuous	27
Turbidity - Continuous	28
Water Temperature - Continuous	29
Salinity - Continuous	30
Specific Conductivity - Continuous	31
Oyster	31
Density	33
Natural	33
Percent Live	34
Natural	35
Shell Height	36
Natural	36
References	38

Funding & Acknowledgements

The data used in this analysis is from the Export Standardized Tables in the SEACAR Data Discovery Interface (DDI). Documents and information available through the SEACAR DDI are owned by the data provider(s) and users are expected to provide appropriate credit following accepted citation formats. Users are encouraged to access data to maximize utilization of gained knowledge, reducing redundant research and facilitating partnerships and scientific innovation.

With respect to documents and information available from SEACAR DDI, neither the State of Florida nor the Florida Department of Environmental Protection makes any warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose arising out of the use or inability to use the data, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

This report was funded in part, through a grant agreement from the Florida Department of Environmental Protection, Florida Coastal Management Program, by a grant provided by the Office for Coastal Management under the Coastal Zone Management Act of 1972, as amended, National Oceanic and Atmospheric Administration. The views, statements, findings, conclusions and recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of the State of Florida, NOAA or any of their sub agencies.

Published: 2026-06-10



Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection's (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR_QAQCFlagCode* and *SEACAR_QAQC_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Dissolved Oxygen	mg/L	-0.000001	50
Dissolved Oxygen Saturation	%	-0.000001	500
Fluorescent Dissolved Organic Matter	QSE	-	-
Salinity	ppt	-0.000001	70
Specific Conductivity	mS/cm	-0.000001	200
Turbidity	NTU	-0.000001	4000
Water Temperature	Degrees C	-5.000000	45
pH	None	2.000000	14

Table 2: Discrete Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Ammonia, Un-ionized (NH3)	mg/L	-	-
Ammonium (NH4)	mg/L	-	-

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Chlorophyll a, Corrected for Pheophytin	ug/L	-	-
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Colored Dissolved Organic Matter	PCU	-	-
Dissolved Oxygen	mg/L	-0.000001	25
Dissolved Oxygen Saturation	%	-0.000001	310
Fluorescent Dissolved Organic Matter	QSE	-	-
Light Extinction Coefficient	m ⁻¹	-	-
NO2+3, Filtered	mg/L	-	-
Nitrate (NO3)	mg/L	-	-
Nitrite (NO2)	mg/L	-	-
Nitrogen, inorganic	mg/L	-	-
Nitrogen, organic	mg/L	-	-
Phosphate, Filtered (PO4)	mg/L	-	-
Salinity	ppt	-0.000001	70
Secchi Depth	m	0.000001	50
Specific Conductivity	mS/cm	0.005000	100
Total Ammonia (N)	mg/L	-	-
Total Kjeldahl Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Phosphorus	mg/L	-	-
Total Suspended Solids	mg/L	-	-
Turbidity	NTU	-	-
Water Temperature	Degrees C	3.000000	40
pH	None	2.000000	13

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 and 2

<i>SEACAR QAQC Description</i>	<i>Include</i>	<i>SEACAR QAQCFlagCode</i>
Exceeds maximum threshold	0	2Q
Below minimum threshold	0	4Q
Within threshold tolerance	1	6Q
No defined thresholds for this parameter	1	7Q

Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>MDL</i>	<i>Description</i>
STORET-WIN	H	0	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	0	0	Estimated value
STORET-WIN	V	0	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	0	0	Lab analysis from an improperly preserved sample; data may be inaccurate

Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

H - Value based on field kit determination; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

I - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

Q - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

S - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

U - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported

Systemwide Monitoring Program (SWMP) value qualifier codes

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	1	Optional parameter not collected
SWMP	-2	0	Missing data
SWMP	-3	0	Data rejected due to QA/QC
SWMP	-4	0	Outside low sensor range
SWMP	-5	0	Outside high sensor range
SWMP	0	1	Passed initial QA/QC checks
SWMP	1	0	Suspect data
SWMP	2	1	Reserved for future use
SWMP	3	1	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	1	Historical: Pre-auto QA/QC
SWMP	5	1	Corrected data

Water Column

The water column habitat extends from the water's surface to the bottom sediments, and it's where fish, dolphins, crabs and people swim! So much life makes its home in the water column that the health of marine and coastal ecosystems, as well as human economies, depend on the condition of this vulnerable habitat. Local patterns of rainfall, temperature, winds and currents can rapidly change the condition of the water column, while global influences such as [El Niño/La Niña](#), large-scale fluctuation in sea temperatures and climate change can have long-term effects. Inputs from the prosperity of our day-to-day lives including farming, mining and forestry, and emissions from power generation, automobiles and water treatment can also alter the health of the water column. Acting alone or together, each input can have complex and lasting effects on habitats and ecosystems.

SEACAR evaluates water column health with several essential parameters. These include nutrient surveys of nitrogen and phosphorus, and water quality assessments of salinity, dissolved oxygen, pH, and water temperature. Water clarity is evaluated with Secchi depth, turbidity, levels of chlorophyll a, total suspended solids, and colored dissolved organic matter. Additionally, the richness of nekton is indicated by the abundance of free-swimming fishes and macroinvertebrates like crabs and shrimps.

Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as "insufficient data to conduct analysis". Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use_In_Analysis* = **TRUE**. Any that fail either test will be excluded from the analyses and labeled as *Use_In_Analysis* = **FALSE**. The points for all Water Column plots displayed in this section are monthly averages. Trend significance will be denoted as "Significant Trend" (when $p < 0.05$), or "Non-significant Trend" (when $p \geq 0.05$). Any parameters with insufficient data to perform Seasonal Kendall-Tau test will have their monthly averages plotted without a corresponding trend line.

Water Quality - Discrete

The following files were used in the discrete analysis:

- *Combined_WQ_WC_NUT_Chlorophyll_a_corrected_for_pheophytin-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Chlorophyll_a_uncorrected_for_pheophytin-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Colored_dissolved_organic_matter_CDOM-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen_Saturation-2026-May-18.txt*
- *Combined_WQ_WC_NUT_pH-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Salinity-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Secchi_Depth-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Total_Nitrogen-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Total_Phosphorus-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Total_Suspended_Solids_TSS-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Turbidity-2026-May-18.txt*
- *Combined_WQ_WC_NUT_Water_Temperature-2026-May-18.txt*

Dissolved Oxygen - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 1: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No detectable trend	759	26	1997 - 2025	5.41	0.0213	4.9734	-0.0018	0.7954

Dissolved oxygen showed no detectable trend between 1997 and 2025.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 7: Programs contributing data for Dissolved Oxygen

ProgramID	N_Data	YearMin	YearMax
5002	745	1997	2025
95	24	2017	2017
69	4	2001	2004

Program names:

5002 - Florida STORET / WIN¹

95 - Harmful Algal Bloom Marine Observation Network²

69 - Fisheries-Independent Monitoring (FIM) Program³

Dissolved Oxygen Saturation - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 3: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 8: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen Saturation

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Increasing trend	464	12	2009 - 2025	71.4	0.1547	68.5207	0.6876	0.0231

Monthly average dissolved oxygen saturation increased by 0.69% per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 9: Programs contributing data for Dissolved Oxygen Saturation

ProgramID	N_Data	YearMin	YearMax
5002	466	2009	2025

Program names:

5002 - Florida STORET / WIN¹

pH - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 5: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Trend Analysis for pH

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No detectable trend	748	26	1997 - 2025	7.54	0.1093	7.4087	0.0042	0.1062

pH showed no detectable trend between 1997 and 2025.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 11: Programs contributing data for pH

ProgramID	N_Data	YearMin	YearMax
5002	745	1997	2025
69	4	2001	2004

Program names:

5002 - Florida STORET / WIN¹

69 - Fisheries-Independent Monitoring (FIM) Program³

Salinity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 7: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 12: Seasonal Kendall-Tau Trend Analysis for Salinity

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No detectable trend	752	25	1998 - 2025	15.915	-0.0075	14.5419	0.0028	1

Salinity showed no detectable trend between 1998 and 2025.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 13: Programs contributing data for Salinity

ProgramID	N_Data	YearMin	YearMax
5002	725	1998	2025
95	24	2017	2017
69	4	2001	2004

Program names:

5002 - Florida STORET / WIN¹

95 - Harmful Algal Bloom Marine Observation Network²

69 - Fisheries-Independent Monitoring (FIM) Program³

Total Nitrogen - Discrete

Total Nitrogen Calculation:

The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1) $TN = TKN + NO3O2$;
- 2) $TN = TKN + NO3 + NO2$;
- 3) $TN = ORGN + NH4 + NO3O2$;
- 4) $TN = ORGN + NH4 + NO2 + NO3$;
- 5) $TN = TKN + NO3$;
- 6) $TN = ORGN + NH4 + NO3$;

Additional Information:

- Rules for use of sample fraction:
 - Florida Department of Environmental Protection (FDEP) report that if both “Total” and “Dissolved” components are reported, only “Total” is used. If the total is not reported, then the dissolved components are used as a best available replacement.
 - Total nitrogen calculations are done using nitrogen components with the same sample fraction, nitrogen components with mixed total/dissolved sample fractions are not used. In other words, total nitrogen can be calculated when TKN and NO3O2 are both total sample fractions, or when both are dissolved sample fractions. *Future calculations of total nitrogen values may be based on components with mixed sample fractions.*
- Values inserted into data:
 - ParameterName = “Total Nitrogen”
 - SEACAR_QAQCFlagCode = “1Q”
 - SEACAR_QAQC_Description = “SEACAR Calculated”

Seasonal Kendall-Tau Trend Analysis




Figure 9: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 14: Seasonal Kendall-Tau Trend Analysis for Total Nitrogen

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Decreasing trend	417	22	1997 - 2025	0.7879	-0.215	1.0321	-0.0099	0.0003

Monthly average total nitrogen decreased by 0.01 mg/L per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 15: Programs contributing data for Total Nitrogen

ProgramID	N_Data	YearMin	YearMax
5002	381	1997	2025
514	36	2000	2001

Program names:

5002 - Florida STORET / WIN¹

514 - Florida LAKEWATCH Program⁴

Total Phosphorus - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 11: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Trend Analysis for Total Phosphorus

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	775	26	1997 - 2025	0.0962	0.0507	0.0979	0.0001	0.7126

Total phosphorus showed no detectable trend between 1997 and 2025.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 17: Programs contributing data for Total Phosphorus

ProgramID	N_Data	YearMin	YearMax
5002	746	1997	2025
514	36	2000	2001

Program names:

5002 - Florida STORET / WIN¹

514 - Florida LAKEWATCH Program⁴

Turbidity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 13: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 18: Seasonal Kendall-Tau Trend Analysis for Turbidity

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	533	26	1997 - 2025	7.12	0.0456	7.1953	0.0338	0.3347

Turbidity showed no detectable trend between 1997 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 19: Programs contributing data for Turbidity

ProgramID	N_Data	YearMin	YearMax
5002	534	1997	2025

Program names:

5002 - Florida STORET / WIN¹

Water Temperature - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 15: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 20: Seasonal Kendall-Tau Trend Analysis for Water Temperature

Activity Type	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Increasing trend	775	26	1997 - 2025	25	0.1446	24.4489	0.0429	0.0092

Monthly average water temperature increased by 0.04°C per year.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 21: Programs contributing data for Water Temperature

ProgramID	N_Data	YearMin	YearMax
5002	750	1997	2025
95	24	2017	2017
69	4	2001	2004

Program names:

5002 - Florida STORET / WIN¹

95 - Harmful Algal Bloom Marine Observation Network²

69 - Fisheries-Independent Monitoring (FIM) Program³

Water Quality - Continuous

The following files were used in the continuous analysis:

- *Combined_WQ_WC_NUT_cont_Chlorophyll_a_Uncorrected_for_Pheophytin-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Dissolved_Oxygen-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Dissolved_Oxygen_Saturation-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Fluorescent_Dissolved_Organic_Matter-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_pH-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Salinity-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Specific_Conductivity-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Turbidity-2026-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Water_Temperature-2026-Mar-06.txt*

Continuous monitoring locations in Tomoka Marsh Aquatic Preserve

Table 22: Station overview for Continuous parameters by Program

<i>ProgramID</i>	<i>ProgramLocationID</i>	<i>Years of Data</i>	<i>Use in Analysis</i>	<i>Parameters</i>
10003	TMGR	2	FALSE	pH , Dissolved Oxygen Saturation, Dissolved Oxygen , Turbidity , Water Temperature , Salinity , Specific Conductivity

Program names:

[10003](#) - Tomoka Marsh Aquatic Preserve Continuous Water Quality Monitoring⁵




Figure 17: Map showing continuous water quality sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. Sites marked as *Use In Analysis* (green) are featured in this report.

pH - Continuous




Figure 18: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 23: Seasonal Kendall-Tau Results for pH - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	7.8	-	-	-	-

There was insufficient data to fit a model for one location.

Dissolved Oxygen Saturation - Continuous




Figure 19: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 24: Seasonal Kendall-Tau Results for Dissolved Oxygen Saturation - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	40492	2	2021 - 2022	88.4	-	-	-	-

There was insufficient data to fit a model for one location.

Dissolved Oxygen - Continuous




Figure 20: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 25: Seasonal Kendall-Tau Results for Dissolved Oxygen - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	40492	2	2021 - 2022	6.5	-	-	-	-

There was insufficient data to fit a model for one location.

Turbidity - Continuous




Figure 21: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 26: Seasonal Kendall-Tau Results for Turbidity - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41169	2	2021 - 2022	7	-	-	-	-

There was insufficient data to fit a model for one location.

Water Temperature - Continuous




Figure 22: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 27: Seasonal Kendall-Tau Results for Water Temperature - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	25.5	-	-	-	-

There was insufficient data to fit a model for one location.

Salinity - Continuous




Figure 23: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 28: Seasonal Kendall-Tau Results for Salinity - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	23.3	-	-	-	-

There was insufficient data to fit a model for one location.

Specific Conductivity - Continuous




Figure 24: Scatter plot of monthly average specific conductivity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 29: Seasonal Kendall-Tau Results for Specific Conductivity - All Stations

Program Location	Statistical Trend	No. of Samples	No. Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TMGR	Insufficient data to calculate trend	41374	2	2021 - 2022	36.84	-	-	-	-

There was insufficient data to fit a model for one location.

Oyster

The data file used is: **All_OYSTER_Parameters-2026-Jun-04.txt**

Tomoka Marsh Aquatic Preserve

Oyster sample locations available in SEACAR by Program and Year



Program name

- Northern Coastal Basins and Mosquito Lagoon Oyster Condition Assessment

Figure 25: Maps showing the temporal scope of oyster sampling sites within the boundaries of *Tomoka Marsh Aquatic Preserve* by program name.

Click [here](#) to view spatio-temporal plots on GitHub.

Density

For natural reefs, there was insufficient data to calculate a trend for density.

Natural




Figure 26: Scatter plot of oyster density over time. If the time series included five or more years with observations, an estimated trend (blue line) and a 95% credible interval (purple band) may also be plotted. Data points are jittered horizontally to reduce overlap.

Table 30: Model results for Oyster Density - Natural

Habitat Type	Shell Type	Statistical Trend	Period of Record	Estimate	Standard Error	Credible Interval
Natural	Live Oysters	Insufficient data	2015 - 2015	-	-	-




Figure 27: Map showing location of oyster density sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Percent Live

For natural reefs, there was insufficient data to calculate a trend for percent live cover.

Natural




Figure 28: Scatter plot of percent live oysters over time. If the time series included five or more years with observations, an estimated trend (blue line) and a 95% credible interval (purple band) may also be plotted. Estimation method is represented as percent (circles), point-intercept (triangle), or estimated percent (square), with data points jittered horizontally to reduce overlap.

Table 31: Model results for Oyster Percent Live - Natural

Habitat Type	Shell Type	Statistical Trend	Period of Record	Estimate	Standard Error	Credible Interval
Natural	Live Oysters	Insufficient data	2015 - 2015	-	-	-




Figure 29: Map showing location of oyster percent live sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Shell Height

For natural reefs, there was insufficient data to calculate a trend for percent live cover.

Natural

Table 32: Model results for Oyster Shell Height - Natural

Habitat Type	Shell Type	SizeClass	Statistical Trend	Period of Record	No. of Samples	Estimate	Standard Error	Credible Interval
--------------	------------	-----------	-------------------	------------------	----------------	----------	----------------	-------------------

Map showing location of oyster shell height sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Figure 30: Map showing location of oyster shell height sampling locations within the boundaries of *Tomoka Marsh Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

References

1. Florida Department of Environmental Protection (DEP). [Florida STORET / WIN](#). (2024).
2. Florida Fish and Wildlife Conservation Commission (FWC); Florida Fish and Wildlife Research Institute (FWRI). [Harmful Algal Bloom Marine Observation Network](#). (2018).
3. Florida Fish and Wildlife Conservation Commission (FWC). [Fisheries-Independent Monitoring \(FIM\) Program](#). (2022).
4. University of Florida (UF); Institute of Food and Agricultural Sciences. [Florida LAKEWATCH Program](#). (2024).
5. Florida Department of Environmental Protection (DEP); Office of Resilience and Coastal Protection (RCP); Tomoka Marsh Aquatic Preserve. [Tomoka Marsh Aquatic Preserve Continuous Water Quality Monitoring](#). (2022).