

Cape Romano-Ten Thousand Islands Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 15 July, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	15
Water Temperature - Continuous	19
Water Temperature - Discrete	24
pH - Continuous	26
pH - Discrete	28
Specific Conductivity - Continuous	30
Water Clarity	32
Turbidity - Continuous	32
Turbidity - Discrete	34
Total Suspended Solids - Discrete	36
Chlorophyll a, Uncorrected for Pheophytin - Discrete	38
Chlorophyll a, Corrected for Pheophytin - Discrete	40
Secchi Depth - Discrete	42
Colored Dissolved Organic Matter - Discrete	44

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Monthly average total nitrogen increased by less than 0.01 mg/L per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Increasing trend	2023	32	1989 - 2025	0.38645	0.13153	0.37438	0.00375	0.0024




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Total phosphorus showed no detectable trend between 1989 and 2025.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	2423	32	1989 - 2025	0.04349	0.07862	0.04309	0.00013	0.0582




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: At three program locations, monthly average dissolved oxygen decreased between 0.02 and 0.02 mg/L per year. No detectable change in monthly average dissolved oxygen was observed at one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	No detectable trend	612445	23	2002 - 2026	5.4	-0.03	5.21	0.00	0.51
rkbfuwq	Decreasing trend	644958	22	2002 - 2026	5.1	-0.30	5.27	-0.02	0.00
rkmbwbq	Decreasing trend	665650	24	2000 - 2026	4.4	-0.24	4.86	-0.02	0.00
rkbpbwq	Decreasing trend	341310	11	2016 - 2026	5.0	-0.15	5.17	-0.02	0.04




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Monthly average dissolved oxygen decreased by 0.01 mg/L per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Decreasing trend	10769	35	1989 - 2025	5.8	-0.08181	6.02669	-0.0108	0.0288




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

At three program locations, monthly average dissolved oxygen saturation decreased between 0.17 and 0.29% per year. No detectable change in monthly average dissolved oxygen saturation was observed at one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	No detectable trend	616751	23	2002 - 2026	79.1	0.03	78.42	0.04	0.51
rkbfuwq	Decreasing trend	645300	22	2002 - 2026	72.7	-0.20	76.89	-0.17	0.00
rkmbwbq	Decreasing trend	671974	24	2000 - 2026	65.4	-0.21	73.09	-0.23	0.00
rkbpbwq	Decreasing trend	344747	11	2016 - 2026	72.9	-0.15	77.04	-0.29	0.04




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 5: There was insufficient data to fit a model for dissolved oxygen saturation.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Insufficient data	112	7	2008 - 2025	78.1	-	-	-	-




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 6: Salinity showed no detectable trend between 1956 and 2025.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No detectable trend	11152	45	1956 - 2025	31.9	-0.04349	33.00994	-0.02251	0.2601




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous

National Water Information System - 7




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 7: At two program locations, monthly average salinity increased by 0.25 ppt per year at one site and by 0.28 ppt per year at the other. At two program locations, monthly average salinity decreased by 0.06 ppt per year at one site and by 0.44 ppt per year at the other. No detectable change in monthly average salinity was observed at six locations. There was insufficient data to fit a model for two locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
255123081321300	Decreasing trend	1809	8	2007 - 2017	32	-0.23	35.11	-0.44	0.02
255138081321701	Increasing trend	3074	10	2017 - 2026	32	0.21	29.33	0.28	0.01
255432081303900	No detectable trend	6556	20	2006 - 2026	22	-0.05	26.67	-0.06	0.29
255443081314700	Insufficient data	1465	4	2007 - 2011	32	-	-	-	-
255532081314300	Insufficient data	902	3	2009 - 2011	31	-	-	-	-
255534081324000	No detectable trend	6250	19	2007 - 2026	32	-0.02	33.56	-0.03	0.62
255654081350200	No detectable trend	6271	19	2007 - 2026	32	-0.03	33.07	-0.03	0.52
255732081363700	No detectable trend	1434	5	2011 - 2015	34	0.25	32.15	0.52	0.1




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 17: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: At two program locations, monthly average salinity increased by 0.25 ppt per year at one site and by 0.28 ppt per year at the other. At two program locations, monthly average salinity decreased by 0.06 ppt per year at one site and by 0.44 ppt per year at the other. No detectable change in monthly average salinity was observed at six locations. There was insufficient data to fit a model for two locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	No detectable trend	711718	25	2002 - 2026	30.0	0.05	30.80	0.04	0.21
rkbfuwq	No detectable trend	725512	25	2002 - 2026	26.5	0.03	27.82	0.03	0.51
rkbbmbwq	Decreasing trend	744660	27	2000 - 2026	33.4	-0.09	34.80	-0.06	0.02
rkbpbwq	Increasing trend	341625	11	2016 - 2026	30.7	0.17	29.69	0.25	0.02




Figure 18: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous National Water Information System - 7




Figure 19: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: At six program locations, monthly average water temperature increased between 0.03 and 0.06Deg C per year. No detectable change in monthly average water temperature was observed at four locations. There was insufficient data to fit a model for two locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
255123081321300	No detectable trend	1818	8	2007 - 2017	27.3	0.1	26.46	0.03	0.38
255138081321701	No detectable trend	3093	10	2017 - 2026	27.0	0.01	26.94	0	0.97
255432081303900	Increasing trend	6615	20	2006 - 2026	27.0	0.17	26.23	0.05	0
255443081314700	Insufficient data	2011	4	2007 - 2011	29.3	-	-	-	-
255532081314300	Insufficient data	906	3	2009 - 2011	29.3	-	-	-	-
255534081324000	Increasing trend	6313	19	2007 - 2026	26.6	0.13	25.67	0.03	0.01
255654081350200	Increasing trend	6310	19	2007 - 2026	26.7	0.11	26.41	0.03	0.03
255732081363700	No detectable trend	1435	5	2011 - 2015	28.4	0.24	27.3	0.23	0.11




Figure 20: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 21: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 10: At six program locations, monthly average water temperature increased between 0.03 and 0.06Deg C per year. No detectable change in monthly average water temperature was observed at four locations. There was insufficient data to fit a model for two locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	Increasing trend	727557	25	2002 - 2026	27.0	0.22	26.26	0.04	0.00
rkbfuwq	Increasing trend	736040	25	2002 - 2026	26.9	0.26	26.08	0.06	0.00
rkbnbwq	Increasing trend	770564	27	2000 - 2026	26.8	0.24	26.09	0.04	0.00
rkbpbwq	No detectable trend	346335	11	2016 - 2026	27.3	0.05	27.12	0.02	0.53




Figure 22: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 23: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 11: Monthly average water temperature increased by 0.02Deg C per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Increasing trend	11375	43	1956 - 2025	26.8	0.09271	26.08235	0.01805	0.0056




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 25: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 12: At one program location, monthly average pH decreased by less than 0.01 pH units per year. No detectable change in monthly average pH was observed at three locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	No detectable trend	679220	25	2002 - 2026	7.8	0.04	7.83	0	0.36
rkbfuwq	No detectable trend	697654	25	2002 - 2026	7.7	0.01	7.73	0	0.88
rkbnmbwq	Decreasing trend	735914	27	2000 - 2026	7.7	-0.10	7.70	0	0.01
rkbpbwq	No detectable trend	332136	11	2016 - 2026	7.8	0.10	7.80	0	0.19




Figure 26: Map showing location of pH continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 27: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 13: Monthly average pH decreased by 0.01 pH units per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Decreasing trend	4301	36	1956 - 2025	7.9	-0.23696	8.18343	-0.00802	0




Figure 28: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Figure 29: Scatter plot of monthly average specific conductivity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: At two program locations, monthly average specific conductivity increased by 0.32 mS/cm per year at one site and by 0.35 mS/cm per year at the other. At one program location, monthly average specific conductivity decreased by 0.07 mS/cm per year. No detectable change in monthly average specific conductivity was observed at six locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
255123081321300	No detectable trend	1295	6	2010 - 2017	49.10	-0.20	53.63	-1.03	0.20
255138081321701	Increasing trend	3074	10	2017 - 2026	48.70	0.21	45.21	0.35	0.01
255432081303900	No detectable trend	5600	17	2010 - 2026	34.90	0.00	31.28	0.01	0.98
255534081324000	No detectable trend	5597	17	2010 - 2026	48.80	0.06	48.14	0.11	0.24
255654081350200	No detectable trend	5635	17	2010 - 2026	48.80	0.04	47.21	0.05	0.43
rkbfbwq	No detectable trend	711718	25	2002 - 2026	46.24	0.05	47.30	0.06	0.20
rkbfuwq	No detectable trend	725511	25	2002 - 2026	41.45	0.03	43.06	0.04	0.51
rkmbwq	Decreasing trend	744660	27	2000 - 2026	50.93	-0.09	52.43	-0.07	0.03
rkbpbwq	Increasing trend	341625	11	2016 - 2026	47.30	0.16	45.74	0.32	0.02




Figure 30: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 31: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 15: At four program locations, monthly average turbidity decreased between 0.08 and 0.36 NTU per year.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
rkbfwbq	Decreasing trend	691451	25	2002 - 2026	7	-0.39	11.21	-0.23	0
rkbfuwq	Decreasing trend	664104	25	2002 - 2026	6	-0.19	8.57	-0.08	0
rkmbwbq	Decreasing trend	722768	27	2000 - 2026	9	-0.23	12.72	-0.13	0
rkbpbwq	Decreasing trend	339797	11	2016 - 2026	10	-0.29	13.23	-0.36	0




Figure 32: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 33: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 16: Monthly average turbidity increased by 0.08 NTU per year, indicating a decrease in water clarity.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Increasing trend	4677	31	1989 - 2025	4	0.15654	4.59643	0.07792	0.0005




Figure 34: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 35: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 17: Total suspended solids showed no detectable trend between 1989 and 2025.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	500	17	1989 - 2025	10	-0.01172	10.83431	-0.00554	0.842




Figure 36: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 37: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: Monthly average chlorophyll a, uncorrected for pheophytin, increased by 0.07 ug/L per year, indicating a decrease in water clarity.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Increasing trend	2188	29	1994 - 2025	3.66293	0.27318	3.27391	0.07359	0




Figure 38: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 39: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 19: Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2002 and 2025.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	655	15	2002 - 2025	3.8	0.08519	3.09688	0.04286	0.1675




Figure 40: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.




Figure 41: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 20: There was insufficient data to fit a model for Secchi depth.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Insufficient data	220	9	2001 - 2025	-0.9	-	-	-	-




Figure 42: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 43: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 21: There was insufficient data to fit a model for colored dissolved organic matter.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data	8	1	2001 - 2001	13	-	-	-	-




Figure 44: Map showing location of discrete water quality sampling locations within the boundaries of *Cape Romano-Ten Thousand Islands Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.