

Terra Ceia Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 15 July, 2026

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Continuous	6
Dissolved Oxygen - Discrete	8
Dissolved Oxygen Saturation - Continuous	10
Dissolved Oxygen Saturation - Discrete	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Continuous	18
Water Temperature - Discrete	20
pH - Continuous	22
pH - Discrete	24
Specific Conductivity - Continuous	26
Water Clarity	28
Turbidity - Continuous	28
Turbidity - Discrete	30
Total Suspended Solids - Discrete	32
Chlorophyll a, Uncorrected for Pheophytin - Discrete	34
Chlorophyll a, Corrected for Pheophytin - Discrete	36
Secchi Depth - Discrete	38
Colored Dissolved Organic Matter - Discrete	40

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Monthly average total nitrogen decreased by 0.01 mg/L per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Decreasing trend	2472	32	1995 - 2026	0.374	-0.25382	0.5776	-0.00679	0




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Monthly average total phosphorus decreased by less than 0.01 mg/L per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Decreasing trend	3304	32	1995 - 2026	0.106	-0.12184	0.15617	-0.00123	0.0009




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Continuous




Figure 5: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 3: There was insufficient data to fit a model for one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data	8153	2	2008 - 2009	4.2	-	-	-	-




Figure 6: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 4: Dissolved oxygen showed no detectable trend between 1989 and 2026.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No detectable trend	25200	38	1989 - 2026	6.54	-0.02502	6.63293	-0.00244	0.4855




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 5: There was insufficient data to fit a model for one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data	8153	2	2008 - 2009	59.5	-	-	-	-




Figure 10: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 6: Dissolved oxygen saturation showed no detectable trend between 1993 and 2026.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No detectable trend	5286	34	1993 - 2026	90.4	0.10555	86.30012	0.10209	0.0679




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Salinity showed no detectable trend between 1966 and 2026.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No detectable trend	25653	50	1966 - 2026	30.66	-0.04557	30.9237	-0.01266	0.153




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: There was insufficient data to fit a model for four locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data	645	2	2006 - 2007	12.0	-	-	-	-
02300084	Insufficient data	696	2	2006 - 2007	11.0	-	-	-	-
023000842	Insufficient data	578	2	2006 - 2007	28.0	-	-	-	-
TCBH	Insufficient data	8304	2	2008 - 2009	32.6	-	-	-	-




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 17: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 9: There was insufficient data to fit a model for four locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data	659	2	2006 - 2007	26.6	-	-	-	-
02300084	Insufficient data	717	2	2006 - 2007	27.5	-	-	-	-
023000842	Insufficient data	571	2	2006 - 2007	27.7	-	-	-	-
TCBH	Insufficient data	8305	2	2008 - 2009	20.0	-	-	-	-




Figure 18: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 19: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 10: Monthly average water temperature increased by 0.01Deg C per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Increasing trend	26426	52	1966 - 2026	26.7	0.08246	25.00065	0.01215	0.0158




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 21: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 11: There was insufficient data to fit a model for one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data	8306	2	2008 - 2009	8.1	-	-	-	-




Figure 22: Map showing location of pH continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 23: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 12: Monthly average pH decreased by less than 0.01 pH units per year.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Decreasing trend	23408	38	1989 - 2026	8	-0.09422	8.04912	-0.00161	0.0087




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Specific Conductivity - Continuous




Figure 25: Scatter plot of monthly average specific conductivity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 13: There was insufficient data to fit a model for four locations.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
023000825	Insufficient data	710	2	2006 - 2007	18.60	-	-	-	-
02300084	Insufficient data	775	2	2006 - 2007	17.00	-	-	-	-
023000842	Insufficient data	669	2	2006 - 2007	43.10	-	-	-	-
TCBH	Insufficient data	8306	2	2008 - 2009	49.71	-	-	-	-




Figure 26: Map showing location of specific conductivity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Continuous




Figure 27: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: There was insufficient data to fit a model for one location.

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
TCBH	Insufficient data	8263	2	2008 - 2009	2	-	-	-	-




Figure 28: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Discrete




Figure 29: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 15: Turbidity showed no detectable trend between 1995 and 2026.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No detectable trend	8158	32	1995 - 2026	1.9	0.00289	2.15566	0.00056	0.9219




Figure 30: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 31: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Monthly average total suspended solids decreased by 0.57 mg/L per year, indicating an increase in water clarity.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Decreasing trend	1733	31	1995 - 2025	10.8	-0.39721	19.03459	-0.57333	0




Figure 32: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 33: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Monthly average chlorophyll a, uncorrected for pheophytin, decreased by 0.03 ug/L per year, indicating an increase in water clarity.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Decreasing trend	1104	26	1999 - 2026	3.2	-0.10382	3.45611	-0.02639	0.0138




Figure 34: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 35: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 18: There was insufficient data to fit a model for chlorophyll a, corrected for pheophytin.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data	54	6	2000 - 2022	5.1	-	-	-	-




Figure 36: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 37: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 19: Monthly average Secchi depth became deeper by 0.02 m per year, indicating an increase in water clarity.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Decreasing trend	10427	32	1995 - 2026	-1	-0.35726	-0.89659	-0.02446	0




Figure 38: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 39: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 20: There was insufficient data to fit a model for colored dissolved organic matter.

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data	707	9	2018 - 2026	5.5	-	-	-	-




Figure 40: Map showing location of discrete water quality sampling locations within the boundaries of *Terra Ceia Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.