

Supplemental methods

Animal handling methods, biollogger specifications, and calculation of arrival and departure dates are described in Robinson, et al. ¹. Satellite tracking data were filtered and processed using the R package *crawl*^{2,3} to eliminate inaccurate location points and interpolate between locations. The resulting latitude and longitude estimates were used to calculate great circle distance (in kilometers) from the Año Nuevo breeding beach (37.1083°N, 122.3366°W) for each time-latitude-longitude point in the R package *distsm* (NOTE: I don't think this is right. According to the workflow file, this was done in MATLAB with the *distance()* function.) Across all seals, foraging trip timing (mean \pm SD day-of-year) was as follows: departure 157 ± 9 , turnaround 287 ± 40 , and arrival 15 ± 8 (Figure 1C). Therefore, outbound trip durations were 130 ± 41 days, and inbound trip durations were 93 ± 41 days. Turnaround dates were calculated using Gaussian kernels with standard deviation 6 hours using custom functions in R. Code and data for a subset of animals are available on GitHub (link available upon review of manuscript) (NOTE: The GitHub repo will be archived on Zenodo, so cite that instead when ready.) Drift rate dates were calculated using...⁴ Dates are presented as day-of-year relative to parturition date, with negative numbers indicating dates before pupping. All analyses were carried out in R v4.0.2. A linear mixed-effects model of turnaround date (relative to pupping date) as a function of turnaround distance and buoyancy change date was run in the package *lme4*⁵ after scaling and centering the continuous variables and including individual as a random effect.

Figure S1 will go here. There's a bug in analysis/data/9SupplementalFigure.R:105.

1. Robinson, P., Costa, D., Crocker, D., Gallo-Reynoso, J., Champagne, C., Fowler, M., Goetsch, C., Goetz, K., Hassrick, J., Hückstädt, L., et al. (2012). Foraging behavior and success of a mesopelagic predator in the northeast pacific ocean: Insights from a data-rich species, the northern elephant seal. *PLoS ONE* 7, e36728.
2. Johnson, D., London, J., Lea, M., and Durban, J. (2008). Continuous-time correlated random walk model for animal telemetry data. *Ecology* 89, 1208–1215.
3. Johnson, D., Josh M. London (NOAA), and Kenady (2016). *Crawl*: V2.0.
4. Robinson, P., Simmons, S., Crocker, D., and Costa, D. (2010). Measurements of foraging success in a highly pelagic marine predator, the northern elephant seal. *Journal of Animal Ecology* 79, 1146–1156.
5. Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using *lme4*. *Journal of Statistical Software* 67.