

Homework 3

Due date: February 6th, 2025

Martin Coyne, Flora Dedvukaj, Jiahao Gao, Anton Karabushin, Zhihan Lin, Joshua Morales

1. *Proof.* It is sufficient to show that

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = g'(p)$$

since letting $\lambda = g'(p)$ would complete the proof. By the mean value theorem on $[p_n, p]$, or $[p, p_n]$, there exists a $\xi_n \in (p_n, p)$ such that

$$\frac{|g(p_n) - g(p)|}{|p_n - p|} = \frac{|p_{n+1} - p|}{|p_n - p|} = g'(\xi_n).$$

Since $p_n \xrightarrow{n \rightarrow \infty} p$, we must have $\xi_n \xrightarrow{n \rightarrow \infty} p$. Therefore, by the continuity of g' , we must have

$$g'(\xi_n) \xrightarrow{n \rightarrow \infty} g'(p).$$

Therefore, taking the limit of both sides of the first equation gives

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lim_{n \rightarrow \infty} g'(\xi_n) = g'(p)$$

which is the desired result. □

2. (a) Note that

$$f'(x) = 5x^4 - 7.2x^2 + 1.28x + 1.536 \implies f'(0.8) = 0$$

$$f''(x) = 20x^3 - 14.4x + 1.28 \implies f''(0.8) = 0$$

$$f'''(x) = 60x^2 - 14.4 \implies f'''(0.8) = -14.4 \neq 0$$

Therefore, 0.8 is a root of multiplicity 3 of f .

(b)

n	x_n
0	2
1	1.68
2	1.4363
3	1.2536
4	1.1190
5	1.0216
6	0.9038
7	0.9038
8	0.8703
9	0.8474
10	0.8318

- (c) Using $\mu(x) = \frac{f(x)}{f'(x)}$, we see the following:

n	x_n
0	2
1	0.6286
2	0.78835
3	0.79995
4	0.80000
5	0.80000
6	0.80000
7	0.80000
8	0.80000
9	0.80000
10	0.80000

which clearly converges much faster.

3. (a)
- (b)
- (c)

4. *Proof.* Suppose there exists, two degree n polynomials, p_1 and p_2 , such that

$$p_1(x_i) = y_i = p_2(x_i)$$

for all $0 \leq i \leq n$. It suffices to show that $p_1(x) = p_2(x)$. Therefore, the polynomial,

$$f(x) = p_1(x) - p_2(x)$$

is a polynomial of degree at most n , with $n + 1$ distinct roots, x_0, x_1, \dots, x_n . However, by the Fundamental Theorem of Algebra, f must be the 0 polynomial.¹ Therefore,

$$f(x) = 0,$$

which means that $p_1(x) = p_2(x)$, which is the desired result. \square

5. (a) We can take $(0, 7), (2, 11), (3, 28)$, and $(4, 63)$ as $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$ respectively. We find the following:

$$L_0(x) = \frac{(x - x_1)(x - x_2)(x - x_3)}{(x_0 - x_1)(x_0 - x_2)(x_0 - x_3)}$$

$$L_1(x) = \frac{(x - x_0)(x - x_2)(x - x_3)}{(x_1 - x_0)(x_1 - x_2)(x_1 - x_3)}$$

$$L_2(x) = \frac{(x - x_0)(x - x_1)(x - x_3)}{(x_2 - x_0)(x_2 - x_1)(x_2 - x_3)}$$

$$L_3(x) = \frac{(x - x_0)(x - x_1)(x - x_2)}{(x_3 - x_0)(x_3 - x_1)(x_3 - x_2)}$$

¹Since otherwise it would be a non-zero degree n polynomial with more than n distinct roots.

Plugging in the points from above, we get:

$$L_0(x) = \frac{(x-2)(x-3)(x-4)}{(-2)(-3)(-4)} = \frac{-1}{24}(x-2)(x-3)(x-4)$$

$$L_1(x) = \frac{(x-0)(x-3)(x-4)}{(-2)(-1)(-2)} = \frac{1}{4}(x)(x-3)(x-4)$$

$$L_2(x) = \frac{(x-0)(x-2)(x-4)}{(3)(1)(-1)} = \frac{-1}{3}(x)(x-2)(x-4)$$

$$L_3(x) = \frac{(x-0)(x-2)(x-3)}{(4)(2)(1)} = \frac{1}{8}(x)(x-2)(x-3)$$

Using the y-values from above, we get the following interpolation:

$$P(x) = \frac{-7}{24}(x-2)(x-3)(x-4) + \frac{11}{4}(x)(x-3)(x-4) - \frac{28}{3}(x)(x-2)(x-4) + \frac{63}{8}(x)(x-2)(x-3)$$

Which can be simplified to become

$$P(x) = x^3 - 2x + 7$$

(b) To find the approximation of $f(1)$, we just plug in 1 for x in $P(x)$:

$$P(1) = 1^2 - 2(1) + 7 = 1 - 2 + 7 = 6$$

(c)

$$\int_0^4 x^3 - 2x + 7 = 76$$

6.

7. We know that

$$|f(x) - p(x)| = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{i=0}^n x - x_i$$

Since x_0, x_1, \dots, x_n are evenly spaced,

$$\prod_{i=0}^n (x - x_i) \leq \frac{1}{4} \left(\left(\frac{x_n - x_0}{n} \right)^{n+1} (n!) \right)$$

So,

$$|f(x) - p(x)| = \left| \left(\frac{f^{(11)}(\xi)}{11!} \right) \left(\frac{1}{4} \right) \left(\frac{1.6875 - 0}{10} \right)^{11} (10!) \right|$$

$f(x) = \sin(0.16875x)$, so

$$|f(x) - p(x)| = \max \left(-\frac{((0.16875)^{11} \cos(0.16875)(0))}{11!}, -\frac{((0.16875)^{11} \cos(0.16875)(1.6785))}{11!} \right)$$

.

Since

$$| - ((0.16785)^{11}(\cos(0.16785)(x = 0))) | = 3.15996008e^{-9}$$

and

$$| - ((0.16785)^{11}(\cos(0.16785)(x = 1.6875))) | = 3.15867894e^{-9}$$

The maximum value of f^{11} is at $x = 0$. Then,

$$|(-\frac{(0.16875)^{11}(\cos(0.16875(1.6875)))}{11!})\frac{1}{4}(0.16875)^{11}(10!) \leq 2.178e^{-19}$$

.

Then, the error bound of $|f(x) - p(x)| \leq 2.178(10^{-9})$ on $[0, 1.6875]$.