

## Homework 8

Due date: April 9th, 2025

- 1.
2. First we note that for all  $0 \leq k \leq n$ , we have that

$$\left\langle \frac{1}{\sqrt{2}}, \cos(kx) \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{1}{\sqrt{2}} \cos(kx) dx = \frac{1}{\pi} \left[ \frac{\sin(kx)}{k} \right] \Big|_{-\pi}^{\pi} = 0$$

where we used the fact that  $\sin(k\pi) = 0$  for all integers  $k$ .

Furthermore, note that

$$\left\langle \frac{1}{\sqrt{2}}, \sin(kx) \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{1}{\sqrt{2}} \sin(kx) dx = 0$$

since  $\sin(kx)$  is an odd function for all integers  $k$ .

Next, note that

$$\left\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{1}{2} dx = \frac{1}{\pi} \frac{2\pi}{2} = 1.$$

Now for all positive integers  $m$  and  $n$ , we first note that  $\cos(mx) \sin(nx)$  is an odd function, so they will have an inner product of 0. Considering  $\cos(nx)$  and  $\cos(mx)$ , we have two cases. If  $n = m$ , we have that

$$\begin{aligned} \langle \cos(mx), \cos(mx) \rangle &= \frac{1}{\pi} \int_{-\pi}^{\pi} \cos^2(mx) dx = \frac{1}{\pi} \int_0^{\pi} (1 + \cos(2mx)) dx = \frac{1}{\pi} \left( x + \frac{1}{2m} \sin(2mx) \right) \Big|_0^{\pi} \\ &= \frac{1}{\pi} \pi = 1. \end{aligned}$$

If  $n \neq m$ , then we have

$$\begin{aligned} \langle \cos(nx), \cos(mx) \rangle &= \frac{2}{\pi} \int_0^{\pi} \cos(nx) \cos(mx) dx = \frac{1}{\pi} \int_0^{\pi} (\cos((n-m)x) + \cos((n+m)x)) dx \\ &= \frac{1}{\pi} \left( \frac{1}{n-m} \sin((n-m)x) + \frac{1}{n+m} \sin((n+m)x) \right) \Big|_0^{\pi} = 0 \end{aligned}$$

where we once again used that  $\sin(k\pi) = 0$ .

We finally consider  $\sin(nx)$  and  $\sin(mx)$ , which once again has two cases. If  $n = m$ , then

$$\begin{aligned} \langle \sin(mx), \sin(mx) \rangle &= \frac{2}{\pi} \int_0^{\pi} \sin^2(mx) dx = \frac{1}{\pi} \int_0^{\pi} (1 - \cos(2mx)) dx = \frac{1}{\pi} \left( x - \frac{1}{2m} \sin(2mx) \right) \Big|_0^{\pi} \\ &= \frac{1}{\pi} \pi = 1. \end{aligned}$$

If  $n \neq m$ , then we have

$$\begin{aligned}\langle \sin(nx), \sin(mx) \rangle &= \frac{2}{\pi} \int_0^\pi \sin(nx) \sin(mx) dx = \frac{1}{\pi} \int_0^\pi (\cos((n-m)x) - \cos((n+m)x)) dx \\ &= \frac{1}{\pi} \left( \frac{1}{n-m} \sin((n-m)x) - \frac{1}{n+m} \sin((n+m)x) \right) \Big|_0^\pi = 0\end{aligned}$$

by the same reasons as above. Therefore, the set is an orthonormal set, which is the desired result.

3. (a) We seek the continuous Fourier approximation  $S_3(x)$  of the function  $f(x) = e^x$  on the interval  $[-\pi, \pi]$ , using the formula:

$$S_n(x) = \langle f(x), 1 \rangle \cdot \frac{1}{2} + \sum_{k=1}^n (\langle f(x), \sin(kx) \rangle \cdot \sin(kx) + \langle f(x), \cos(kx) \rangle \cdot \cos(kx))$$

with the inner product defined as:

$$\langle f(x), g(x) \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x)g(x) dx$$

$$\begin{aligned}a_0 &= \langle f(x), 1 \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} e^x dx = \frac{1}{\pi} (e^\pi - e^{-\pi}) \\ &\Rightarrow \frac{a_0}{2} = \frac{1}{2\pi} (e^\pi - e^{-\pi})\end{aligned}$$

$$\begin{aligned}a_1 &= \frac{1}{\pi} \int_{-\pi}^{\pi} e^x \cos(x) dx = \frac{1}{\pi} \left[ \frac{e^x}{2} (\sin(x) + \cos(x)) \right]_{-\pi}^{\pi} \\ &= \frac{1}{\pi} \left( -\frac{e^\pi}{2} + \frac{e^{-\pi}}{2} \right) = -\frac{1}{2\pi} (e^\pi - e^{-\pi})\end{aligned}$$

$$\begin{aligned}b_1 &= \frac{1}{\pi} \int_{-\pi}^{\pi} e^x \sin(x) dx = \frac{1}{\pi} \left[ \frac{e^x}{2} (\sin(x) - \cos(x)) \right]_{-\pi}^{\pi} \\ &= \frac{1}{\pi} \left( \frac{e^\pi}{2} - \frac{e^{-\pi}}{2} \right) = \frac{1}{2\pi} (e^\pi - e^{-\pi})\end{aligned}$$

Using:

$$\int e^x \cos(ax) dx = \frac{e^x (a \sin(ax) + \cos(ax))}{a^2 + 1}, \quad \int e^x \sin(ax) dx = \frac{e^x (\sin(ax) - a \cos(ax))}{a^2 + 1}$$

We get:

$$\begin{aligned}a_2 &= \frac{1}{\pi} \left[ \frac{e^x (2 \sin(2x) + \cos(2x))}{5} \right]_{-\pi}^{\pi} = \frac{1}{5\pi} (e^\pi - e^{-\pi}) \\ b_2 &= \frac{1}{\pi} \left[ \frac{e^x (\sin(2x) - 2 \cos(2x))}{5} \right]_{-\pi}^{\pi} = -\frac{2}{5\pi} (e^\pi - e^{-\pi})\end{aligned}$$

$$a_3 = \frac{1}{\pi} \left[ \frac{e^x(3 \sin(3x) + \cos(3x))}{10} \right]_{-\pi}^{\pi} = -\frac{1}{10\pi}(e^{\pi} - e^{-\pi})$$

$$b_3 = \frac{1}{\pi} \left[ \frac{e^x(\sin(3x) - 3 \cos(3x))}{10} \right]_{-\pi}^{\pi} = \frac{3}{10\pi}(e^{\pi} - e^{-\pi})$$

Evaluating numerically, we have:

$$S_3(x) \approx 3.6767 - 3.6767 \cos(x) + 3.6767 \sin(x) + 1.4707 \cos(2x) - 2.9414 \sin(2x) - 0.7353 \cos(3x) + 2.2060 \sin(3x)$$

(b) We must calculate  $\langle y, \phi_k \rangle$  and  $\langle y, \psi_k \rangle$  for  $k = 0, 1, 2, 3$ . First, calculate  $\langle y, \phi_0 \rangle$ ,

$$\langle y, \phi_k \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \cos(kx_j)$$

$\langle y, \phi_1 \rangle$ ,  $\langle y, \phi_2 \rangle$ , and  $\langle y, \phi_3 \rangle$ .

$$\langle y, \phi_0 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \cos(0) = \frac{1}{10} \sum_{j=0}^{19} y_j$$

$$\langle y, \phi_1 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \cos(x_j)$$

$$\langle y, \phi_2 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \cos(2x_j)$$

$$\langle y, \phi_3 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \cos(3x_j)$$

Next, calculate  $\langle y, \psi_k \rangle$  for  $k = 1, 2, 3$

$$\langle y, \psi_k \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \sin(kx_j)$$

$$\langle y, \psi_1 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \sin(x_j)$$

$$\langle y, \psi_2 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \sin(2x_j)$$

$$\langle y, \psi_3 \rangle = \frac{1}{10} \sum_{j=0}^{19} y_j \sin(3x_j)$$

$$S_3(x) = \frac{1}{2}\langle y, \phi_0 \rangle + \sum_{k=1}^3 (\langle y, \phi_k \rangle \cos(kx)) + \sum_{k=1}^3 (\langle y, \psi_k \rangle \sin(kx))$$

$$S_3(x) = \frac{1}{2}\langle y, \phi_0 \rangle + \langle y, \phi_1 \rangle \cos(x) + \langle y, \phi_2 \rangle \cos(2x) + \langle y, \phi_3 \rangle \cos(3x) + \langle y, \psi_1 \rangle \sin(x) + \langle y, \psi_2 \rangle \sin(2x) + \langle y, \psi_3 \rangle \sin(3x)$$

$$S_3(x) \approx 3.676 + 2.963 \cos(x) + 0.946 \cos(2x) + 0.164 \cos(3x) - 1.263 \sin(x) - 0.740 \sin(2x) - 0.214 \sin(3x)$$

4. (a) Q is orthogonal.

$$Q^t Q = I$$

$$\det(Q^t Q) = 1$$

$$\det(Q^t) \det(Q) = 1$$

Since  $\det(Q^t) = \det(Q)$ ,

$$\det(Q)^2 = 1$$

$$\det(Q) = \pm 1$$

(b)

$$(PQ)^t PQ = Q^t P^t PQ = Q^t (I) Q = Q^t Q = I$$

. So PQ is orthogonal.

5. (a)

(b)

(c)

6. We are given the symmetric matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ -1 & 1 & 2 \\ 2 & 2 & 2 \end{bmatrix}$$

We want to decompose it as:

$$A = Q D Q^T$$

To find the eigenvalues, we solve the characteristic polynomial:

$$\det(A - \lambda I)$$

$$A - \lambda I = \begin{bmatrix} 1 & -1 & 2 \\ -1 & 1 & 2 \\ 2 & 2 & 2 \end{bmatrix} - \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$= \begin{bmatrix} 1-\lambda & -1 & 2 \\ -1 & 1-\lambda & 2 \\ 2 & 2 & 2-\lambda \end{bmatrix}$$

We know can find the determinant of the matrix:

$$(1-\lambda) \begin{bmatrix} 1-\lambda & 2 \\ 2 & 2-\lambda \end{bmatrix} - (-1) \begin{bmatrix} -1 & 2 \\ 2 & 2-\lambda \end{bmatrix} + 2 \begin{bmatrix} -1 & 1-\lambda \\ 2 & 2 \end{bmatrix} = -\lambda^3 + 4\lambda^2 + 4\lambda - 16 = -(\lambda-4)(\lambda-2)(\lambda+2)$$

The eigenvalues are:

$$\lambda_1 = 4, \quad \lambda_2 = 2, \quad \lambda_3 = -2$$

Solve  $(A - 4I)\vec{v} = 0$ , we get:

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \quad \|\vec{v}_1\| = \sqrt{6} \Rightarrow q_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

Solve  $(A - 2I)\vec{v} = 0$ , we get:

$$\vec{v}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \quad \|\vec{v}_2\| = \sqrt{2} \Rightarrow q_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Solve  $(A + 2I)\vec{v} = 0$ , we get:

$$\vec{v}_3 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \quad \|\vec{v}_3\| = \sqrt{3} \Rightarrow q_3 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

Orthogonal matrix  $Q$  (columns are the normalized eigenvectors):

$$Q = \begin{bmatrix} \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \\ \frac{2}{\sqrt{6}} & 0 & -\frac{1}{\sqrt{3}} \end{bmatrix}$$

Diagonal matrix  $D$  of eigenvalues:

$$D = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -2 \end{bmatrix}$$