

Homework 8

Due date: April 3rd, 2025

1. (a)
- (b)
- (c)

2.

3. *Proof.* First, note that since $x^2 \geq 0$ for all $x \in \mathbb{R}$ and $f \in C[a, b]$, we have that

$$\langle f, f \rangle = \int_a^b f(x)^2 dx \geq 0.$$

Furthermore, we have that

$$\langle f, f \rangle = 0 \iff \int_a^b f(x)^2 dx = 0 \iff f(x)^2 = 0 \iff f(x) = 0$$

for all $x \in [a, b]$. This is also due to the non-negativity of $f(x)^2$. Next, for any $f, g \in C[a, b]$ we have that

$$\langle f, g \rangle = \int_a^b f(x)g(x) dx = \int_a^b g(x)f(x) dx = \langle g, f \rangle.$$

Finally, for any $f, g, h \in C[a, b]$ and $c_1, c_2 \in \mathbb{R}$, by the linearity of the Riemann integral, we have that

$$\begin{aligned} \langle c_1 f(x) + c_2 g(x), h(x) \rangle &= \int_a^b (c_1 f(x) + c_2 g(x))h(x) dx = \int_a^b (c_1 f(x)h(x) + c_2 g(x)h(x)) dx \\ &= c_1 \int_a^b f(x)h(x) dx + c_2 \int_a^b g(x)h(x) dx = c_1 \langle f, h \rangle + c_2 \langle g, h \rangle. \end{aligned}$$

Therefore, $\langle \cdot, \cdot \rangle$ is an inner product, which is the desired result. ■

4. We are given that $\mathbf{v} = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 0 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 1 \\ -1 \\ 2 \\ -2 \end{pmatrix}$.

(a) $\langle \mathbf{v}, \mathbf{w} \rangle = 1 * 1 + 3 * (-1) + 2 * 2 + 0 * (-2) = 2$

(b) $\|\mathbf{v}\| = \sqrt{1^2 + 3^2 + 2^2} = \sqrt{14}$

(c) $\mathbf{v} - \mathbf{w} = \begin{pmatrix} 0 \\ 4 \\ 0 \\ 2 \end{pmatrix}$

$$\|\mathbf{v} - \mathbf{w}\| = \sqrt{0^2 + 4^2 + 0^2 + 2^2} = 2\sqrt{5}$$

$$(d) \cos(\theta) = \frac{\langle \mathbf{v}, \mathbf{w} \rangle}{\|\mathbf{v}\| \cdot \|\mathbf{w}\|} = \frac{2}{\sqrt{14} \cdot \sqrt{1^2 + (-1)^2 + 2^2 + (-2)^2}} = \frac{2}{\sqrt{140}}$$

$$\theta = \cos^{-1}\left(\frac{2}{\sqrt{140}}\right) \approx 1.40095$$

5. (a)

$$\begin{aligned} \langle f, g \rangle &= \int_0^1 f(x)g(x)dx \\ &= \int_0^1 (x-1)(x+1)dx \\ &= \int_0^1 (x^2 - 1)dx \\ &= \left[\frac{x^3}{3} - x\right]_0^1 \\ &= \left(\frac{1}{3} - 1\right) - \left(\frac{0^3}{3} - 0\right) = \frac{-2}{3} \end{aligned}$$

(b)

$$\begin{aligned} \|f - g\| &= \sqrt{\int_0^1 |(f - g)|dx} \\ f(x) - g(x) &= (x-1) - (x+1) = x-1-x-1 = -2 \\ |f(x) - g(x)| &= 2 \\ \int_0^1 2dx &= [2x]_0^1 = 2 - 0 = 2 \end{aligned}$$

(c) The angle between f and g is

$$\cos(\theta) = \frac{\langle f, g \rangle}{\|f\| \cdot \|g\|}$$

We already know that $\langle f, g \rangle = \frac{-2}{3}$, so we need to find the norm $\|f\|$.

$$\begin{aligned} \|f\| &= \left(\int_0^1 (x-1)^2 dx\right)^{\frac{1}{2}} = \left(\int_0^1 (x^2 - 2x + 1) dx\right)^{\frac{1}{2}} \\ &= \left(\left[\frac{x^3}{3} - 2\frac{x^2}{2} + x\right]_0^1\right)^{\frac{1}{2}} = \left(\left(\frac{1}{3} - 2\frac{x^2}{2} + x\right) - (0)\right)^{\frac{1}{2}} \\ &= \left(\frac{1}{3} - 1 + 1\right)^{\frac{1}{2}} = \frac{1}{\sqrt{3}} \end{aligned}$$

.

Now we need to find $\|g\|$.

$$\|g\| = \left(\int_0^1 (x+1)^2 dx\right)^{\frac{1}{2}} = \left(\int_0^1 (x^2 + 2x + 1) dx\right)^{\frac{1}{2}} = \left(\left[\frac{x^3}{3} + x^2 + x\right]_0^1\right)^{\frac{1}{2}}$$

$$= \left(\frac{1}{3} + 1 + 1\right)^{\frac{1}{2}} = \sqrt{\frac{7}{3}}$$

We can now use both of these norms to find θ :

$$\cos(\theta) = \frac{\frac{-2}{3}}{\frac{-1}{3} \cdot \sqrt{\frac{7}{3}}} = \frac{\frac{-2}{3}}{\sqrt{\frac{7}{9}}} = \frac{\frac{-2}{3}}{\frac{\sqrt{7}}{3}} = \frac{-2}{\sqrt{7}}$$

Now we can solve for θ :

$$\theta = \cos^{-1}\left(\frac{-2}{\sqrt{7}}\right) \approx \cos^{-1}(-0.7559) \approx 139.1^\circ$$

- (d) To find a nonzero h that is perpendicular to f , we would have to find a function $h(x)$ that satisfies the following:

$$\int_0^1 (x-1) \cdot h(x) dx = 0$$

We can try a simple polynomial $h(x) = ax + b$, and plug it into the above orthogonality condition:

$$\int_0^1 (x-1)(ax+b) dx = 0$$

$$(x-1)(ax+b) = (ax^2 + bx - ax - b) = ax^2 + (b-a)x - b$$

$$\int_0^1 (ax^2 + (b-a)x - b) dx = a\frac{1}{3} + (b-a)\frac{1}{2} - b = 0$$

$$2a + 3(b-a) - 6b = 0 \rightarrow 2a + 3b - 3a - 6b = 0 \rightarrow -a - 3b = 0 \rightarrow a = -3b$$

Let us pick $b = 1$, then $a = -3$, so $h(x) = -3x + 1$

6. We know that $\mathbf{v}_1 = (1, 2, 2)$, $\mathbf{v}_2 = (-1, 0, 2)$, and $\mathbf{v}_3 = (0, 0, 1)$

(a)

$$\mathbf{v}_1 = \mathbf{x}_1 \text{ and } \|\mathbf{x}_1\|_2 = \sqrt{1^2 + 2^2 + 2^2}$$

$$\|\mathbf{w}\| = \frac{\mathbf{x}_1}{\|\mathbf{x}_1\|_2} = \left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}\right)$$

$$\mathbf{x}_2 = \mathbf{v}_2 - \langle \mathbf{v}_2, \mathbf{w}_1 \rangle \mathbf{w}_1 = (-1, 0, 2) - \langle (-1, 0, 2), \left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}\right) \rangle$$

$$= (-1, 0, 2) - 1\left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}\right)$$

$$= \left(\frac{-4}{3}, \frac{-2}{3}, \frac{4}{3}\right)$$

$$\|\mathbf{x}^2\|_2 = \sqrt{\left(\frac{-4}{3}\right)^2 + \left(\frac{-2}{3}\right)^2 + \left(\frac{4}{3}\right)^2} = 2$$

$$\mathbf{w}_2 = \frac{\mathbf{x}_2}{\|\mathbf{x}_2\|_2} = \left(\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}\right)$$

$$\begin{aligned}
\mathbf{x}_3 &= \mathbf{v}_3 - \langle \mathbf{v}_3, \mathbf{w}_1 \rangle \mathbf{w}_1 - \langle \mathbf{v}_3, \mathbf{w}_2 \rangle \mathbf{w}_2 \\
\mathbf{x}_3 &= (0, 0, 1) - \langle (0, 0, 1), (\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) \rangle (\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) - \langle (0, 0, 1), (\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}) \rangle (\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}) \\
&= (0, 0, 1) - \frac{2}{3} (\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) - \frac{2}{3} (\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}) \\
&= (\frac{2}{9}, \frac{-2}{9}, \frac{1}{9}) \\
\|\mathbf{x}_3\|_2 &= \frac{1}{3} \\
\mathbf{w}_3 &= \frac{\mathbf{x}_3}{\|\mathbf{x}_3\|_2} = (\frac{2}{3}, \frac{-2}{3}, \frac{1}{3})
\end{aligned}$$

(b) We know $\mathbf{x} = (7, 5, 1)$

$$\begin{aligned}
\langle (7, 5, 1), (\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) \rangle &= \frac{19}{3} \\
\langle (7, 5, 1), (\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}) \rangle &= \frac{-17}{3} \\
\langle (7, 5, 1), (\frac{2}{3}, \frac{-2}{3}, \frac{1}{3}) \rangle &= \frac{5}{3}
\end{aligned}$$

.

Now, we can write

$$\frac{19}{3} (\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) - \frac{17}{3} (\frac{-2}{3}, \frac{-1}{3}, \frac{2}{3}) + \frac{5}{3} (\frac{2}{3}, \frac{-2}{3}, \frac{1}{3})$$