

Set Theory Notes

by Tyler Wright

github.com/Fluxanoia

fluxanoia.co.uk

These notes are not necessarily correct, consistent, representative of the course as it stands today or, rigorous. Any result of the above is not the author's fault.

0 Notation

We commonly deal with the following concepts in Set Theory which I will abbreviate as follows for brevity:

Term	Notation

Contents

0	Notation	1
1	The Fundamentals	3
1.1	Axiom of Extensionality	3
1.2	Axiom of Pair Sets	3
1.3	Axiom of the Powerset	3
1.4	Axiom of the Empty Set	3
1.5	Axiom of Subsets	3
1.6	Axiom of Unions	4
1.7	Classes	4
1.8	The Set ω	4
1.9	Russell's Theorem	4
1.10	The Universe of Sets	5

1 The Fundamentals

1.1 Axiom of Extensionality

For two sets a and b , we have that $a = b$ if and only if for all x we have that:

$$x \in a \iff x \in b.$$

For two classes A and B , we have that $A = B$ if and only if for all x we have that:

$$x \in a \iff x \in b.$$

1.2 Axiom of Pair Sets

For any sets x and y , there is a set $z = \{x, y\}$. This is the (unordered) pair set of x and y .

1.3 Axiom of the Powerset

For each set x , there exists a set which is the collection of the subsets of x , the powerset $\mathcal{P}(x)$.

For some set x , we have the powerset defined as follows $\mathcal{P}(x) = \{z \mid z \subseteq x\}$.

1.4 Axiom of the Empty Set

There exists a set with no members, the empty set \emptyset .

We have the empty set defined as follows $\emptyset = \{x \mid x \neq x\}$.

1.5 Axiom of Subsets

For some set x , we have that $\{y \in x \mid \Phi(y)\}$ is a set for some well-defined property of sets Φ .

1.6 Axiom of Unions

We have the basic union of two sets x_1 and x_2 :

$$x_1 \cup x_2 = \{y \mid y \in x_1 \text{ or } y \in x_2\},$$

but for cases where we want to unify the members of the sets in a set X , we define:

$$\bigcup X = \{y \mid \exists x \in X, y \in x\}.$$

This axiom states that for a set X , $\bigcup X$ is a set.

1.7 Classes

We have that classes are collection of objects, these could also be sets. Classes that are not sets are called proper classes.

1.8 The Set ω

We have the set of natural numbers, $\mathbb{N} = \{0, 1, 2, \dots\}$, and from this, we define ω :

$$\omega = \{0, 1, 2, \dots\},$$

where for some n in ω ,

$$n = \{0, 1, 2, \dots, n-1\},$$

with 0_ω being the empty set. We can go beyond this definition, defining:

$$\begin{aligned}\omega + 1 &= \{0, 1, 2, \dots, \omega\}, \\ \omega + 2 &= \{0, 1, 2, \dots, \omega, \omega + 1\}, \\ &\dots \\ \omega + n &= \{0, 1, 2, \dots, \omega, \omega + 1, \dots, \omega + n - 1\}.\end{aligned}$$

1.9 Russell's Theorem

We have that $R = \{x \mid x \notin x\}$ is not a set.

Proof. Suppose we have a set z such that $z = R$, is z in R ? If we suppose z is in R , we have that z is not in z by the definition of R (as $z = R$) but z is R so z is not in R , a contradiction. Thus, we have that there is no set z equal to R , so R is not a set but a proper class. \square

1.10 The Universe of Sets

We define the universe of sets as $V = \{x \mid x = x\}$. We have that V is a proper class.

Proof. If we suppose V is a set, we apply the axiom of subsets with $\Phi(x) = x \notin x$ and reach a contradiction via Russell's theorem. \square