Table 1. Model-supported intersections between Asian Koel acoustic activity and planning-sensitive urban spaces in Shenzhen.

A. Functional zones within Primary Results + Moderate Extrapolation HEXes

Functional zone	Overlap area (km^2)	Annual mean ψ	W10–W30 mean ψ	Peak week	Expected dawn-positive sampled minutes	Dawn $P(K_{window} \geq 1)$
Residential	29.23	39.6%	71.4%	W20	0.29	17.2%
Commercial	4.61	40.0%	73.9%	W20	0.40	20.5%
Industrial	28.78	39.5%	73.1%	W20	0.36	19.6%

B. Planning-sensitive places within Primary Results + Moderate Extrapolation HEXes

Sensitive-place class	POIs (n)	Annual mean ψ	W10–W30 mean ψ	Peak week	Expected dawn-positive sampled minutes	Dawn $P(K_{window} \geq 1)$
School and childcare	269	39.5%	70.7%	W21	0.27	16.7%
Higher education	26	37.9%	68.6%	W20	0.23	15.4%
Hospital	23	38.0%	69.7%	W20	0.30	17.7%
Other healthcare and care	18	40.3%	73.9%	W20	0.40	20.1%
Park and garden	374	40.2%	75.0%	W20	0.45	22.0%
Accommodation	53	42.5%	75.5%	W20	0.44	21.2%

Note. Values for ψ and the dawn metrics are posterior medians; 95% credible intervals are reported in Supplementary Data 15 (Tables S15B-S15C). Annual mean ψ covers weeks 1-52, and the seasonal mean covers weeks 10-30. Peak week maximises the posterior-mean ψ curve.

Dawn metrics describe 24 scheduled one-minute samples between 04:00 and 08:00 on a reference day, averaged over weeks 15-21. K_{window} is the number of Koel-positive samples. Expected counts and $P(K_{window} \geq 1)$ were derived from the fitted beta-binomial process with positive-day conditioning.

Summaries include hexagonal (HEX) cells classified as Primary Results or Moderate Extrapolation. Functional zones were weighted by polygon-HEX overlap area; points of interest (POIs) were equally weighted after deduplication. These are modelled acoustic co-location summaries under the stated sampling design.