

-, i? i l



GENERAL INSTRUCTION AND DEVELOPMENT OF SENSITIVITY
IN 30 LESSONS OVER ! 50 EASY EXERCISES

Appendix: Processes of the MENTAL Méfhode, by P BÉASSE

IGNORED RICHES

CLASSES

OF

,

RADIESTHESIA

PRACTICE

FROM THE SAME AUTHORS:

Works published:

Theoretical and Practical Elementary Treaty of Physical Dowsing. by J. CORRENSON. Former student of the Ecole Polytechnique. 1936. (Ep U ized) .

New simple and rational processes of Physical Dowsing. by P. BÉASSE, Consulting Technician. 1938. (Out of print) .

Dowsing explained and practiced by the Physical Method. by P. BÉASSE, Consulting Technician. 1939. {Three editions: French, English, Spanish} .

Practical Dowsing Course for beginners of the Physical Method. by J. CORRENSON, Alumnus of the Ecole Poly technique , and P. BÉASSE, Technician-Advisor. 1943. TOME 1: General Instruction and Development of Sensitivity (followed by an Appendix on the processes of the Mental Method, by P. BÉASSE).

In preparation for : *

Practical Dowsing Course for beginners of the Physical Method. by P. BÉASSE, Consulting Technician. VOLUME II: Various Applications and Prospecting.

Scientific Censorship Control visa, General Secretariat* for Information: N° 8.089, of September 30, 1943 .

Copyright by Pierre BÉASSE, Publisher . • 1943. — Rights of reproduction , translation and adaptation reserved for all countries.

COLLECTION : "RICHESSSES CACHÉES

To learn without a teacher

CLASSES

r-

of

RADIESTHESIA PRACTICE

for use by beginners

PHYSICAL METHOD

you

accessible **to all** , excluding any occultism

by

J. CORRENSON •

**Former student
of the Ecole Polytechnique**

P. BÉASSE

**Technicien-Consoil
Radio-Physicien**



General Instructions. — Development of Sensitivity ■

30 Lessons — 150 Easy Exercises

Appendix:

Processes of the MENTAL Method, by P. BÉASSE

EDITIONS OF SCIENTIFIC PROGRESS:.

37, Rue Rossini — NICE (Alpes-Maritimes)

work crowned by the Union of Journalists and Writers Competition of 1943

AAA»---AAAAAA K.

aupre s de nos Lecteurs.
Due to the consequences of
the war, we not can

offer that ' a paper grade 1

lower, good that of pnx I
e i eve . We apologize

I TABLE OF CONTENTS OF VOLUME I

	Pages
WARNING	V
ABBREVIATIONS . . . v	VII

FOREWORD. — What is Dowsing 1 >

The outside world. Its knowledge "with the help of sensitive perceptions. The 5
usual external senses. The 6TM external sense, in animals, in primitive man, in
civilized peopleIX

Dowsing. Its definition. Its object. The dowsing phenomenon. Is dowsing a

FIRST CHAPTER. — What is Radiesthetics 1 I

§ 1" r Microphysics 3

§ 2. The radiation of matter. (Electromagnetic waves , including specific radiation.
Corpuscular radiation) 3

§ 1 . l ^e ^3 tri ^3 11 ^3 > » ^9

§ 4 Magnetism 1 J

§ 5 Radioactivity 15

§ 6 Light 17

§ 7 telluric rays . Checkerboard of Dr. Peyré
19

§ 8 Infinitesimal Biology. Human polarity 24

CHAPITRE II. — The instruments of the dowser 33

4 I Neutral wand of dowsers 34

5 ^2 . ^3 p^ ^3 n ^dulen ^3 utr ^3 ^3 ^9

6 3z The active instruments 44

CHAPITRE III. — The perception of waves	49
1 ^{r*} Lesson: Experimental adjustment of the pendulum on the sheet of vertical rays of the different bodies. The perception of specific rays	52
2 ^{**} Lesson: Polarity of bodies. Positive bodies. Negative bodies	64
3 th Lesson: Swing of the pendulum, the index of the free hand in antenna. Different reaction when this index is put in direct contact with the radiant object	72
4 ^{m*} Lesson : Swing of the pendulum by the intervention of the radius	77
5 ^{**} Lesson : Swing of the pendulum by the intervention of the ray	180
6 ^{**} Lesson : Work at Pointe -Antenne. The reflection. The Refraction	84
7 ^{m*} Lesson: Adjustment of the precision spherical neutral pendulum	89
8 ^{m^} Lesson: Self-Tuning the Pendulum. Case of the hollow pendulum. Case of the full pendulum. The colored pendulum. . .	100
9 ^{r*} Lesson: Radii at 45° and 135°	103
10 TM Lesson: Series. The Particular Number. The Number Characteristic.	1 bone
11 ^{**} Lesson: The fundamental angle. The Basic Plan.	110
12 ^{**} Lesson: The “Witnesses”	118
13 [^] Lesson: Instructions for use of the SCHUMFELL Developer of precision, anti-image	121
CHAPITRE IV. — Some microphysical phenomena of practical use	126
14 ^{**} Lesson: Fluidic Lines and Planes	126
15 ^{m^} Lesson : Fluidic fields. The magnetic Circix. The fluidic field of a rectilinear magnet. Photogenic triangle	131
16 ^{**} Lesson: The Ray of Uni^o	13 ^9
17 ^{**} Lesson: The radioactive radiation of bodies. The specific axis of simple bodies	145
18 TM Lesson: Amplification by the Action of Radioactivity ..	147
19 ^{r*} Lesson: Shape Waves	150

20^{TM*} Leçon : Les Rayons Telluriques nocifs. La pointe verticale.

21^{ra °} Lesson : The search for an underground stream 162

» •

CHAPITRE V. — The Radiation of Living Beings 17t ■

22^{ne} Lesson : The nerve rays. The vital wave. The sexual wave. The human wave

1.72

23^{TM*} Lesson: The ray lies a l . Species waves 176

24sm Lesson : The sympathetic ray. The personal wave 179

CHAPTER VI. — Images in Dowsing 183

siques. 18^{^t}

25^{mc} Leçon : Les diverses catégories d'images. Leur élimination. Tableaux
synoptiques de leurs propriétés phy- 2 6 :rn^c Lesson :

Daytime images 196

27^{mc} Lesson: Permanent Images 205

28^{r °} Lesson: Remanence and Impregnation 207

oomc 11

29 Lesson: LI m □ p reg na ti on m □ ent ale 215

30^{'''} Lesson : The Pitfalls 217

WARNING

t

•

At the time of the publication of the work "La RADIES THÉSIE explained et pratique par la METHOD PHY SIQUE" by P. BÉASSE, several correspondents had expressed the wish to see an exclusively practical guide added to it, but at the same time sufficiently clear and precise so that, with its help, a beginner in this method could easily instruct himself, without having to resort to the experience of a teacher.

It is to this wish that this manual, presented to the public today, attempts to respond. If this book is addressed before . entirely for a reader, who possesses, at least in broad outline, the essence of the aforementioned treatise, it has been composed in such a way that it could, if need be, be self-sufficient, and that the 'together. its text constituted a sufficiently complete whole. .

This is why each series of lessons is preceded by a very succinct documentation summarizing, in the minimum of words, the exposition of the questions dealt with in the exercises proposed after it, and allowing them to be carried out easily with logic and method.

A method, however good it may be, is not sufficient **on** its own. A big part of success is execution; this is why we have endeavored to adjust the execution methods as precisely as possible to further reduce the student's effort. Our goal is to be complete, because what often worries a neophyte is considered quite natural for the already informed reader.

On the other hand, Microphysics being in perpetual progress, this documentation has been supplemented with a few precisions resulting from new acquisitions.

This Course being, above all, intended for a beginner, the text strictly essential to its initiation was printed in large characters, reserving the text in smaller characters to the confirmed dowser or to the neophyte equipped with a sufficient scientific background. Made available to everyone, this practical course does away with any scientific presentation of too high an order not to be grasped immediately.

Finally, one. appendix on the processes of the MENTAL Method, by Pierre BÉASSE, has been added to this course. It is intended for practiced followers of the PHYSICAL Method, whose special sensitivity or simply developed thanks to this course, or even particular aptitudes, allow the practice of the Mental Method, for more convenience.

ABBREVIATIONS

EM waves	Electro-magnetic wave.
ES waves	Specific wave of electricity, with electrical support cudgel. ■
MS waves	•Specific wave of magnetism, with •magnetic support . ■
EM	image Electro-magnetic image.
Field El	Electric field of Physicists.
Field Mag ..	Magnetic Field of Physicists.
ES+ field. .	Limit space traveled by specific rays with positive electrical supports.
MS field — . .	Limit space traversed by specific rays with negative magnetic supports.
Field RS	Boundary space traveled by specific rays^ with radioactive support.
Image M	Magnetic image .
AR image	Radioactive image .
ERM image	Electro-radio-magnetic image .
ERA image	Electro-radioactive image .
IM	picture Mental impregnation.
Nm . - .	Magnetic North.
S ^m	Magnetic South.
N S	North-South.
EO	Is West.
NOSE ..	North. West. South. East.
	.NESO. . North.
East. South. West.	
Pf	Fundamental plan.
Pef	Electro-fundamental plan.
CL spokes.....	Heat rays.
rays	Chemical rays.
CM	Magnetic cross.

.P ou + Positive.

N ou — Negative.

More or less, following . the case.

Fundamental radius.

The words: wave, ray, radiation, radiation, vibration, .. have, in principle, the same meaning in this book. These terms are used to expose with the maximum possible clarity facts that are still more or less obscure. .

Name of the letters of the Greek alphabet used:

'1. alfa	7naked
H beta	I omicron
• gamma	ιwp«
£ epsilon	' · · ro
": dzeta	I sigma
(j theta	. tau
Υ. kappa	Ci) omega
IJ. mu	

Reports to the work : " Dowsing

Méthode Physique »

During the lessons which will follow, we will often refer the reader to the work "Dowsing explained and practiced by the Physical Method", by P. BÉASSE. He will find additional details that will facilitate his understanding and help his progress. The indication of these pages will appear in characters as follows: (Page 57 of "Dowsing Physical Method"). On the other hand, when we refer the reader to a lesson in this manual, the indication of the page will be in current characters: (Page 57).

.FOREWORD

This course includes two types of texts:

One, printed in ' normal characters, essentially practical, within everyone's reach, is intended for the prospector : — the other, printed in smaller characters. is more theoretical and aimed at technicians.

WHAT IS RADIESTHESIA

Of the external world around us, we know almost nothing. The ideas that we form about it are suggested to us only by the impression that this world exercises on the organs of each of the external senses with which the Creator has provided us. It is to the images that this impression represents, and to the relationships that they allow us to deduce between real objects that escape us. ..that our knowledge of matter is limited.

The physiological excitation determined on these organs by the action of a reality impossible to grasp produces an instantaneous modification on our sensibility. we call **sensation**. Alongside the **affective fact** that is observed in every sensation, there is produced. at the same time, in our intelligence, a fact of knowledge, a **representative fact**; and it is the sum of these two facts, indissolubly linked to each other by consciousness, which determines the sepsis image which we represent to ourselves.

Thus, when, transmitted through the air, the vibrations of a piano string strike the **drum skin** that constitutes our eardrum, we experience a sensation of shaking that our intelligence immediately represents as the sound of stamped and of a determined height . When the rays of light diffused by a colored object come, after refraction in our visual organ, to strike our retina, we instantly picture to ourselves a determined shade of color.

For the clarity of the understanding of the phenomena of which we are the perpetual witnesses, the psychologists have agreed to call sensation, their affective side; and perception their representative side, and our consciousness. reveals to us the existence, in each of them, of both the first and the second of these two aspects.

Their mechanism can, moreover, be analyzed as follows:

1. The external excitation produces, on the sensitive organ, a physiological impression;
 2. This impression is transmitted to the brain by the sensitive nerves;
 3. In the brain, the movement then takes place, following which appears, in the consciousness, the sensation, agreeable, disagreeable or indifferent, in connection with which the representation of is formed almost instantaneously. the sensible image which is born in our intelligence ;
- ''' 4-0 Finally, in most cases, the cycle is completed by a motor reaction which the motor nerves transmit to the affected organ.' It is the violent start determined by an unusual noise or the flash of a sudden clap of thunder or the flame of an unexpected explosion, it is the withdrawal of the hand in contact with an object at high temperature.

But it may also happen that the nervous shock, instead of being transmitted to the brain, is simply registered by the spinal cord and goes no further. In this case, consciousness does not appear, but the motor sensation is triggered outside by gestures, and inside, in the form of unreflected automatic movements, to which we give the name of reflexes.

The six external senses

In addition to the five usual senses to which we are indebted visual sensations. auditory, tactile, gustatory, olfactory, there is an external sixth sense. more or less perfect, more or less developed in animals or in primitive man. strongly attenuated and most often completely atrophied in civilized man, it is the dowsing sense or sense of perception on the spot or at a distance of the radiation of matter.

It is on the indications of this direction that the migratory bird goes, that the pigeon joins its dovecote, the bee its hive, that the Negro of the Center of Africa or the Redskin of the Amazon finds their hut or their camp lies, that the dog or the ferocious animal discovers the first the game, the second its prey.

The organ of this sense. in humans, is the human receptor for which each nerve end constitutes a receiving antenna. Its structure and its mechanism are still insufficiently studied, whereas they deserve to be studied with as much care as those of the organs of hearing and vision, the functioning of which is exposed in all detail in physics works. '

"In dowsing. the animals are our masters". the commander of the Bastide could say very truthfully; and, to animals, we will add primitive man, of whom the Comte de Marsay, in his work "Electricity, Magnetism, Dowsing ", **has described some traits of dowsing perception which leave him little to envy the privilege of his inferior brothers .**

In perfect beings **from** this point of view. it is infinitely probable that the mechanism of perception by the o' ^{mp} sense is absolutely identical to what it is in normal man for perception by his usual senses. The shock reaches the brain and determines the formation of a conscious image. In beings with an attenuated reception, this image is almost unconscious. Finally, in beings whose dowsing receiver is almost atrophied, the shock does not go beyond the spinal cord and determines simply **automatic** movements whose reality only appears to us when their amplitude has been considerably multiplied by the intervention of an artificial detector. It is only on the observation of these movements which have become perceptible that intelligence intervenes to interpret them.

If we extend by the wire of a regulated pendulum the nervous extremities of one of our hands, the movements of this hand, without always being visible, will nevertheless become pronounced enough to determine either the sway or the gyration. of the instrument, if it is placed in a radiant zone.

If, after having grasped the wand with both hands under the required

conditions, we enter a radiant zone, the muscles which will force it to rise or fall will obey a reflex action so violent that our will will be absolutely incapable of control it or thwart it. Our hands will have ceased to belong to us to momentarily become the plaything of an impulse, provoked by an external action and with regard to which we can only observe our complete helplessness.

With regard to an attenuated radiesthesic sense organ, the amplifier will play the role of the acoustic horn for the reduction of deafness, of glasses for the correction of abnormal sight, finally that of the microscope or the telescope when we charge these instruments to supply the imperfection and insufficiency of our visual organ.

The perception of the radiation of matter

It is in the form of corpuscular radiations or **simple** electromagnetic waves that the radiation of matter manifests itself. but it is the latter, alone, which excite the dowsing sense.

The shock of electromagnetic waves can, however, to impress sensitive organs other than that of the external sixth sense. Some dowsers see sorts of vapors on the vertical of underground streams.

rains. Others, when prospecting for a Jilon of copper ore, perceive, **according to** M. Viré, Doctor of Science, the bitter taste of verdigris; others perceive the smell of the radiant object they are looking for from a distance.

But it is the sense of touch that is most frequently affected. The EM waves of infinite tesimal wavelength produce on the hand that they strike sometimes a sensation of heat, sometimes a sensation of cold; sometimes they cause stiffness in the tendons and muscles.

This is how with a little practice some doctors manage to distinguish at first sight by the sensation of the particular radiation which escapes from it a sick organ from a healthy organ and end up becoming very expert in this kind of discrimination. .

Let us seize in our right hand a fairly large steel rod ending in a fairly slender point. Let us direct this point towards the wrist of our left hand, but in such a way that it remains contained in a vertical plane oriented either NOSE or NESO We will experience, after a few seconds, at the point struck by the radiation of the point, a very clear feeling of cooling which will make even more sensitive the feeling of warming which will follow it, when we will definitively remove the point of the left wrist. If, on the contrary, we let its action continue for some time. we will note a certain stiffness of the muscles of the hand, stiffness which, after having reached the forearm, will extend little by little from the elbow to the shoulder.

Some people very easily distinguish the full from the empty, with the only presentation of the hand and recognize not only the presence of underground cavities, but also the different depths of certain of their parts.

* What is Dowsing ?

From the word dowsing, each, following. culture and temperament, has given a definition which, most often, diverts the term from its etymological meaning, which is none other than sensitivity to radiation (from the Greek word *aisthesis* which means sensitivity). We will therefore allow ourselves to propose the one which, taking into account all that precedes, seems to us the most logical and the most in conformity with the reality of the facts, by saying that dowsing is the faculty of perceiving the radiation of matter. The term dowsing therefore becomes absolutely analogous to those of vision, hearing, touch, faculties to perceive smells or tastes.

Just as we call optics and acoustics the branches of physics which concern the study of perceptible phenomena with the help of the senses of sight and hearing, so it will be logical to call *fadiaesthetics* the branch of physics which is in the process of being born, and which concerns the study of perceptible phenomena using the impressions of the sixth sense.

It is by placing ourselves at this point of view that we will expose the summary study which is the subject of the first chapter, before beginning that of the practical re-education of the sixth sense which forms, for us, the main object of this manual and which will form the subject of the other chapters.

- The dowsing phenomenon

Having fixed, insofar as it has been possible for us, the nature of the external sixth sense, we are all the more at ease to define the dowsing phenomenon such as we see it produced with the strongly diminished organ of the normal civilized. As we exposed it in the work : " The Dowsing explained and practiced by the Physical Method", page 14, this phenomenon is not simple in the true sense of the word, but comprises a succession of three facts:

1^o A physical fact, external to the operator and objective, which is none other than the emission of a radiant source, radiation of matter of an infinitesimal order of magnitude; our human receiving apparatus being as passive and as unconscious as possible.

2^o A physiological and subjective fact, It is the shock of the emission of this source on the attenuated organ of our sixth sense; and, as

a consequence, the birth of a reflex, as well as that of the subsequent movement that it determines , a movement amplified by the detector (pendulum or rod).

A reflex, by itself, is unconscious, but its effect is not always so; and, in the case of the ray phenomenon , this effect, movement of the rod or of the pendulum, gives us the consciousness of a sensation of the organ of touch, which constitutes the spring perception.

3° A psychic fact. It is a second intervention of our intelligence which, by deducing from the physiological fact the existence of the physical and objective fact, comes to give to the second fact the interpretation that reason seems to dictate. When the role of intelligence is limited to the sole observation of the relationship that exists between the first and the second fact, it is because the operator has let the cycle occur automatically, it is because he has adopted the passive attitude, that which is appropriate for the processes of dowsing by the Physical Meihode.

But it is beyond doubt that, in the perception of the dowsing phenomenon, the soul can intervene actively, in particular by the exercise of the will. The operator will be able, for example, to adopt the selective attitude specific to the processes of dowsing by the Mental Method ; that is to say that, in a cloud of different radiations, he will only allow his organ to be receptive to those coming from a determined radiant source, of which he will have mentally made a choice, excluding others. His attitude will be absolutely comparable to that of the listener of a concert who, of all the music of an orchestra, will force himself to hear only the notes of a determined instrument. (We

will notice, on the other hand, that if the selection of sounds is possible, it is the same for that of smells and tastes). ■

It is the exercise of this faculty that Brother Padey calls the use of the “fluid of intention” and that other dowsers designate under the name “call to mental orientation ”.

The selective attitude, in the cases where it is essential, must precede the operation of research. You have **to** separate before grabbing.

Finally, there is a third attitude on which we will not insist in this manual. because, seeming to require special faculties, it does not appear to be of any interest to the beginner; it is the exercise of mental **concentration**.

Certain dowsers, in whom psychic intervention is preponderant, have denied the possibility of perception in the passive attitude. But this

opinion is contrary, not only to the laws of psychology, but above all to the facts themselves. How many times does it not happen to hold between the fingers a pendulum adjusted to the radiation of a determined substance. Suddenly, in the middle of a conversation, when we are no longer even aware of having it in our hands, the instrument begins to spin. Looking for the reasons for such an unexpected fact, we find that the pendulum was actually on the vertical of a substance, identical to that for which it had been adjusted. The human receiver had behaved like an absolutely automatic machine ! • •

When two neighboring operators have in their hands a pendulum set for the same substance, yes. one of the two devices begins to turn, the other inevitably follows the same movement, by the fact of a sort of resonance, by synchronism, or of a transmission of reflexes analogous to the fact of the contagion of yawning in society .

. Passivity can be unconscious and involuntary; in this case. it is absolute, as in the two preceding facts . But she. can also be imposed by will; in this case, it is only relative, and comes all the closer to absolute passivity as the subject has more empire over himself.

Is dowsing an art or a science?

If we keep the word dowsing in its etymological sense, or if we give it the definition we have chosen, there can no more be scientific dowsing than there can be scientific hearing, scientific vision, scientific tasting. . We have more or less fine hearing, more or less sight. less piercing, I palate more or less educated or refined, but one cannot have the eye, the ear or the palate scholars.

On the other hand, Radiaesthetics, the study of the phenomena revealed by the perceptions of the sixth sense, is a branch of physics, and, consequently, a science of observation . •

For the same reasons, dowsing cannot be an art; but the dowsing technique, collection of the most advantageous processes to facilitate or amplify the dowsing perception, is, incontestably, an art. It is not enough to be sensitive; it is also necessary to know how to make the most of the exercise of this faculty.

The objections' against Ja Dowsing:

Like all novelties likely to disturb vested interests, it was not possible

that at its beginnings, **dowsing** did not provoke violent reactions. That's what happened. indeed. .

We will not return on a question which was treated in the work : “ The Dowsing explained and practiced by the Physical Method” with all the necessary width; but **concerning** the writings published on this subject, we will allow ourselves to add a simple word:

To be able to render a fair judgment, the judge must be impartial, that goes without saying, but, above all, competent. To deserve to be heard, it is essential that whoever speaks or writes knows perfectly the

question he treats, that his knowledge is not only the result of the study of books written on the subject, but indeed an experimental knowledge. "It is by forging that one becomes a blacksmith". What would we say of a person blind from birth who would like to persuade us that colors do not exist? What would one think of a sufficiently re-educated deaf-mute who offered to give a lecture on the works of Mozart or Chopin? To condemn a priori the thesis radiates, without having tried, beforehand, to do . to turn a rod or a pendulum is to attempt to achieve such a prodigious tour de force.

Because, finally, as Marshal Foch would say : " What is all this controversy about? It is a question of knowing if, yes or no, the dowsing sense exists. This is the real, the only question that arises. All the others are accessories. How to solve it if one is not a dowser? If we have only acquired a bookish idea of it? Why refuse to experience it personally ? We are not **asking** to "**believe.**" We ask to "know" by experience.

Those who want to evade it come to tell us : " There **are** dowsers who make mistakes". We will answer: Which doctor has never made a diagnostic error? Others will add: "There **are** dowsers who, when they talk about science, say terrible nonsense". And when would that be? You play out of tune on an instrument. You're a bad musician, that's all. But no one would dare to claim that your instrument does not exist, nor that others more skilful than you will not one day be able to draw wonders from it.

As the railways started **there** . hundred years, the dowsing begins at this time. Those who look at the marvelous ultra-modern machines now in circulation find that they only remotely resemble the locomotive of the train in which Dumont d'Urville lost his life.

When it will count the same number of years as the railways, who will be able to count the immense services that Dowsing will have rendered to **humanity** ?

What would have become of the sensational discovery of the real movement of the earth without the calm contempt of the physicist Galileo, who went so far as to be denounced a heretic by the Inquisition in 1633, thus coming up against general incomprehension?

What would have become of one of the most beautiful discoveries of modern physics, dynamic electricity? without the reckless courage of the physicist Galvani, in the face of the skilful calumnies of his detractors in 1780 ?

What would have become of the marvelous discovery of the

TSF without the energetic and ardent convictions of the engineer Hertz, on the propagation of electric waves. overturning prejudices and routines and sparking its victorious spark in 1885? -

The great Pasteur himself, a true revolutionary in the art of healing, in 1890, was translated "as accused" according to the expression of Brouardel, at the tribune of the Academy of Medicine "when he had to hear that his discoveries were unscientific. ineffective, dangerous and that they constituted a double danger: social danger, homicide; — intellectual peril. unreason ! »

Let us currently leave the spirits that the Dowsing grieves to write in peace the pages which relieve them and let us pass... but not without having noted that if the Dowsing still counts detractors, it also counts among its defenders of the illustrious names.

Among these, let us mention the glorious Edouard Branly. who accepted until his death the title of Honorary President of the International Association of Friends of Dowsing, and MM. Deslandres and d'Arsonval (the latter recently deceased). both members of the Institute, I a current Honorary President, the second Honorary Member* of the French Scientific Association of Dowsing.

FIRST CHAPTER

WHAT IS RADIESTHETICS

Many definitions of science have been given. Some have called it the exact and reasoned knowledge of a set of facts of a well-determined category; others have been able to say with more accuracy that science is an effort to reduce the fleeting complexity of facts to the stable unity of law. Science is only science, in fact, on **the** condition of disengaging itself from particular facts and establishing general laws.

This being of course, the sciences have been classified into three categories:

1 ° Mathematical sciences or exact sciences whose acquisitions are based on deductive reasoning.

Mathematics, or the science of measurable magnitude or of quantity considered in the form of numbers or figures, is the simplest of all sciences because one only has to reflect in order to grasp the true relations of the ideas that she presents.

2 ° The natural sciences, the development of which depends exclusively on the observation of facts and induction. It is in this second category that physics, or reasoned knowledge of what we can know of the external world, falls into.

3 ° The moral sciences which hold the middle between the two.

Our five known external senses can be considered as windows opening onto the external world and enabling us to receive from it five impressions of a DIFFERENT nature. It is the senses of sight, hearing and touch which have made it possible to construct the branches of classical physics which have been called: optics, acoustics, heat, electricity, magnetism, radioactivity. It is obvious that if we really have a sixth sense and if **we** manage to put it into action, it will allow us to perceive new aspects of the external world and will reveal to us, between the realities which escape us, the existence new reports still ignored. Now, as there is no doubt that this sixth sense exists, we can say that it is a new branch of physics, the development of which we will soon be able to see: and this branch, by analogy with the acoustics and optics, we will call it the radia-

thetic. This term, in fact, seems to us more proper than that of scientific dowsing, by which some have designated it up to now.

Among the practitioners who have dealt with scientific dowsing, some have forced themselves to undertake their research only in the exclusive domain of the knowledge of official science and to ruthlessly reject, as false, everything that does not seem to them a confirmation or addition of what they knew.

do so is, in our opinion, to condemn oneself to eternally trampling on the spot. As, as Pascal said, we shall never know the whole of nothing; just as, on the other hand, the more man advances in the study of the sciences, the more he realizes his ignorance, it must be admitted that present-day physics, chemistry, biology only know 'particular cases of a much more general science, barely suspected. To persist in wanting to attach everything to these particular cases is to fall into the error of a narrow conformism, the immediate effect of which is to oppose all progress in the natural sciences, in perpetual evolution.

We will therefore deviate from a dangerous pitfall which would make us sink and, making as far as possible a clean slate of all that we know, we will quite simply seek the truth by following the path which leads to it in the natural sciences, c. ie using the experimental method.

Experience being the source of all truth, it is to experience that we will ask to provide us with the knowledge we seek. The experimental method to which we will have feedback will consist of :

1°) A. observe and note the facts.

2°) To formulate a hypothesis that we will seek to verify as soon and as often as possible.

3°) To generalize, that is to say to pass from the particular to the general and to establish laws.

4°) Finally, to deduce, that is to say to pass then from the general to the particular, and, from the established laws, lead to the existence of facts which will inevitably have to occur if these laws are correct.

If official science repudiates the inadmissible connections that several dowsers have allowed themselves, if it vigorously protests against the borrowings they have made from its vocabulary to designate facts quite different from those it has in view, it is with the most legitimate reasons.

Several phenomena met with in dowsing, although having a common cause with facts acquired and known for a long time, are effects absolutely distinct from them. It is therefore essential, at the risk of causing chaos and confusion, to designate them by the proper denomination that suits them.

It is to this rule that we will adhere strictly in the following account, an account which we have tried to make as simple and as elementary as possible, in order to put it within the reach of all our readers.

§ 1 — Microphysics

. As long as the dowser works with reflection, either in the laboratory or outdoors, he quickly realizes that his pendulum puts it in contact with all the forces of nature, studied in the treatises of physics: gravity, heat, electricity, magnetism radio-activity',

light: but it also realizes that these modes of energy are presented to it, more often than not, with an infiniteesimal order of magnitude, the measurement of which escapes the most perfected apparatus. These are, for example, weights of a millionth of a milligram, electro-motive forces of a ten-thousandth of a millivolt, intensities of electric current of a thousandth of a micro-ampere, radioactive intensities of a millionth of micro-curie, etc., etc. The whole of the phenomena which result from the bringing into play of these infinitesimal forces constitutes what one can call **microphysics**.

At the same time that he is impressed by this very **clear particularity** presented by the manifestations of forces of nature which are familiar to him, he finds himself in contact with new facts, completely unknown to physicists, because nothing has **allowed** these, up to now, not only to seize them, but, even, to suspect their existence.

The study of the phenomena of microphysics, which have hitherto only been revealed by the perceptions of the sixth sense, constitutes **dowsing**.

§ 2 — The radiation of matter

Among the waves in general, we retain essentially two kinds of radiation: electromagnetic waves; corpuscular radiation.

1 ° Electro-magnetic waves : We will not go back here to the considerations relating to electromagnetic (EM) waves. One will find them sufficiently developed in chapter II, page 48, of the treaty of “Dowsing by the Physical Method” of P. BÉASSE.

Just as audible waves, called sound (result - of a vibrating body) move in a solid, liquid or gaseous medium (generally air in the latter case), so electromagnetic waves propagate in the interstellar elastic medium, called ether, which fills all spaces and penetrates all bodies.

Are electromagnetic waves the result of the vibration of ether, an elastic substance, the nature of which we are completely ignorant of? Could this vibration cause the formation and propagation of electric and magnetic fields ? Simple assumptions.

Among the electromagnetic waves in general, let us note the existence:

a) electromagnetic waves specific to simple bodies (compound bodies emit waves specific to each of the simple bodies which enter into their composition). All these specific waves are of very short length (a few millimeters for small masses); _

b) waves of the terrestrial electromagnetic field, which seize and transport the specific electromagnetic waves of the aforementioned simple bodies and serve as their support. It is these waves of the terrestrial electromagnetic field which will allow the specific electromagnetic waves of simple bodies (of the animal, vegetable and

mineral kingdoms) to propagate over long distances.

These support waves are, depending on the case, electric, magnetic, luminous, nervous, etc.

Electro-magnetic waves, called " specific " Returning to the specific electromagnetic waves of simple bodies, we will point out:

a) that polarized electromagnetic waves (positive or negative) have two planes of vibration: one electrical, always vertical; the other magnetic perpendicular to the first, and passing through the ray of propagation, which thus happens to be at the intersection of the two planes;

b) that when the wave propagates vertically, its magnetic plane of vibration is always oriented in a fixed direction (direction of the fundamental plane). The angle that this plane forms with that of the magnetic meridian taken for origin, is called the fundamental angle of the wave;

c) that each wave is characterized by a special number, which appears in numerous phenomena, a number which is called the series objective number or particular number of the wave (and which itself depends on another number, called ' characteristic number with which it is linked by a very simple mathematical relationship: $N = 2n + 1$);




Fig. 1
Rayon vertical




Fig. 2




Fig. 2 bis
Horizontal radius




Fig. 3

d) that by adding to these data the wavelength of the wave, if it has been possible to determine it, and several millimetric adjustments of the pendulum suitable for it, one can establish a sort of identity card which makes it possible to recognize and identify the wave in all circumstances;



e) that the polarity of a wave depends on the direction of the orientation of its magnetic field with respect to the corresponding electric field. Two identical waves, but of different polarity, are offset from each other by half a phase and have their fundamental angle at 180° from each other. ¹

It is probably under the effect of the planetary atomic gyration that the bodies emit the special electromagnetic radiation, discovered by the engineer Voillaume and which is identified with that which the ancient dowsers called "effluvia". This radiation, which he called : specific radiation of the body, characterizes each substance.

Of very short length, as we have just indicated above, this radiation would only be perceptible at a very short distance, if it were not seized and carried away by the waves of the terrestrial electromagnetic field which serve as its support . .

This radiation is not unique, it includes a beam of EM-radiations of different wavelengths, but which have the particularity of possessing the same tondamenral angle, the same particular number at the same time as several common physical properties such as the faculty of walk through certain screens, or be stopped by them.






Fig. 4



Figs. 5



Figs. 6




Fig. 7

Fig. 8





Fig. 9




Fig. 10




Fig. 11




Fig. 12

Thus, under the action of the terrestrial electromagnetic field, a body or object emits : .. .

1°) In both directions (heaven or earth), specific vertical electromagnetic rays, very short, with an ES support (waves of radiant electricity);

2°) In the directions at 45° and 135° with respect to the vertical, specific electromagnetic rays inclined, very short, with MS support (radiant magnetism);

3°) In the horizontal direction of specific electromagnetic rays, of limited length, with eu support (waves emitted by magnetic fields);

4°) In a fixed diredion, characteristic of each • substance, a beam of specific electro-magnetic rays, very . short, with MS support (**Plane Phenomenon**

fundamental) ;

5°) Under the combined action of light and the terrestrial EM field, and in the direction of each of the four magnetic cardinal points of the object, a beam **of** very short specific rays, with MS support. This radiation constitutes the magnetic cross of the object.

We pass over in silence the "particular ray", described on page 87 of the treatise " Dowsing by the Physical Method", because, without practical interest from the prospecting point of view, it seems useless to encumber the baggage with it. of the beginner. >

2. Corpuscular radiations : There also exist in nature material corpuscles which are infinitely small but heavy and endowed with energy. Their displacement in a cloud or in a beam constitutes a corpuscular radiation.

The discoveries of modern physics have shown that these corpuscular radiations are formed of corpuscles. transported by electromagnetic waves, particular

for each of them. It is these waves which communicate to them the movement, without which they would remain motionless.

biles, and which Louis de Broglie calls the wave associated with the corpuscle.

Among these corpuscles, let us quote:

- a) Photons transported by light electromagnetic waves and whose impact on our retina produces the sensation of light;
- b) The Electrons transported by negative electromagnetic waves ES and which constitute, either the electric current, or the rays γ of the radioactive bodies;
- c) The rays α : radioactive bodies, atoms of helium, transported by the specific rays of this gas.

The corpuscle and the wave being closely associated, the movement of the corpuscle and the propagation of the wave are not independent (1) ; and, to the mechanical quantities (quantity of movement and energy) of the corpuscle, are linked the characteristics of the associated wave (speed of propagation, wavelength and all other elements which, from the dowsing point of view constitute its map of identify).

- The dowser therefore does not recognize a corpuscular radiation only by identifying the wave associated with the moving corpuscle. When the quantity of movement of the corpuscle cancels out, the wave is damped.

One easily distinguishes an EM wave from a corpuscular **radiation** by the fact that while the first is picked up instantaneously by an antenna placed at a suitable distance from the beam of propagation , the second remains absolutely insensitive to the call of this device. The wave closely associated with the corpuscles follows their path without being able to be either dissociated from it or diverted.

The example of rays. α , helium atoms, whose associated wave is none other than the specific radiation of helium; that of the electron, whose associated wave is the ray ES, shows that what physicists have named **the associated wave** is identified with the **specific ray** of the α corpuscle. It is the corpuscle which produces it and naturally follows it.

It accompanies him everywhere, since it is he who gives birth to him. She has, of course, no influence on the movement of the corpuscle which is due to an external force.

§ 3 - Electricity

One of the first acquisitions of microphysics carried out by the way of the dowsing processes, was the discovery of ES waves, or waves specific to electricity , which constitute what some call radiant electricity. These waves, with either positive or negative polarity, permanently escape from any static electric charge .

(1) Louis de Broglie : “ *Matter and Light* ”

• • □ ii □ i ■ □ □ ■ • f
 tick or any conductor carrying a direct current
 - or alternative. They constitute the waves associated with the electron or
 the positron (1) . Their wavelength, 11 microns 363, which the engineer
 Voillaume determined in 1932, places them in the infrared region of the
 spectrum. For several years, their existence seems to have been recognized
 by. dfe American physicists, under the name of extra-short waves emitted
 by lighted incandescent lamps, and their action then used, across the
 Atlantic, for various effects .
 fJerapeutics.

By a mechanism which space does not permit of describing here, these ES[^]
 waves constitute the active agent of the phenomena of electric induction. They are
 also the same waves, which export vertically in both directions, and, at dis- .
 considerable tances, the specific waves of all bodies (with the exception of zinc),
 waves which, reduced to their own energy, would not exceed a range of some
 a few millimeters. ~

But, if these ES waves escape from an electric charge, on the other hand, we
 find no trace of their emitted

sion in an electric **field** , unless provoked there artificially by processes of
 induction.

The electric **fields** emit other electromagnetic waves of a wavelength which
 seems infinitesimal and of the order of that of the X-rays , waves of which it was
 possible, after having isolated them, to determine the characteristics. These are the
 rays (**J** with double polarity, positive (J 7) in the ' positive electric fields;
 negative (fj —) in the negative electric fields. ' '

Ij rays of limited length stop at the extreme equipotential surface of the electric
 field, which makes it possible to trace their contour using the pendulum in a chosen
 plane, for example horizontal or vertical.

EM radiations in electric fields . of shorter length and which are damped on
 intermediate equipotential surfaces. .Those are

„UJ *Positon*: Name that Louis de Broglie gave to the positive electric **corpusu** !e positive
 electron . At **first it is called a positron**. The electron e6t • the negative electric body. •

α and β rays of positive polarity in positive fields; the rays γ and δ of polarity . negative in negative fields.-




Fig. 13

The rays γ , δ , and α are emitted by the electric field which the phenomenon of disintegration determines in radioactive bodies, and, as we shall see further on, they are encountered in the beam of harmful telluric rays. which escapes from certain parts of the ground.


§ 4 — Magnetism

The MS waves whose wavelength, also determined by the engineer Voillaume (10 microns 787), places them in the infrared, play, with regard to magnetism, a role which seems analogous to that ES waves with respect to electricity. Could they be, like the first, the waves associated with a corpuscle analogous to the electron and whose existence official science still ignores ? It's not implausible. They escape from any magnetized body.

The terrestrial globe seems permanently traversed by a rectilinear emission of MS rays parallel to the magnetic meridian. This emission appears to correspond to the geometrical system of radiations discovered and described by Doctor Peyré.

Under the same conditions as ES rays, waves

MS export from all bodies specific rays which escape under inclinations between 45° and 135° ; the extreme rays of this zone being by far the most intense. '



Distribution of radiation θ , O. L. and M. in a magnetic field developed by the charge of a rectilinear magnet

The MS rays therefore constitute the emission waves of the magnetic charges (1). They are found in a magnetic field only if their

Fig.14

emission is provoked there by an induction process.

But if the magnetic fields do not emit MS rays, they emit, on the other hand, w rays with always positive polarity and o rays whose polarity is double. corresponds to that of the field from which they emanate.

The rays <=>, of limited length, stop at the extreme equipotential surface of the magnetic field, which makes it possible to trace, using the pendulum, the outline of this field in a chosen plane, for example, a vertical plane. or a horizontal plane. The rays o of shorter length stop at an intermediate equipotential surface.

The classic physicists' magnet consists of a bar of steel. The magnetic charges there appear concentrated in two points situated near the extremities of the bar and which are called poles. The part between these two points is called the neutral zone. However, this area is not completely devoid of magnetism, since the pendulum, set to MS rays, rotates along the entire length of the bar. In reality this very weak charge, after the pole, decreases more and more until a neutral line, from which it changes polarity.




Fig. 15

All bodies, some to a measurable degree, some to an infinitesimal degree, possess a charge magr . etiquette. The iron objects form a single magnet with two poles. But

(1) Per • charge ". ll In ut hear the amount of magnetism or electricity contained in a body. •


By " field " the region subject to. the influence of a magnet (magnetism) or a current (electricity).

Objects. of any substance other than iron. consist of two magnets placed side by side.

. We give below the example of a silver spoon on which have been indicated the location of the poles as well as the layout of the neutral lines of its double magnet (Fig. I 6) .

When an object has a point, one of its magnetic poles is generally located near this point.

To finish, let us say that the rays (**tt**) escape from all the points and from all the extreme parts of the elongated bodies carrying with them electricity and specific rays of these bodies, rays to which they serve as supports.



Location of the poles and 'tracing of the neutral lines
of the double magnet of a silver spoon

Fig. 16

§ 5 — Radioactivity

For physicists, radioactivity characterizes the property of a body to continuously emit α and β **corpuscular rays** at the same time as EM waves, called by them γ *rays*. This emission is due to a phenomenon of atomic disintegration and transmutation which takes place continuously but extremely **slowly**.

By the action of a **negative magnetic field** produced by a fragment of pitchblende, for example, it is easy to separate the different radiations of a radioactive body. The α rays then escape into the fundamental plane of the substance such as it is oriented after the deviation which the magnetic field causes it to undergo; and the rays β in a plane perpendicular to the previous one. The rays *there* continue to escape vertically.

The α -rays **are helium atoms**, the path of which can be followed with the aid of a pendulum adjusted to this gas; the radii of the electrons, of which an adjustment on radius ES makes it possible to recognize the path.

A positive magnetic field is without any action.

If one carefully studies with the help of an impregnation on paper the radiation of a radioactive body, one realizes that it is surrounded by a negative electric field, itself prolonged by a magnetic field. negative tick. • ■ .

Figure 17 below shows the result of the study carried out on a fragment of pitchblende.

The negative electric field within which the body itself is comprised, emits β and γ **rays**. As a result of an anomaly, the cause of which has not yet been found, the rays (α which, associated with the rays β and γ should be of negative polarity, are positive in radioactive bodies.

To the rays emitted by the electric field is added a fourth radiation, the RS rays discovered by the engineer Voillaume and which really constitutes the specific radiation specific to radioactivity. It is found on living beings. The set of four rays, RS α + β and γ waves (EM waves of radioactive bodies), undoubtedly constitutes the group of physicists' γ **-rays**. It should be noted that in a study published in 1926 by M. F. Honoré and edited by the Gauthier-Villars bookshop, the author already mentioned several rays of different wavelengths.

Radio-active radiation from bodies : Physicists who have occupied themselves with the fascinating question of radio-activity have taught us that if radioactive bodies

1. et des ondes électromagnétiques dénommées rayons γ et radia-

emit corpuscular rays




Fig. 17

. Antenna-tip examination
of a pitchblende fragment

Suppose it was easy to separate from each other **using** a negative magnetic field, **all** non-radioactive bodies also emitted this same radiation, but to an infinitesimal degree.

1.4 dowsing allows us to make, with the greatest facility, the observation of this radioactive radiation of all the bodies, which is completely distinct from the radiation **RS**, of which Mr. Voillaume discovered the laws.

Without it being necessary, this time, to resort **to** the action of a magnetic field (because, because of the weak intensity of this radiation, the terrestrial magnetic field is enough **to** separate the elements), we will note, with appropriate settings, that the **rays** emitted

by a substance, escape in the fundamental plane of this substance and the rays $\wedge$ in a vertical plane perpendicular to the direction of this plane.

If we study, for example, the radioactive radiation of a silver coin, we find that the '**J rays**' go in the direction of magnetic East and the $\wedge$ **rays** in that of magnetic North.

From all that precedes, it follows that, considered from the microphysical **point** of **view**, all bodies are electrified, magnetic and radioactive .

. § 6 — Light

If we know very little about matter, we are even more ignorant about light, many of whose effects still remain completely mysterious to us. By the additional perceptions which it brings to us, the dowsing will be able to come to our assistance, and to

contribute, probably, to 'give us the solution of certain disconcerting problems.

The most recent works of modern physicists teach us that the cause of light reaches us in the form of the shock of corpuscular radiations.

Infinitely small material bodies, called photons, which it shoots out with considerable speed, are associated with unpolarized EM waves called light waves. If it is correct, as we recalled in § 4 (page II). that there is a link between the mechanical magnitudes of the corpuscle and the characteristics of the wave associated with it, each shade of color would correspond, it seems, to a different energy of the photon, which would determine the length of wave different from the light wave; and, consequently, a different impression on our retina.

The number of shades of color is not infinite, but limited; because, in the dispersion spectrum, each corresponds to a homogeneous line of a relative width. White light constitutes the impression made on our retina by the composition of the elementary impressions determined by each shade of color.

As physicists taught us more than a century ago, light waves are accompanied by calorific waves (CL rays with negative polarity) which produce heat, and chemical rays (CH rays with positive polarity). which play a - considerable role in photography.

Accompanying Rays of Colors : The Spectrum. dispersion of white light is clearly divided into two very distinct zones, separated by the shade of medium green: an electric zone and a magnetic zone.

In the electric zone which extends from this shade of green to the extreme violet, and, even beyond, in the ultraviolet, the light rays are accompanied at the same time as CH rays, ES rays presenting frequent polarity alternations. These alternations determine a large number of subdivisions separated by neutral shades on which a pendulum set on ES rays does not react. In the positive subdivisions, the rays visible or not, are accompanied by rays $J + e$ and y and, in the negative subdivisions, by rays $0 \rightarrow$ and $\wedge$

In the magnetic zone, the light rays are accompanied at the same time as calorific rays (the effect of which, at first imperceptible, only begins to become noticeable from the yellow), MS waves whose alternations of polarity look perfect


These luminous rays are 'accompanied
by specific waves of Magnetism.

Magnetic Zone

These light rays are accompanied
by specific waves of electricity.

Electrical area

N+	n- _	n+	11 —	E +	E -	E+	E -
----	------	----	------	-----	-----	----	-----



Spokes MS 6 polaritt alternoti...

Alternate polarity ES spokes

Fig. 18 COLOR ACCOMPANYING RAYS (I)
on a solar spectrum bringing together all the colored rays
resulting from the decomposition of sunlight

(1) Recent experiments have shown that color alternations are much more frequent than is indicated by lo tn b Ica u do la flg , 18. Thus , there are two shades <indigo, uno positive, other negaUvo : do môme, two shades of o rang é do po lar i 16 illtréj-cnte. We are nevertheless reproducing the table which appeared in our last edition because it gives in an intelligible way the other general nature of the phenomenon of the polarity of colours . Mals It must be well cn teitdu For our readers , the parts of the table must themselves include several subdivisions . We will give more details in our next edition.

concordance with those of the previous zone. In the positive subdivisions, the light rays are accompanied by waves w of Q^+ ; and, in the negative subdivisions, of $\{\bullet\}$ and Q^- WAVES, that is to say of the same waves which are found in magnetic fields. The magnetic zone extends from medium green to extreme dark red and extends beyond that into the infrared. The number of alternations do. polarity in the visible part of the spectrum appears to be about fifty.

Physiological action of the colors : For several years already, several dowzers, Mr. Mellin, in particular, have pointed out the great physiological influence of the colors, a pernicious influence in certain cases, beneficial in others. There is no doubt that the different energy of the photons, the different wavelength of the associated wave, the nature, the polarity and the intensity of the accompanying rays are all factors which react on the vibratory movement. of the cells of living beings and, consequently, influence their health. Hence the principle of chromotherapy.

Colors can also act on the brain and, consequently, on the mental state. A person accustomed to intellectual work said to have noticed that by covering her head with a yellow scarf she felt her lucidity and her intellectual power considerably increased.

It goes without saying that the color capable of producing such an effect varies according to the person.

§ 7 — Harmful telluric rays

Doctor Peyré checkerboard

It is no longer disputed today that certain areas of the land emit radiation harmful to the health of living beings, by the disturbance they bring to the normal vibration of the cells they reach.

We will only mention here the rays which normally escape from certain regions of the ground, deliberately leaving aside any other radiation emanating in particular from microbial cultures, which would find themselves accidentally mixed there.

For reasons which will be explained, the **harmful zones**, sometimes of very small dimensions, are extremely widespread. The intensity of their emission, very variable, moreover, is, fortunately, not always sufficient to cause appreciable disturbances. It is therefore, above all, **areas of intense radiation** that we are going to deal with.

Harmful radiation is revealed by the more or less rapid, more or less accentuated uneasiness that it causes in most people staying in the irradiated zone, by the observation, in certain fractions of the ground, if not of a total absence, at least from a serious attenuation of the vegetation.

Nature of harmful telluric radiations : If, after exposing a sheet of white paper for a few seconds to radiation from a harmful zone, it is studied in the laboratory, it is found that this paper is impregnated with several distinct radiations, including a selection method allows to determine the characteristics and • the respective description. These are, invariably, negative ES and MS waves (ES—, MS—): then positive (**J rays** ((j + J rays ; 'i and £ rays: RS rays, that is to say the EM waves which normally escape from the vertical of any **radioactive body** .

ES and MS rays not being harmful, *n* harmful are therefore nothing other than rays further away in les radiations telluriques I. Nous verrons un peu

s'échappent d'une fraction radio-active du so

which cases, the **y rays** can be accompanied by other equally harmful radiations, the **p, - rays**; and **x**. (See page **98**, pendulum adjustment table on each of these waves).

(

Source of harmful rays. — The emission of harmful rays can come from several radiant sources:

1. Generally quite extensive region in which the composition of the soil includes **radioactive mineral elements**, the presence of which, because of theirⁱ⁻ infinitesimal dosage, escapes chemical analysis.

mic. This is the case for certain marls, certain ochres. There is no doubt that living in or staying on such land for a long time can only be detrimental to health.

2° Underground hermetically closed or communicating with the atmosphere only in a precarious way: this is the case of a natural underground cavity, cave, cavern, crack in a fault, or of a cavity established by human hand, closed shaft, gallery, tunnel. sewer, **etc.** . .

As it is easy to see in the laboratory using a stoppered bottle or a tightly closing cardboard box, **the air confined** in an enclosure ionizes and **becomes radioactive**.

A hermetically sealed underground therefore emits the radiation of a radioactive body. If its bottom or its walls contain ores of radium (**I**), thorium, actinium, the ionization of the air causes the formation of gaseous emanations discovered by Rutherford, subsequently called radon, thoron, actinon, and whose specific rays come out of the underground at the same time as the **y rays**. To be sure, in the laboratory, it suffices to put in the flask or in the experimental cardboard box, a fragment of ore containing the aforementioned elements. After a few seconds, the emission of the **p, - rays** will be observed ; and **x**. which are also harmful.

„

„

CI) It is only by analogy that we cite actinium, the experiment having been made up to the present. than on radium and thor um.

3° Subterranean stream passing through a cavity or radioactive. Steeped in their show, the stream carries tions and emits them vertically un terrain throughout its course. ces radia-

Subdivisions of the clcmc-nhs of the checkerboard : Each élément d u

4° Rectangle du damier du Docteur Peyré.

Damier du Docteur Peyré : « A la surface du globe, dit le Docteur Peyré, on trouve -partout, et d'une façon géométrique, des rayonne ments telluriques .parallèles ou perpendiculaires ou méridien mc?.gné tique et formant un damier de 8 mètres environ de côté entre axes».

Sous réserve des dimensions des éléments du damier, qui peuvent varier légèrement d'un lieu à l'autre suivant les influences locales, on peut, après avoir tracé à l'aide du pendule, la figure géométrique sur le terrain, constater que les lignes Nord-Sud sont porcqurees par des rayons MS et les lignes Est-Ouest par des rayons ES, **accompagnés de rayons** ..! . Le sens. de propagation et la polarité des rayons ES et MS, qui

ne sont pos verticaux. mais inclinés, paraissent al- terner brusquement toutes les douze heures.




Fig. 19

L'émission des rayons nocifs qui est, normalement, peu intense, devient, ou contraire, parfois extrême- ment violente aux points de croisement du damier. En ces points, ("émission •des rayons GiES et MS s'arrête complètement pour ne laisser subsister qu'une émission de rayons y. Sont qu'on puisse s'en expliquer les raisons, l'intensité do cette croisement. à l'autre. La nocivité

émission est extrêmement variable d'un

d'un rectangle peut être très bénigne, alors que celle voisin peut du rectangle et être extrêmement préjudiciable à la santé, provoquer lo notamment formation du cancer.

checkerboard itself comprises several secondary divisions determined by bands traversed by the same rays and whose crossings enjoy the same properties as those of the extreme bands of the element.

Positive ground : The earth's electric field being **negative**, these are ES- rays which escape from it everywhere; However, these rays are generally of an intensity much too weak to impress a pendulum adjusted to this radiation. It was M. Orcet who **first** observed the existence of zones of land of small dimensions

emitting with a more or less high intensity **positive ES rays**. **11** has named these grounds positive grounds and pointed out their extremely beneficent influence on vegetation. He recommended planting trees only on positive ground. It is easy to realize that the recommendation is excellent, because any tree living on such ground thrives in a remarkable way.

If we study in detail a Peyré checkerboard element, we find that each crossing is flanked by two smaller rectangles, c=onfigus, emitting ES+ rays.

Ter **Rectangl** **Ter**
TG:n **e - Dow!** **rain**
•pCI **•** **posi**
Sihf **(JAUNE)** **hl**
•faITl **(vfr**

Fig. 20

Croquis de la page 24 bis: Le croquis colorié de cette page représente un élément du damier du Docteur Peyré.

Les raies teintées en rouge représentent les bandes parcourues par des rayons MS; celles teintées en bleu, les bandes parcourues par des rayons ES; les rectangles jaunes figurent les croisements émetteurs de rayons nocifs, les rectangles verts adjacents les zones de terrains positifs.

On the element studied on site, given as an example, the extreme magnetic bands were spaced 7 m apart. **66** from axis to axis, the electric bands of **8 m. 60**. This element was subdivided by secondary bands parallel to the main bands. Among the secondary bands sufficiently wide and intense to be perceptible were **8** electric bands and **4** magnetic bands.

The width and spacing of the strips that have been scaled **to** the drawing. may vary slightly depending on the location, and perhaps also depending on the time of year.

Explanation of the Damier Pcyré phenomenon: The formation of the Peyré checkerboard is an effect of the Earth's electromagnetic field, as proven by the small experiment that we describe below and which allows the phenomenon to be artificially reproduced. On a table, let's put a small electric TSF capacitor, the negative side up. On this condenser, let us place a bar magnet as it is indicated on figure 21. We thus carry out an EM field analogous to the terrestrial EM field, at least such as it occurs in the boreal hemisphere (I).



On this assembly, let rest for a few seconds a sheet of white paper, the center of which is placed above the capacitor, and the major axis of the sheet oriented parallel **to** that of **Fig. 21** from the line of the poles of the right magnet.

(1) The North pole or positive pole of the compass being attracted "Towards the North, the boreal pole, of the terrestrial magnet, is of opposite polarity, therefore negative.

This done, let us study using the pendulum the fluidic impregnation of the radiation of the device- which is printed on the sheet of paper. These are the elements of this impregnation which are reproduced on the sketch in 4 colors of the Peyré checkerboard, page 24bis. We find there:

1° Bands parallel to the axis of the right magnet traversed by MS rays.

-,

2° Bands perpendicular to this axis, traversed by - rays ES accompanied by

rays "((RS, (j + , '1, £).


3° Crossings on which the MS and ES rays stop and which only let through "(rays of greater or lesser intensity.

4° Finally, in each element of the grid, subdivisions analogous to those of the Peyré checkerboard. ■

5° To these various radiations is added the emission of hydrofluoric streams or rivers. These rivers discovered and reported by Mr. Orcel, would be extremely numerous. They would flow from east to west and to a fairly large depth, from 80 to 150 meters. Their emission is all the more pernicious as it is generally unknown to dowsers .

While streams carrying y-rays (RS, 0 + .V, £) are simply impregnated with these radiations, fluorine in combination is found in a state of solution in hydrofluoric streams, at a dose absolutely infinitesimal that no chemical analysis is able to detect. But this "homeopathic" dose *suffices* in the long run to produce great ravages on health.

By a method, which would take too long to explain here, it is possible to measure the relative intensity of several identical EM waves. These are the numbers corresponding to this measurement for the radii 0 +



which were, on **the 4 -color** sketch of the Peyré checkerboard, placed next to each crossing rectangle. It will be realized that they are extremely variable from one rectangle **to** another, which explains why, in nature, the number of truly harmful rectangles is much less than one might suppose.

A ship with a steel **sheet hull** constitutes a magnet whose line of poles may not be in the axial plane of symmetry of the ship, although most often it is along its major **axis**.

As its installations comprise an electric charge corresponding to an electric field, it follows that the ship gives rise **to** an **EM field and produces a Peyré** grid identical to that which we have just described, and oriented along the line of its magnetic poles .

The observation of this fact **was** made by Doctor Peyré himself who pointed it out.

The same phenomenon occurs in a metal railroad car, tank truck, or motor car, ote ...

§8. — Infinitesimal Biology


If there exist, in inanimate nature, a large number **of** phenomena which develop an imperceptible energy and the knowledge of which constitutes what may be called microphysics, it is indisputable that manifestations of a weakness of intensity of the same orders of magnitude also occur in living beings.

Their knowledge constitutes an infinitesimal • **biology** that could have been called microbiology, if the term had not already been chosen to designate a completely different science: that of rudimentary micro-organisms. This biology, which is that of the future, and which is linked to radiesthetics, will make it possible, with the assistance of the sixth external sense, to pierce many mysteries of organic life (mode of vibration of cells, laws of their development, mechanism of external influences which act on this development, mechanism of vegetal tropism I, etc..., etc...).

We shall content ourselves, in this chapter, with making a discreet borrowing from this science, confining ourselves **to a** brief study of the human body from the electric and magnetic point of view.

human polarity

A study by the engineer Voillaume, entitled " Essay on the **radiation of man and** living beings", the publication of which in **1933** can be considered as a historical event, **reported** the emission by the human body of a certain number of hitherto unknown radiations, the characteristics of which he gave.



• * ' s 'NOTE
?

relating to the design of the front, representing with its subdivisions an element of Doctor Peyré's checkerboard

The earth is traveled in the magnetic North - South direction by MS magnetic rays (with magnetic support) and , in the East - East direction , West by electric rays ES (with electrical support) , accompanied by 1 ■ rays “ y (harmful rays)

- The **resulting** checkerboard is not formed of squares, but of rectangles whose dimensions are not strictly constant, but may vary slightly depending on the premises. •

■ In **the** colored drawing below, •which is the statement of an element of,; do r n i e r . measure. on the. ground , the main magnetic North - South bands are separated by 8 meters 60 from axis to axis , and the main East - West bands by 7 m e t e r s 66. The East-West bands are always a lot . wider than the m a g n e t i c North - South bands .

. The **rectangles** (in yellow) . formed by the crossing of the *bands* only emit r ray (RS , J , v . ^A) ; these are harmful areas . _ _

On either side of each yellow rectangle are two small rectangles (in green) . _ _ _
_ _ _ _ _ _ _ _ _ _ emitting much narrower ES+ rays :
positive ground , indicated by M. Orcel in his works as particularly favorable to vegetation . ■

The Peyré checkerboard is itself subdivided by secondary bands parallel to the main bands . _ _ There are 10 electrical secondary bands and 6 m a g n e t i c secondary bands that are wide enough and of a sufficiently intense radius to be _ _ _ _ _ _ _ _ r e p e r c e p t i b l e s . '

The element represented has been drawn and measured on p l a c e . But the dimensions of such elements, and, a f o r i o r i , those of their subdivisions . i m a y vary from moment to moment , there was no need to indicate measurements ' rigorous. The important thing for this drawing is to give • an exact idea of the p h e n o m e n . As an indication however , here are the widths of the bands _ _ _ measured in this experiment :

1 = 0	Mr.58	a - 0	Mr.28
2 = 0	M 38	b - 0	Mr. ¹⁴ ■
3 = 0	Mr 27	c . 0	Mr.0 4
4 = 0	M 08	d - 0	Mr.01
5 = 0	Mr 03		
6 = 0	Mr 01		

These are in particular:

1 ° **The Nerve wave** with double polarity, as well in the man as in the woman and which is emitted by the whole of the cerebral spinal system.

2 ° **The sympathetic Wave** also with double polarity emitted by all the nerves, forming part of the sympathetic system.

3 ° **The always positive Human Wave** which is characteristic of the human species. To this wave corresponds a racial wave characteristic of the species in both animals and plants: canine wave, porcine wave, feline wave, abalone or little bend wave, etc.

4 " **The vital wave** which is always positive and which is common to all living beings, animals and plants. The vital wave is not emitted by the sick parts of the living being.

5 " **The positive sexual wave** in the male, negative in the female and which is also common to all living beings, animals and plants.

6 ° **The always positive mental ray** , emitted by the brain and which is a material effect of thought.

7 ° **The personal** or individual Wave, positive in the male, negative in the female and which is special to each living being, man, animal or plant. This characteristic wave of each individual, varies from one being to another and makes it possible to recognize it.

Detailed information concerning human waves will be given in chapter V, on the radiation of living beings.

To speak of the polarity of a living limb or organ without at the same time specifying the emission one has in view is nonsense. The living being emits, indeed, simultaneously, several different waves and of variable polarity.

The nervous wave and the sympathetic wave are of double polarity. From a certain finger, it is a positive nerve wave that springs; of the neighboring finger, that is, au. contrary, a negative nerve wave. It is therefore only by specifying - although we place ourselves or point of view of the nervous wave, that we will be able to say that such a finger is positive, such another negative. In the same way, C.: is by placing ourselves from the point of view of the polarity of the **sympathetic wave** which it emits, that we will be able to say that such an organ is positive, and another negative. '

To claim that, while the man is positive, the woman is negative , is an evasive assertion that does not correspond to any tangible reality.

Electrical Polarity : If we examine **electrically** , we find that the human body is comparable to a dry battery for a flashlight. Shared by a median vertical plane, passing through the middle of the face, the body human behaves

a positive half and a negative half. The hands play the same role as the strips of copper, the end of which serves as the poles of the **pocket** battery.

If we seize with each of the two hands the two ends of a copper **wire** on which will have been inserted a micro-ammeter, we note the deviation of the needle **following** the production **and** the passage of an electric current. (experiment of Doctor Leprince).

The right side of the normal man is positive. the negative left side: it is the opposite in the normal woman. But there are individuals in both sexes whose polarity is abnormal and reversed (**approximately 5 %**).

In a household, the difference in electric polarity seems to be one of the conditions of the attraction and consequently of the happiness of the two spouses.

Direction of gyration of the Pendulum : A positive body is a body whose specific radii are specific positive EM waves; a body called negative emits, on the contrary, specific negative EM waves.

A pendulum held by an **electrically** positive hand rotates clockwise above a positive body on which it is tuned; it turns in the opposite direction above a negative body. If the pendulum is held by an electrically negative **hand**, its movements are reversed.

Human electric fields : When the poles of the human battery are not united, this battery forms an electricity capacitor whose contour of the electric charge is easy **to** delimit **using** an adjustment on ES rays.

This capacitor determines a double electric field, positive on one side, negative on the other, the limits of which can be determined very easily **0** using an adjustment on **y-rays** (**fj -r** on one side, **J** - the other).

Figures 23 and 23 bis show the main lines of these fields, the shape of which varies according to the individual and the extremities of which terminate **approximately** at the rear third of the upper part of the head and at the rear third of the feet.

The human body not being immobile, it goes without saying that the shapes of the fields change at the same time as those of the body and in particular according to the movement of the limbs.

Magnetic polarity humzinc : Considered from the **magnetic point of view**, the human body, like all objects of a substance

other than iron, forms a group of two magnets located in the extension of each other and whose line of separation is on a surface terminating at the level of the solar plexus.

The upper magnet has the shape of a bar magnet **0** rectangular section (and even square in young people). At the

old men, this section takes on a slightly more complicated form:




Fig. 23

Figs. 23a _

Limit of the human electric field seen from the front

seen from the side

Roglogo on rays 0 . The two halves of the field are rarely symmetrical. At the lower part the field ends in the posterior third of the foot. When the feet are not joined the field takes the form indicated above. .

upper magnet starts from the top of the head (negative pole) and ends or solar plexus (pit of the stomach or point of the sternum).

lower magnet has the shape of a sharp shear . It starts at the level of the solar plexus (negative pole) and ends at the heel of each foot (positive pole). (Fig. 24 and 24a) .

This lower magnet sits between the human capacitor and the spine. 11' is very easy, using a pendulum set on MS rays, to determine the contour, either on the person himself, or on an impregnation on paper of this person, or on a photograph.

The human magnetic charge, concentrated as indicated in figure 24 bis, is generally very weak and difficult to perceive with the instruments of current physics laboratories. However, a fairly small number of people possess it to a high enough degree that their approach is enough to **deflect** the needle of




Fig. 24 Vu de

profil

Seen from the front

Location of the poles and neutral lines of the human double magnet. As with electric fields, the points where the magnetic fields exit the body are cusp line points. Both on the head and on the feet, these cusps are behind the similar electrical ones.

This load seems to present itself in an identical way in man and in woman.

Double human magnetic field: To each of the two human magnetic charges corresponds a magnetic field whose contour can be determined very easily **using** a pendulum adjusted to radii (,).

Figures 25 and 25 bis give an idea of this contour both **in** an individual seen in profile **and** in an individual seen from the front.

Human electromagnetic field : The superposition of an electric field **and** a magnetic field **determines**, for the game. common to these two fields, a human electromagnetic field possessing physical properties analogous **to** those of the terrestrial **EM field**. ■




Fig. 25

Figs. 25a _

Profile

view Front view

Double magnetic field of the two human magnets
The neutral surface of separation of these .
two magnets passes through the solar plexus
(tip of the sternum).

Double human halo : The belief in the existence of a fluidic halo in the vicinity of the human head is very old. The artists who paint the portraits of holy personages often represent them provided above their heads with a luminous horizontal circle.

The dowsing clarifies this question. It allows us, in fact, to see that the possession of a halo is not the prerogative of saints alone, but that it is one of the attributes of every human being.

However, instead of being circular, horizontal, and hovering above the head, the aureole is generally oval, almost vertical, and placed behind the head. Moreover, this halo is not unique, because if the human being has a halo at the level of the head, he also has a second one, more flattened, above the feet.

Figure 26 reproduces, profile views, the two human aureoles; Foutes two are located in perpendicular planes or verti cal plane and inclined 20 to -25 ° on the vertical. Figure 26 bis represents ____ i ____... _ _ _ _ _ _1 _ r» ii _1_ 1_ _1 _ _1_ _1_ _1_ _1_ _t_____




Fig. 26

Plans des deux auréoles humaines (supérieure et inférieure). Elles sont dans des plans verticaux faisant angle de 20 à 25° avec la verticale.




Fig. 26 bis

Front views, the two human halos. The lower part of the upper aureole is shown in dotted lines because it is hidden by the head. The superior aureole exceeds the superior level of the head; its lower level rests on a horizontal plane passing through the lower third of the nose. The lower aureole seems to rest on the insteps.



Fig. 26 te

Imprégnation sur papier d'un aimant droit vertical et de ses deux anneaux (ou auréoles).

The shape **and** location of these fluidic rings obviously vary slightly between individuals.

rests on a horizontal plane passing through the lower 1/3 of the nose , and , through its upper part, protrudes by 0.05 and 0 , 06 the upper level of the head. Much more flattened than the previous one, in the horizontal direction, the lower ring generally rests on a horizontal plane. 0 0.05 or 0.06 above the top of the insteps.

The human double halo is a magnetic phenomenon: If we present a sheet of white paper behind the back of a person's neck, so **that** it goes beyond the top of that person's head, and if, **0** the help of a pencil forming a tip-antenna and a pendulum adjusted to MS rays, we study the impregnation thus produced in the laboratory, we observe:

1° That an oval ring about one centimeter wide has formed on the sheet of paper, which

emits MS + rays over its entire surface.

2° That, from the left to the right of the person, the ring is run by MS + rays which, if they circulated with the speed of light (which is probable). would do 8,500,000 revolutions per second.

The shape of the ring can be influenced by the action of a magnetic field (sewing machine for example).

If we operate in the same way for the lower halo, after having placed a sheet of paper resting on the insteps of a person, we observe:

1. That there is formed on the sheet of paper a flattened oval ring, from 5 to 6 millimeters in width, which emits on all its surface MS rays.

2° That from the right **to** the left of the person the ring is run by rays MS—probably circulating with a speed of 8,500,000 revolutions per second.

The formation of the human double halo corresponds to a general magnetic phenomenon.

Let's put a rectilinear magnet on a table and keep it vertical, so **that** its positive end rests on the table and its negative pole is in the air.

The position of this magnet will correspond to that of a standing human being. Behind this magnet, let us slide a vertical sheet of white paper, and **using** a pencil forming a point-antenna and a pendulum regulated on MS rays, let us study in the laboratory the impregnation produced.

We see around the negative pole a ring of MS+ rays flowing counter-clockwise and around the positive pole a ring of MS- rays flowing cos-clockwise.

It should **be** noted that the MS rays which circulate **inside** a ring are of an inverse polarity to that of the pole that this ring surrounds.

will say, by placing from the microphysical point of view, that **a**ll bodies, some **t**o a sensible degree, **o**thers **t**o an **i**nfinitesimal degree , are electrified, magnetic and radioactive. Just like a copper **w**ire , **a** cotton ball is a conductor of electricity : a silver spoon has a magnetic charge; an object of n **i**c **i** e l emits rays '**1.**, **p** and **y**. Seizing imperceptible energies, so far, our sixth sense will allow us to **a**dvance in the **k**nowledge of the physical properties of matter . and the practical applications that will be **m**ade of new discoveries, will bring to humanity immense **a**dvantages of **a**ll kinds .

A - Sources of emission from

BODY RADIATIONS

whose displacement correlates -
with the propagation of an
electromagnetic wave, called the
“ associated wave ”.

dowsing sense -

I" Corpuscles ' PH OTO N S

A support of light rays

• 2° ELECTRONS corpuscles ,
which constitute either the
electric current, or the rays 3
of the radioactive bodies,

A sup p o r t either ES—, ' or radioactive.

3" RAYS Corpuscles '2 .
radioactive bodies,
atoms of Helium.

Supported by specific rays of
this gas .

• (Pages 8, 9, 61)

Specific radiation_
of ELECTRICITY
exporting vertically!
specific rays- of bodies.

ES escaping from
electrical charges ; _ _

• J, Z, z, 7 escaping
electric fields.

(Pages 9, 57, 131)

II
Specific radiation
of MAGNETISM
exporting between 45° and
135"


the specific rays
of the bodies .
MS escaping loads
magnetic;

f>). f> escaping
magnetic fields. •

' (Pages II, 132)


B. Sources of
emission

ELECTROMAGNETIC WAVES. Only excite the radiesthetic sense




Specific vertical
electromagnetic
rays .
Has ES support.
(Pages 3, 57, 61 |


111
Radiation from BODIES, SUBSTANCES or OBJECTS
Humans, Animals, Plants, Minerals (Example of a copper urn)




Specific electromagnetic
beams angled at 45" and
135".
A s up po rt MS. (Pages
7, S. 58. 61, 103 I



Ray ons _
specific horizontal tro -
magnetic elements . _ _
ni support , (Pages S. 59,
6'2, 127)




Specific
electromagnetic rays
of the 'Fundamental Plane.
Has MS support.
(Pages 8, 110. 133|



ectro -magnetic rays
specific
to the M agne ti c Cross .
A s u p p o r t MS.
(Pages 8, 126, 133)

IV
TELLURIC vertical radiations from certain
regions of the ground



Harmful zone affecting part of a dwelling, originating from
radioactive minerals, or closed underground , or impregnated
streams . or Rectangles Peyré, etc... ' .
■(Pages 19, 59, 153)


V
Radiation of
LIGHT

the wave of sl u m ine u
ses (light and colors) .

2 " Caloric
waves ; (heat) . _
)" Chemical
rays . (photograph).

(Page 17)

VI
Radiation of
GEOMETRIC SHAPES




Sauvasti ka Sv a stik a
(Pages 60, 1501

VII
Radiation of
RADIOACTIVITY

. 1“ E m iss io n de r ad iat ion s
cular " 1. et fi '

2" Emission of certain -
electromagnetic waves called
y. bringing together the waves
v. fj . . T.

3° Emission of RS waves, specific
radiation proper to
radioactivity, also attached to
rays.



(p °9 is 15, 59, 61, 145,
147.)

HIV
Radiation of
LIVING

BEINGS (Humans, Animals,
Plants)

1 " Nerve wave . _ _ _ _ _

2 " » friendly _ _ _ _ _

J" » h u m ain e.

4 " » vital . _ _ _ _ _


5" » s ex u she .

6" " mental.

7 " » p e r s o n a l . _

8 ")) of specie . _

We find on living beings the
specific radiation specific to
radioactivity (RS) . _ _ _ _ _



(Pages 24, 60, 1701 .

Hearts of RadiedhAie Béasse ot
Corrensoit.

CHAPTER II

THE INSTRUMENTS OF THE RADIESTHESIST

Documentation

While the *raison d'être* of the sense of hearing is the perception of sound waves, that of the sense of sight, the perception of light waves, proper to the 6th sense, as it has been called Professor Charles Richet, is the perception of **EM** (electro-magnetic) waves, polarized or not. But, except for primitive man and ultra-sensitive subjects, the action of these waves on the human nervous centers is so weak that it only becomes conscious after the intervention of an amplifier, whose effect can be compared to that of the acoustic trumpet in the deaf.

Out of **100** people, at least **80** react to the first contact with EM waves. Although atrophied, the 6th sense does not differ from the others.

The **amplifiers** most commonly used by dowsers are the wand and the pendulum. both more or less perfected. They can be **neutral** or **active**. depending on whether or not they have any disturbing action on the radiation that it is desired to capture by their action.

We are going to describe them very briefly, one and the other, confining ourselves, for the moment, to neutral amplifiers . • .

MÉTHODE PHYSIQUE

§ 1*^r — Neutral Wand of Seekers

The rod is a Y made of a material that is sufficiently elastic and permeable to electromagnetic waves. The first ones. dowzers used a flexible wooden fork, hazel or hazel, 40 to 50 centimeters in length. -




Fig. 27




Fig. 27 bis

First way to hold the wand

Some still preferably use two flat whale rods half to one centimeter wide, 25 to 30 centimeters long, bound together by a ligature 4 to 5 centimeters long. Some dowzers find the length of 30 centimeters insufficient and

substitute those of 40 to 50 centimeters which give them, they say, more sensitivity.

Everyone must find, through personal trials, the way of holding the baguette that gives them the best sensitivity. Here are two great ways to choose from:

First way of holding the wand: According to figure 27, the wand must be held by the tips of the fingers, the palms of the hands above, the tip of the wand forward and slightly raised; it should not be squeezed between the fingers, so as to remain, supple, always in unstable equilibrium. When it enters a radiant zone, towards which the operator must move, the rod raises or lowers, at first slowly, then more or less abruptly, depending on the case (fig. 27 bis).

Second way : According to figure 28, the stick must be held lightly, but with both hands, the palms and nails above, the tip forward, slightly raised, the forearms horizontal, in contact with the body, the elbows apart. This position, which is not recommended by all the dowzers, is however, in our opinion, the most advantageous to obtain the maximum of sensitivity. Take care again that the stick, point slightly raised at the height of the solar plexus of your hand , flexibly forms an angle of **approximately** 45" , more or less according to the needs of your personal case. the branches, more or less, or stay at 45", depending on whether you feel the position most favorable to the reaction of the instrument, as it passes over the object of your study: known underground stream, or large cast iron weights .from 5 to 10 kg., etc ...



It does not seem, at least for the majority of the operators, that the direction of the movements of the rod has any relationship whatsoever with the polarity (+. or —) of the radiant object.




Fig. 28




Fig. 28 bis

Second (recommended) way to hold the wand

During the first attempts, the beginner is often too absorbed in the search for the best position, and, passing over an underground stream which takes the place of an exercise, he completely forgets to remember the object of his search,

the water ; it is not surprising that the reaction of his wand is slow... But with a little practice, the beginner will adopt, without thinking **about** it, the

good position of his wand, and worrying only about the object studied (or sought) the wand will manifest itself.

As Mr. Mellin pointed out in his work " La Dowsing agricultural and domestic" after a brief use, each of the ends of the branches of the rod, as indeed each of the faces of the point are charged with electricity of opposite name (fig. 29). This finding can easily be silenced using the black spherical neutral pendulum PS (1), supplemented by a polarity indicator (pocket battery or magnet) (2). Each operator has the greatest interest, by engraving on the branches of his




Fig. 29

The verification of the polarity of the extremities of the rod is done using a pendulum adjusted to electricity, using the control process indicated on page 52 of " Dowsing Physical Method :t, and using preferably a magnet. The + or — signs will then be marked on the stick.

(1) On sale at Editions du "Progrès Scientifique", 37, rue Rossini, Nice.

(2) See -page 52 d e «Dowsing Physical Method • polarity control .

wand, the signs + or -, to take note of the results of the experiment he has made in this respect, and to use them by always placing the same branch in the hand of the same electric polarity. He will thus realize that his instrument is much more sensitive. The man, in 95/ of the cases, has the positive right hand, and the negative left hand; the woman, in the same proportion, has the right hand negative and the left hand positive.

To be neutral, the rod must be made of a non-radioactive substance

whose electrical or magnetic charge is infinitesimal. The radiance of colors, having fairly accentuated physical properties, it must be black and its wire ligatures of the same color. The most advantageous wand and that which has been adopted by the greatest number of dowzers, is the wand **with** flat whale stems.

It is through the tip of the wand and not through the operator's feet that EM waves enter the human receiver. Conducted by the nerve threads, they reach the nerve center which triggers the violent reflex, which the will cannot oppose, and which causes the instrument to tilt in one direction or another.

To realize this, it suffices to present the point of the rod above a radiant source of small dimensions, a copper object, an aluminum box, a diseased point of the human body, and to observe the reaction.

- Wearing rubber soles or silk socks, condemned by some authors during dowsing operations, is therefore completely irrelevant.

Impregnation of the rod : A He ring which has received radiation from an object or a substance for some time (metal mass, or mass of water, etc.) becomes impregnated with radiation from this object or this substance, and Jes shines in turn. This is a temporary impregnation. Thus impregnated, the rod has the disadvantage of upsetting or distorting the results of the operations.

next steps, if they are not remedied. The operator puts an end to this temporary phenomenon by striking the tip of the stick against the exercise table, or against a wall, a tree, or the ground. One can still simply abandon one of the branches^ of the wand.

Impregnation of the hands : The same phenomenon by impregnation occurs for the hands which hold the wand. After each different operation, or from time to time during a prolonged operation, vigorously rub the palms of your two hands, then shake them separately as if you wished to get rid of the impregnation. This precaution is especially recommendable to the operator who insists on completely disimpregnating himself when he operates on microbial radiation, in medical dowsing.

§ 2 — The neutral pendulum

The current pendulum consists of a mass (sphere, cylinder, etc...) in wood, ivory, sandstone, metal. etc .., suspended from a thread .about 30 centimeters in length that can be wrapped around a stick of 0.04 to

0.06 centimeters in length, 1 to 3 millimeters in diameter approximately.




Fig. 30

**Pendule neutre
sphérique noir P. S.,
d'étude.**

As we explained for the rod, to be neutral the current pendulum must have each of its three elements (mass, wire and stick), composed of a non-radioactive substance , and of an infinitesimal electric and magnetic charge. and these three elements must be black in color.

The most advantageous shape of the mass is the spherical shape. We recommend our black spherical neutral pendulum , brand PS, built according to the data of the engineer Voillaume: with its suspensory thread in black silk, the black rod of which is used to conveniently adjust the length.

The neutral pendulum has the disadvantage of capturing both the multiple " images" of bodies and radiations, than these bodies and radiations themselves. Now, these images, in theory , are appearance in relation to reality. They are very often a cause of errors in the search for invisible objects. ■

On the other hand, we recommend the neutral pendulum PS to beginners who only work at first on visible objects, in order to perceive their waves. Indeed, the neutral pendulum PS constitutes a very

convenient study pendulum for the initiation to the first exercises, even if it means adopting as soon as possible the Revealer " Schumfell " anti-image, pendulum scientifically constructed and of first order, especially for subsequent professional work .

The pendulum should be held by the rod, squeezing as little as possible, between the thumb and the index (or, for some people, between the thumb and the middle finger), which the hand, like the fingers which suspend the pendulum, hang , relaxed without tension, motionless, the other fingers raised and spread apart without effort (fig. 34). Suitably adjusted in length and weight (we will see later what it is), and presented above a radiant body, the pendulum begins to turn, depending on the case, either clockwise a clock (direct or positive direction), or in the opposite direction (reverse or negative direction).

In the greatest number of cases, the pendulum has the advantage of being able to be handled at the first attempt by the most inexperienced operator, and of being more sensitive and more precise than the rod, the use of which requires some skill. experience and requires some flexibility.

Impregnation of the pendulum : A pendulum which has received some time the radiation of an object, becomes impregnated with the radiations of this object and radiates them in its turn. This is a fact of temporary self-regulation by impregnation. Thus impregnated with the radiations of an object, the pendulum has acquired the inconvenience of turning when, without changing the length of its suspension thread, it is presented to any other object of a different substance. This would be a cause of serious errors, for the following operations , if it were not remedied. The operator puts an end to this temporary phenomenon, after each prolonged operation, by giving a slight jolt to the pendulum by raising it suddenly , or better still, for greater safety, by knocking its base lightly against the exercise table or against the ground.

Impregnation of the operating hand : The same phenomenon by impregnation occurs for the hand which supports the pendulum. After each prolonged operation, it is appropriate to obviate this by rubbing the back of the free left hand with the free fingers of the right hand, which has the effect of bringing the two hands back to the same electrical potential .

The passive attitude in perception, according to the Physical Method : To obtain almost immediate, if not immediate, success, it

suffices in the majority of cases to adjust the length of the suspension wire, either of the neutral study pendulum PS, or of the Schumfell pendulum, at 126 millimeters, following the manner indicated in figure 46, page 95. This adjustment length corresponds to one of the millimetric adjustment lengths of silver. Thus, the beginner only has to present one like the other of these pendulums, at the indicated length of wire, on the vertical of a small mass of any silver coins. It is rare that the reaction of the pendulum does not occur imperceptibly and after a few tries for the most refractory. But, even in the absence of this precise adjustment indicated, the first following exercises will be amply sufficient for the beginner to quickly reach normal perception.

Under the same conditions as for the tip of the stick, it is through the oscillating weight and the suspension wire that the captured EM wave penetrates the human receiver and reaches the nervous center, which in turn produces the reflexes. of the hand which determine the movements of the pendulum — imperceptible movement of the hand forwards or backwards, in the event of swinging; double movement with orthogonal orientations (forwards or backwards, to the right or to the left), in the event of gyration.

Everything therefore takes place as if the assembly constituted by the body of the operator and his tool formed an apparatus receiving EM waves, the pendulum playing both the role of amplifier and adjustment accessory.

If, in general, a reflex action is unconscious, its effect is not always so. This effect which, in the dowser, translates into the movement printed on his detector, gives him awareness of a sensation of the sense of touch.

And it is the consciousness of this sensation which in the **civilized will** constitute the source perception, a perception **which** will then be well or badly interpreted by the intelligence.

In dowsing perception, we are free to reduce to a minimum, that is to say to the simple awareness of the reflex, this intervention of the soul, so as to make our receiving apparatus as passive and as unconscious as possible; it is the **passive attitude**. But we can also adopt various **active attitudes** which we will examine later.

Certain dowsers in whom psychic intervention is preponderant, have denied the possibility of perception in the passive attitude. But this opinion is contrary, not only to the laws of psychology, but to the facts themselves. Here are some proofs:

Proofs of the possibility of purely physical perception in the passive attitude: Whatever method he uses, any trained practitioner has noticed, during his prospecting work, that if by chance his pendulum was found on the vertical of an object, for which it has been voluntarily or not regulated, its pendulum suddenly began **to turn**; this completely **outside** of its intention. .

In another circumstance, it may happen that two operators have prospected the same substance and kept the precise adjustment of their respective pendulum on this substance. If one of them takes it into his head to renew the experiment on the substance, and the other holds his pendulum correctly, even though he is no longer aware of having it in his hand, his pendule — thus well held in the passive attitude, mentally and physically — also begins **to** turn. This physical phenomenon is based on equality of length and frequency of waves between the two pendulums considered, a sort of resonance or agreement, which most often is the basis of dowsing. It is by the same resonance process that TSF devices operate. The transmitting station only hooks the receiving station if the wavelength tuning button is in synchronism with the emitted Hertzian waves.

Let us point out a third proof. Let a pendulum set on a body turn. Let's even close our eyes. Unbeknownst to us, let a third party slide a thick plate of mica between this body and this pendulum, and the third party will see that the pendulum will stop immediately. This proves with sufficient evidence, the action of this screen on the waves emitted by the body and the impossibility of perception of the operator.

A fourth proof can be furnished by the phenomenon of "remanence" or "impregnation", obtained by the contact of any body whatsoever on a sheet of paper. Maintain this contact for about half an hour and then present the pendulum ^{above} the place previously occupied by the body: we will record, at this location, all the successive reactions of the pendulum (polarized gyrations, series, fundamental **rays**, etc...) although the body in question has not left any atom of itself **at** this place, but only a physical action of its specific sounds.

Many other proofs could still be given which would not bring more confirmation or physical fact of **the** dowsing. '

Passivity can be unconscious and involuntary; in this case, it is absolute, as in the two preceding facts. But it can also be imposed by the will; in this case, it is only relative and comes all the closer to absolute passivity as the subject has more control over himself.

The attitudes of the mental method : But there is no doubt that the soul can also intervene actively, notably in the order of desire, and that certain particularly gifted subjects arrive, by this studied and disciplined action, at considerable possibilities. It is the **expectant attitude** or that of "mental desire" or "mental orientation" recommended by certain authors and which, according to them, must precede research. It is **the selective attitude**, by which, in a beam of various radiations, we choose the one to which we want to be receptive, forbidding us the reception of the others. Finally, **the intuitive attitude** by which we ask, in a way, the subconscious to discover itself. Such is perhaps the attitude adopted by people who feel their pendulum lighten or weigh down, or who, above oil, for example, have the impression of '**throwing** it in a bath of oil. Such is also, it seems, the attitude taken by certain operators in order to assess **the** flow or the pressure of an underground watercourse.

If the passive attitude, absolute or relative, is the one that best suits the beginner or the operator who wants to prospect by the processes of the **physical** method, none of the other active attitudes suitable for "the **Method mental**", is forbidden to him, and he can resort to it successively, and at his pleasure, according to his particular faculties and the needs of

prospecting.

And by there, fall all the objections which one made **to** the method of physical dowsing. Nothing prevents the same person from having recourse to precision methods in **cases** where they are necessary, and, on the contrary, in completely different cases, from having recourse to other procedures (mental, psychic), in harmony with their special cerebral faculties and recognized as particularly effective in the cases in question.

But, as explained by P. Béasse in the appendix relating to Dowsing by the **Mental Method**, page 229, the beginner will have the greatest interest in remaining, during the first year, faithful to the Physical Method, for it is when he will be perfectly familiar with all the waves that one encounters in microphysics that the expectant, **seductive and intuitive attitudes** of the Mental Method will render him the most service.

§ 3 — Active instruments

Active **instruments** are pendulums with their **own emission**.

1 ° A pendulum with electric or magnetic charge determines on **a** near object effects of induction as well as, sometimes, effects of attraction or repulsion which are not without influence on **Jo** specific radiation of this object. In addition, it deviates more or less considerably from its fundamental angle.

2 ° A pendulum comprising a radioactive substance which amps up considerably the intensity of the radiation from a very close radiating source; in addition, by electric fields. and magnetic which it develops, it produces effects analogous **to** those of the preceding pendulums.

3 ° The pendulum terminated by a point is subject to the physical properties presented by this point, when it is directed towards the ground.

Among the **pendulums with magnetic charge**, we will cite the Arar B pendulum, by M. Baradat, which contains a small copper spiral; the Bovis pendulum, the aluminum spiral pendulum of the engineer Pitois, the electro-magnetic pendulum of Mr. Badoit, terminated by an ebonite rod acting as a point; the pendulum of M. Turenne. All these pendulums present this characteristic that they are not regulated and that they turn on any radiation, whatever the length of the suspending wire. With these instruments, the search for the identity of substances requires recourse either **to** mental **selection** or **to** the use of a **witness**.

Among the **electrically charged pendulums**, let us **quote** : A pendulum whose mass, made up of a rubber ball, is charged little **by** little with negative electricity.

A pendulum terminated by a point: 'this point was it a cone, **has the** property of extinguishing the EM and RA images located in the radius of action of this point.

A statically charged pendulum does not rotate above the EM images encountered **by** its field.

A pendulum traversed, at least partially, by an **electric current, even of** infinitesimal intensity, produced ' for example, by different suitable metals . This pendulum , its swinging produces electrical induction effects which are not without action on the radiation of the bodies it reaches. It **has**, on the other hand, the property of erasing **electromagnetic images**.

If with an **active pendulum**,. **not provided with a witness**, we take **using** the process of the tremors the **number of series** of a substance or a radiation, we find numbers very different from those which one obtains with the suitably adjusted neutral pendulum . These numbers differ, of course, from the particular number of the wave or substance.

Thus, while the "particular number" (or series objective figure) of money is **3** , we find the figure 7 with an Arar B pendulum, or with the Turenne pendulum ; 5 with the Pendulum Pitois. If we equip these pendulums with a "witness ". we find the number **3**.

The **colored pendulum** is also an active pendulum and its action is different according to its color and also according to whether this color is **with** electric or magnetic accompaniment.

All that we have said, in a general way, for the active pendulum, is equally true for the active rod. -

An electric or magnetic pendulum, however, deviates the fundamental angle of the substance or radiation above which it is presented: and when, after a series of jerks, its gyration stops, it oscillates in the direction deviated from the fundamental plane of the object.

The " Schumfell " Revealer

Patented S. G. D. G. — No. 734.234 (Registered model)


This electro-radio-magnetic device, of Alsatian origin , of very studied construction (1) is a perfected " anti-image " **pendulum**. Extinguishing images of all categories, **including** Afterglow and Imprint. its use relieves the operator Of the worry arising from this delicate question, while considerably simplifying prospecting.

Consequently , by using it, the acquisition of knowledge about the "images " from page 183 to page 206 is no longer essential, in principle, **unless** you seek them out in order to use them.

Right hand for right tier

correctly holding the " Schumfell Radio-Magnetic Revealer " , i.e .:

- 1° The hand arranged without stiffness in a swan's neck:
- 2° Fingers raised and spread apart:
- 3° The median plane of the ring (A, B, C, D of Fig. 32) and the bluish needle of the compass directed in the North-South line. '



Bring the median plane of the ring as well as the bluish needle of the compass into the magnetic North-South line:

- 4° Make the ring oscillate slightly , keeping it in this plane;
- 5° Let the thread slide between the fingers until maximum circular action.

(1) The "Schumfell" Precision Developer anti-image can be found at Editions du Progrès Scientifique , 37, rue Rossini . Nice.

Fig. 31 Suspension du « Schumfell »




Fig. 32 - Orientation convenable du plan médian de l'anneau

The Schumfell Revealers consists of a hollow metal ring, the upper part of which is surrounded by a winding of insulated copper wire, comparable to the device of a horseshoe electromagnet, the poles of which would be located at the level of the lower part of the cylindrical box which contains the compass used to orient it. •

This box, hermetically sealed, contains confined air which, for this reason, ionizes after a very short time, and becomes radioactive. The threaded rod which fixes the case of the compass, slightly exceeding the lower part of the ring, is extended by an ebonite rod and plays the role of a point.

To orient the device in the North magnetic direction

South (1), the device should be rotated so that the blue tip of the compass needle is positioned at 340° (Magnetic North), i.e. 20° to the left of the letter N (Geographic North) which is at 0° .

When the device is at rest, placed either horizontal or vertical, but in an orientation other than that of the North-South magnetic meridian, we can see with the help of the neutral pendulum, that it develops a magnetic field around it. The device is more than a radio-magnetic pendulum, which then turns on any radiation or body, and with any length of wire. It is thus suitable only for the method of mental dowsing.

If, on the contrary, placing the ring vertically, we make it swing in the plane of the magnetic meridian, North-South, we realize the conditions of Faraday's experiment, and we find that the winding is traversed by an induced electric current by terrestrial magnetism.

The apparatus is then transformed into an electro-radio-magnetic pendulum, and no longer turns except on the radiations for which it is suitably adjusted. It is then essential to the method of Physical Dowsing, object of this course. It is important to maintain it in the plane of the magnetic meridian (340° — 160°) while modifying the length of the wire until obtaining the perfect gyration of maximum intensity, which characterizes the appropriate adjustment.

The electric current which runs through the ring, the magnetic field which surrounds the electromagnet, the point, the radioactive radiation of the device, make it unsuitable for the perception of images of any kind and make it the pendulum anti-ideal picture. Let us add that the special composition of which a certain quantity of lead, which enters the alloy of the suspensory wire, constitutes a dowsing screen which protects the operator against the perception of radioactive images.

For a quick introduction to the use of the "Schumfell" electro-radiomagnetic anti-image developer, see page 121.

1 U 1. Magnetic North is that indicated by the magnetic needle. Geographic North is approximately 20° East of North magnetic, i.e. at 0° .

CHAPTER III

THE PERCEPTION OF WAVES

FIRST LESSON

EXPERIMENTAL RECLAIM OF THE PENDULUM ON THE TABLECLOTH OF VERTICAL RAYS DIFFERENT BODIES

(Sc refer to “ Physical Method Dowsing”, pages 27 and 28)

SUMMARY AND PRECISE DOCUMENTATION

Except in a few rare and accidental cases, any object emits over its entire surface a **layer of specific vertical rays**, transported towards the sky at a very great distance by electrically supported waves (ES—), and corresponding to each of the simple bodies of which it is composed.

To perceive these specific rays (or electromagnetic waves) using our sixth sense, it will be necessary to adjust our human receiver (that is to say the unit formed by our body and our pendulum) to the length of wave of radiation to be captured, as we do with our TSF receiver. Just as with this device, the more accurate the setting, the more perfect the hearing, the more the adjustment of the length of the suspension thread of our pendulum will be precise, the more our perception will be intense.

It was Viscount Henri of France who was the great popularizer of pendulum adjustment, the principle of which is due to Abbé Ferran. In that he rendered to the dowsing **an** invaluable service of which one will recognize, certainly, **one** day, all the extent. Many dowsers do not hesitate to say that their main secret of success in dowsing lies in a **carefully precise** adjustment. It is the regulation of the pendulum which will have allowed the drawing up of the Map of identity of the waves (see Table of identification, page 96) and consequently, to determine the incalculable consequences which will result from this fact.

The manifestations of the Pendulum : This manifests itself in four different ways: by gyration or rotation, . the ellipse, the oscillation, the stop.

Rotations

Ellipse oscillations or gyrations Ellipse Oscillations r

. 1t --^ --




Fig. 32a

The gyration or rotation takes on a meaning, depending on the polarity of the body studied or sought. •Thus a gyration is of •positive polarity when it is carried out in the direction of clockwise. A gyration is of negative polarity, in the opposite direction. A body is thus positive if, on the vertical of this one, the pendulum makes a positive gyration. This body is negative, if the gyration is in the opposite direction.

The ellipse is the transitory movement that the pendulum takes between oscillations and gyrations.

Oscillation is a beating, or rocking, back and forth in the same plane, of the pendulum. It occurs) in particular, on the ends of a bar magnet, or horseshoe (magnetism), to indicate the polarity, positive or negative, of each end or pole. It is the same with regard to a pocket battery (electricity) ; the direction of the oscillation on each end (or pole) constituted by the two flexible copper blades, is different, according to the polarity, positive or negative, perceived by the pendulum. It is also the same with regard to certain points in the study of the waves of form.

The stopping of the pendulum sometimes takes on its full meaning.

These various reactions must be received **normally** by the pendulum, that is to say following an objective and mechanical experimentation constant and uniform to all the dowzers. This will then result, quite naturally, in the unity of interpretation, which is essential for constituting a science.

One will understand, therefore, all the importance that **there** is, before experimenting on the unknown, of exhausting all the exercises that the observation of visible bodies can provide.

But before approaching the first stretching exercise, we wish to allow you to appreciate yourself, your dowsing possibilities:

. You can yourself know your provisions for dowsing : This is the process advised by RP Bourdoux: Hang your pendulum, with any length of wire, on the middle of your left forearm (from the wrist to the elbow), to find the direction of blood flow through your veins. If your pendulum does not take long to oscillate in the longitudinal direction of your

forearm is that you have sufficient dispositions to become a dowser. Even if the oscillation is weak, but exists, you must be satisfied. Some exercises, among those which will follow, will quickly increase your dowsing sensitivity. The faster the reaction, and the faster the oscillation, the better prepared you will be for the future. If, on the other hand, the reaction was slow to appear, do this experiment at other times, without worrying. The state of your health, or a bad physical disposition, can temporarily render you incapacitated. In the next few weeks you may have better luck.

Exercises:

1. To take their first steps, the beginner will get a black spherical neutral pendulum, of the “Scientific Progress” **type**.

He will be able to use this neutral spherical study pendulum during the first exercises on **visible objects**. These lessons can just as well be followed immediately with the electro-radio-magnetic pendulum “Schumfell”. Subsequently, the operator, no longer being a beginner, will preferably use, under the same conditions, the “Schumfell” **pendulum**, whose qualities are indisputable in the practical prospecting of **invisible bodies**, and in particular, in the constant struggle against **images which** makes it a valuable control instrument.

Momentarily equipped with the black spherical neutral pendulum PS mentioned above, which he will hold, depending on the case, either with his right hand or with his left hand, according to his manual conveniences, in the manner previously prescribed; he will present it on the vertical of any fairly voluminous object (bronze candlestick, aluminum box, nickel alarm clock, etc...).

Of course, without worrying about his own polarity, he will choose to hold the pendulum the hand with which the **handling** of this instrument will seem to him the easiest.

The mass of the pendulum being on the vertical of the object, about 5 centimeters away, he will cause it to oscillate slightly, then will gradually modify, more or less, and very slowly the length of the suspending thread, by rolling the rod between thumb and forefinger that will support it. This is the **experimental search**, by trial and error, for precise adjustment (fig. 35).

During this operation, he will follow the movement of the pendulum obliquely, without looking at it: his gaze will be fixed, on the contrary, on the object studied, while keeping his mind attentive as to the research, passive as to the result.

The hand suspending the pendulum will be arranged hanging, relaxed, "gooseneck", the two fingers squeezing the suspension rod as little as possible, in the manner of figure 34:

That all your thought is then focused on the object studied (or sought): above all do not think of anything else.

At some point, the rocking motion will begin to turn into a very elongated elliptical motion; then the minor axis of the ellipse will gradually enlarge and its shape will approach more and more that of a circle. Finally, the gyration will become absolutely circular and will take the maximum speed and will not be able to stop. **At that moment, the instrument will be correctly adjusted "experimentally"**.

At its beginnings, the neophyte, without worrying about the meaning of *gira*..




Fig. 34

**Manner of
holding and rolling the rod of the pendulum between the fingers.**




Fig. 35

Perception sur les rayons verticaux

tion of the pendulum, will have to multiply this first exercise a great number of times, so as to acquire a great flexibility of the fingers. It is on the acquisition of this flexibility that the success of the exercises that follow will largely depend .

2. Adjust, as it was said above, the pendulum on an aluminum object, for example, and when the adjustment will seem perfect (either in the neighborhood of 143 or **164** millimeters of length of wire), without disturbing this length of wire , present the instrument on the vertical of another aluminum object: start it by a slight rocking and note that it turns. If it doesn't turn, the first setting **was** wrong.

3. As soon as the first exercise has been successfully completed, repeat **it** , under the same conditions, on another **object in** copper, iron, nickel, silver, tin, lead, etc. Note that each time the If you change the substance, the setting must change (except in the exceptional case where almost identical setting lengths would suit two different substances, such as 133 millimeters for silver, 1-33.50 for copper).

4. These first notions being well acquired and possessed to the full, multiply the stretching exercises by passing very quickly from one radiant source to another, proceeding each time to a rapid adjustment, but as precise as possible of the pendulum, by the simple rotation of the stick between the thumb and the index finger.

To do this, present the pendulum successively, on a table. in wood, on a marble slab, on a glass object, on a bowl filled with water, on the back of the hand, etc. ..., etc.... while imposing itself to

obtain each time a fast and perfect adjustment. Passing from one substance to another, **do** not fail each time to de- **impregnate** the suspended pendulum by subjecting its base to a slight shock on **the** table. This precaution is essential to avoid any error.

5. Adjust the pendulum above (between the two blades: negative gyration) a pocket battery, upright, even used (adjustment on ES rays. Specific rays of electricity), then present the pendulum thus adjusted above of a wire leading to a lighted electric bulb and thus observe that the electric current passing through this wire emits ES rays.

6. Set the pendulum on one of the poles of a rectilinear or horseshoe magnet, placed **flat** , horizontally, that is to say on the MS rays, the specific rays of magnetism . According to the pole, there will be oscillation, positive or negative . Present this pendulum, thus adjusted, above any iron object. Note thus that this object is magnetized, since it emits MS rays which are manifested by a positive or negative oscillation or gyration.

7. Set the pendulum successively on cardboard colored with each of the seven colors of the spectrum. '

Then note that if we present the pendulum above a sheet of white paper, it turns, whatever the length of the suspension wire, which is not surprising because of the many nuances of which the assembly makes up the color white, there is always one for which the setting corresponds to the length of the pendulum wire.

*
Hz±

Read the 3Ci^{e now} lesson, on the pitfalls in ray thesis, page 217.

PERCEPTION OF SPECIFIC RAYS

(Refer to “Physical Method Dowsing”,
pages **51** to **53** and **65** to **76**)

SUMMARY AND PRECISE DOCUMENTATION

Let us specify, as of now, that the radiation of the matter by the electromagnetic waves and by the corpuscular radiations, is carried out in two ways which we will study in detail in this course: the first, by **direct radiation** (the rays specific, the radioactive rays, the rays of form); the second, by indirect radiation (images).

Considered from another angle, all substances or simple bodies (of the animal, mineral and vegetable kingdoms) and all radiations (electricity, magnetism, radioactivity, etc.) radiate, emitting rays or waves, whose origin is, presumably, atomic, and due to the movement of planetary electrons around the nucleus of each atom.

The engineer Voillaume, who discovered special electromagnetic waves and studied their common physical properties, called them “waves specific to simple bodies”. (Review the preliminary notions on this subject, page 4).

These specific waves are **electromagnetic**. They are not unique for the same body, but, on the contrary, multiple, although in limited number. They are also of different wavelength. But they have common physical properties, notably that of having the same polarity, the same Particular Number and the same Fundamental Angle.

Always identical, they characterize the simple body; and, by the recognition of their particular elements, make it possible to distinguish and identify it.

Left to their own devices, these waves are damped very quickly and are extinguished after a few millimeters of travel. It is therefore probable that they would have remained unknown if they had not been seized and carried away, at greater or lesser distance, by other more powerful waves, **which serve as their support**. These are the ES waves (with electrical support), the MS waves (with magnetic support) and sometimes RS (with radioactivity support) which escape from the ground. and the waves with support (•>, which escape from the extreme parts of all the bodies, and, in particular from the points.

Sources of emission of electromagnetic waves : A source of emission of an electromagnetic wave is a space containing energy capable of producing the wave in question.

Let us first recall that by **charge**, we must understand the quantity of electricity or magnetism contained in a body, and by **field**, the region subjected to the influence of a charge, either of a current (electricity), or of a magnet (magnetism).

Thereby,

a) Regarding radiation:

1° An electric charge, which can be positive or negative, is a source of wave emission with double polarity : ES-t- rays, or ES- rays (1). The ES rays (i.e. with an electrical support, rays emitted by the earth's electric field) have been called : **specific rays of**

2 1) The polarity, positive or negative. of an electro-magnetic wave, results from the relative direction of its magnetic vibrations with respect to its electric vibrations.

electricity. They are found on a pocket battery, or on the untwisted wires of an electric current.

Adjustment to this ES wave can be made either experimentally , that is to say by trial and error, by shortening or increasing the suspension wire of the pendulum, on the vertical and above the center of a pile of pocket, even used, held upright, or even on a light bulb, or also, automatically, on electricity by using the automatic millimeter settings in the Table of settings, pages 96 to 99.

vertical rays are electrically supported (ES). The **radius of union** has a double ES field.

2° A magnetic charge, which can be positive or negative, is a source of wave emission with double polarity : MS+ rays, or MS— rays. MS rays (that is to say with magnetic support , rays emitted by the earth's magnetic field) have been called: **specific rays of magnetism**. They are mainly encountered around each of the two poles of a magnet. .

Tuning to this MS wave can be done either mentally. by trial and error. towards the ends of a magnet. placed horizontally. be still. automatically on the Magnetism, according to the table of automatic millimeter settings , pages 96 to 99.

The emission can be vertical, or at 45° , or at 135° . or horizontal. The fundamental ray is magnetically supported. The union radius has an MS field.

3° An electro-magnetic charge, which can be positive or negative. is a source of emission of ES and MS waves (that is to say with electrical and magnetic support, rays emitted by the earth's electromagnetic field).

The setting can be done either on the ES wave. or on the MS wave, as indicated in the Automatic millimeter adjustment table , pages 96 to 99.

The 45° and 135° spokes are electromagnetically supported .

4° A negative magnetic field, is a source of emission of waves with support **w** and **o** — . **The w** ' rays are very penetrating waves, which escape from the extreme parts of all bodies, and, in particular, from the **points**, or, in a horizontal direction, from all the points of the line of apparent contour. horizontal of the bodies. The **o** rays , of a much weaker intensity, seem to accompany the **w** rays , at least in the emission of the points and in the horizontal radiation. '

To adjust experimentally on the **w rays** , adjust on the end of a fine point, fixed vertically. It is also possible to use, both for the **w rays** and for the **o rays**, the automatic millimetric adjustments indicated in the Table on pages 96 to 99.

The accompanying **to rays** , especially the **digital** and **visual nerve rays**, as well as the **mental ray**, are truly the active mechanical agent of remote perception . .

5 ° A positive magnetic field is a source of wave emission with support **w** and **o** + . issued under the same conditions. .

Same settings and same . considerations as the previous **4°** .

6° A negative electric field is a source of emission with support **U-** , $\frac{1}{e}L$ **Ç**. Automatic millimetric adjustments, page 98.

The perception of these waves is necessary in the search for or the suppression of **harmful telluric waves**.

7° A positive electric field is a source of wave emission with support **Ij** + , **E** and **a**. Automatic millimetric adjustments, page 98.

The perception of the **(J+)** **wave** is necessary in the search for and suppression of **harmful telluric waves**.

b) With regard to substances:

8° A radioactive body, is a source of emission of waves with RS support, **0** +, **v**, **Ç** The RS rays have been

called : **specific rays of radioactivity**. These RS rays, which are always positive, belong to the group of physicists' **y rays**. They must be an expression of cellular vibrational energy, as they are found on all living things.

Automatic millimetric adjustments, pages 98 and 99.

The perception of these various waves is necessary in the search for **harmful telluric waves**. .

9° A body, a substance or a material object, is a source of emission of electromagnetic waves **specific** to the simple bodies, of which it is composed.

The setting is made on “ control ”, experimentally; or, for some of the most common simple bodies, by referring to the Automatic Millimeter Adjustment Table.

The perception of these waves is necessary in the search for **hidden and lost objects**, **various deposits**, **sources**, infinitesimal traces of substances in analyses, fraud in food and industrial products, etc..., by vertical rays, or by the radii at 45° or 135°.

10° An object of determined shape is a particular source of electromagnetic wave emission, called a "shape wave", special for each geometric shape : sphere, cube, circle, spiral, etc. ..

The adjustment is made experimentally on the shape in question.

Some forms can be used for the formation of **harmful waves**, others for their elimination.

11° A living being, (man, animals, plants), is a source of emission of many electromagnetic waves . '

Particular adjustment on the waves: sympathetic, personal, human, vital, of species, etc..., indicated in the Special table of automatic millimeter adjustments, page 171.

Search for **diseases** and appropriate **remedies** . Search for criminals, **missing persons**, on photo, writing, witnesses. Life or death of a living being, etc.

Note, in passing, that the same space can emit one or more different emission sources.

We. let us insist on the advantages that the student will find on pages 96 to 99 in the Tables of adjustment and identification of simple bodies and living beings. These settings are identical for the two pendulums, either PS study or Schumfell Revealer, whose design has been provided for this purpose.

Sources of emission of corpuscular radiation: The **a.-** and **^** rays, corpuscular rays (atoms of Helium and Electrons), are only perceptible for the radiation therapist by the associated wave (expression of Louis de Broglie) , which accompanies them, specific rays of Helium for the first, ES waves for the second.

The setting of the pendulum will therefore be on Helium for the perception of the **a-rays** . , and on .the ES waves for the .rays p . Refer to automatic millimeter settings , pages 97 and 98.

Radio-active bodies emit corpuscular rays a . and p, among others. The **a** rays emitted

by a substance escape in the fundamental plane of this substance, and the p rays in a vertical plane perpendicular to the direction of this plane.

The radiation of a body : If we examine a body subjected to the action of the two terrestrial electric and magnetic fields (Fig. 33), we note that their effects give rise to the formation of special fields, exporting waves in various fixed directions.

For clarity, let's place the urn of figure 33 on a very horizontal table. This object is simple in shape. If from bottom to top we move our pendulum or a point-antenna in the vertical plane, then in the horizontal plane, passing through the plane at 45° , the reaction of the pendulum will signal to us the various rays emitted: vertical, at 45° , horizontals, etc. ..., with their various supports.




Fig. 33

Ensemble des rayonnements à inclinaisons fixes et lignes ou zones d'influence

Let us add to clarify figure 33, which it is important to fully understand, that :

The specific vertical rays of a body or an object have a negative electrical support when they are directed towards the sky; and positive, when they are directed towards the earth . An emission of vertical rays may possibly be of positive RS (radioactivity) support , when RS rays escape from the ground; as it can be an emission of harmful telluric rays fJ^- , v , ζ , when directed towards the sky, or $(J^+, e, 'j)$, when directed towards the earth.

The horizontal rays are supported ($\bullet>$), launched by the magnetic field determined by the body's own magnetic charge.

Rays at 45° and 135° are both electrically and magnetically supported.

Normal emission and its two anomalies : Note, from now on, that all bodies obey • the laws of normal emission. Their vertical rays, directed towards the sky, are under the **influence of the earth's electric field**, with negative electric support; those directed towards

the earth have a positive electrical support.

However, two simple bodies, **iron** and **zinc**, for reasons still unknown, show different anomalies with respect to this rule. **This** should be remembered in the choice of metals for exercises.

vertical rays of **iron**, therefore directed towards the sky, have a positive electrical support.

Zinc does **not** emit vertical rays towards the sky, but only vertical rays towards the earth, and these rays have a positive electrical support, like all other bodies.

However, some uninformed people will claim that their pendulum, if set and presented above a piece of zinc, starts to turn. But, if they observe carefully, they will recognize that it is under the influence of the " visual nerve ray" (gaze), which we will deal with later, and not under the influence of the earth's electric field that the gyration takes place. Let them stop looking at the object, and thinking about it, and the pendulum will stop.

In the same way, put the finger in contact with a piece of zinc, or direct our hand towards it as an antenna, in other words acting by our "digital nerve rays" (finger), and we will find that the pendulum regulated on the zinc will turn; but it will not be under the influence of the earth's electric field.

SECOND LESSON

BODY POLARITY BODY POSITIVE — BODY NEGATIVE

(Refer to “ Physical Method Dowsing”, pages 51,52, 53)

SUMMARY AND ACCURATE DOCUMENTATION *

Polarity of bodies : The phenomenon of electrolysis, by teaching us that the molecules of certain simple bodies are attracted by the positive electrode of the device, and those of certain others by the negative electrode, constitutes for us a first revelation of the existence because of the polarity (positive or negative) (1) of all bodies in Nature, whether they belong to the animal, vegetable or mineral kingdom.

It completes that given to us by our masters when we stammered the first notions of electricity and when they taught us that there were two kinds of electricity, one positive, the other negative, whose physical properties are different.

We are going to define here the polarity of simple bodies.

A simple body is said to be **positive** when the **specific rays** it emits are **positive**; it is said to be **negative**, on the contrary, when these same **rays** are negative. When a compound body is decomposed by electrolysis, the negative body is attracted by the anode or the positive electrode, and the positive body by the cathode or electrode. negative. The positive or negative bodies are therefore confounded with those which, in chemistry, are designated by the name of electro-positive or electro-negative.

(1) For technology, the polarity, positive or negative, of an electro-magnetic wave; results from the relative direction of its magnetic vibrations in relation to its electrical vibrations.

It is based on the phenomenon of electrolysis that the scientific dowsers had until now given to the positive bodies the denomination of **electro-positive**, and to the negative bodies, that of **electro-negative**. We believe that it is preferable to renounce a terminology which confuses beginners and makes them believe in the simple radiation of matter in an electrical phenomenon which does not exist.

Among the positive bodies are: Aluminum, Silver, Calcium, Carbon, Copper, Iron, Hydrogen, Mercury, Gold, Sodium, Zinc, etc...

Among the negative bodies we can cite: Arsenic, Bismuth, Chlorine, Tin, Nickel, Oxygen, Lead, Phosphorus, Sulphur, Tungsten, etc...

Polarity of the operating hand : But if, in the experiment of electrolysis, the positive bodies precipitate on the negative electrode, and the negative bodies on the positive electrode, on the other hand, the human receiving apparatus radiates the thesist reacts in a **different way** when the pendulum, regulated, is presented on the vertical of substances either positive or negative. Because in a dowsing research does not intervene only the polarity of the considered substance, also intervenes the **polarity of the operating hand of the operator**, that which suspends the pendulum.

Far from being a source of confusion, this question is easily resolved when one takes the trouble to study it summarily.

We learned earlier that electrically the human body is like a flashlight battery. One of the halves (divided by a medium vertical plane) is positive, and the other negative. The two hands play the same role as the two strips of copper which serve as the poles of the electric battery.

Usually the man is positive on the right side; negative on the left side. It follows that his right hand is positive; his negative left hand.'

Usually, the woman is, on the contrary, positive on the left side, therefore with positive left hand; and right side negative, with right hand negative.

But **there are** in one or the other sex individuals whose polarity is abnormally reversed (approximately 5%). It is also sometimes observed that some people change their direction of polarity one or more times during their life. It is thus essential to know perfectly the polarity of the hand usually holding the pendulum, and to check this polarity from time to time, to avoid errors of results, or a false • interpretation of the movements of the pendulum.

Let us recall in passing that the difference in polarity seems to be for Professor Bosset, of the University of Lausanne, and for many well-informed dowsers, one of the conditions for the happiness of a household.

This principle of human polarity being established, we will henceforth, for simplification, disregard sex and we will say that such a person has a positive **operating hand**, another a negative operating **hand**, without consideration of sex. By operating hand we mean the one that suspends the pendulum.

Consequently, a pendulum held by a **positive hand** will automatically rotate in the positive

direction above a positive substance or radiation . It will spin in the negative direction on a negative substance or radiation.

Similarly, for a pendulum held by a **negative hand**, the gyrations, in both cases, will take place in **the** opposite direction to that indicated above.

will **therefore** always be ^{the} polarity **of the** operating hand . regardless of sex, who will decide the direction of gyration on a substance, whether positive or negative.

Determination of the polarity of the hand which suspends the pendulum : It follows that to know the polarity of its operating hand, which is of the greatest importance for the understanding of this course and the exact interpretation of the movements of the pendulum , the beginner, after having regulated his pendulum on the copper, for example, positive body, will experiment the direction of gyration of his pendulum


on this body. If he spins positive, his hand will be positive; if it turns negative, his hand. will be negative. At various times, he will control this experience by renewing it. He will also check it on various bodies, at random, of which he will check the polarity obtained according to the Table of settings and identifications, pages 96 and 97.

He can still check the positive polarity of his operating hand in the following way: if the pendulum suspended by this hand rotates negative in the hollow of his free left hand. If, on the contrary, the pendulum turns positive in the hollow of his free left hand, it is because his right hand operator is negative.

Polarity of carrier wave : A pendulum tuned to the specific rays of a substance rotates in the direction given above, regardless of the polarity of the carrier wave which carries its specific rays; indeed, this support wave itself has its own polarity, either positive or negative, depending on the case.

If, our pendulum being set to the specific radius of a body, we place our left free index finger first above and then below that body, our pendulum will always turn in the same direction, despite the different polarity of the support.

If, instead of setting our pendulum to the specific radius
du corps, nous le réglons sur



ES +

Fig. 36

Réglage sur une onde A
support électrique d' u n
corps positif.

electricity (Rayons ES), it will no longer be the same. Held by a positive hand, it will spin counter-clockwise overhead. of the body, and in a direct, positive sense, below. (Fig. 36).

negative electrical support , while those directed towards the earth have a positive electrical support.

To conclude, a body

positive e'h a body whose specific radii are positive. A negative body, a body whose specific rays are negative. In the phenomenon of electrolysis, the electro-positive bodies are none other than the positive bodies, and the electro-negative bodies the negative bodies.

One will find on page 50 of "Dowsing Physical Method", the explanation of the phenomenon of the polarity of the electromagnetic waves on which we will not return. •

We give below, however, the very interesting means of checking the exactness of this polarity.

Control of the polarity of a body: Normally, a pendulum tuned to a positive wave must, on reception of this wave, turn clockwise; on the contrary, set to a negative wave, it must rotate in the opposite direction. It sometimes happens that in some people, the reflex is badly accentuated and that the pendulum turns indifferently in one direction or the other. There is room for improvement in calm and regular short-term training for one or two weeks. Sometimes the correction is facilitated by holding the pendulum adjusted, not between the thumb and the index

finger, but between the thumb and F major. In any case, in the most disadvantageous - circumstances, the essential thing at the beginning is that the pendulum should turn, oscillate or stop when appropriate; the polarity will become clear later with a little practice.

But sometimes it is necessary to determine exactly, with serious control, the polarity of a body or of a radiation. This is achieved by using a " polarity indicator ". It must be remembered first that only precise adjustments make it possible to obtain a normal gyration, that is to say in the exact direction of the polarity of the wave received. The pendulum being regulated on a positive body, for example (silver or copper), one touches with the middle finger of the left hand the positive pole of a pocket battery, even used (for the batteries of French manufacture, generally the blade the shortest), and the pendulum must continue to turn positive. If, on the contrary, we touch the negative pole of this battery, it begins to oscillate.

It would be the same if we put the middle finger of the left hand in successive contact with the poles of a magnet (the branch marked N or — being positive). ""

The reactions of the pendulum, after the intervention of the " polarity witness ", become normal again and are good indicators of the polarity of the wave studied.

The polarity of the most usual bodies : Among the simple positive bodies, we will mention: aluminum, silver, nitrogen, barium , boron , calcium, carbon, potassium, chromium, copper , iron, hydrogen, magnesium, platinum, radium, sodium, zinc, gold, mercury.

Among the compound bodies: water (negative polarity).

Among the complex bodies: air.

Among the simple negative bodies: antimony, arsenic, I bismuth, bromine, chlorine, cobalt, tin, fluorine, iodine, manganese, nickel, phosphorus, lead, I sulphur, 1 st te. l ure, the uran .um.

it is understood from now on that the methods described in the continuation of this course apply to a positive hand, that is to say: at approximately 95 % of the case of men. It will therefore be up to a male operator with a negative hand, or about 95 cy from queens, to reverse all the moves described later in this manual.

Polarity of electricity and magnetism: the reaction of the Pendulum is not different, as previously supposed , according as this reaction is provoked by the bodies, whether positive or negative, on the one hand ; or that this reaction is caused by the radiations of electricity or

magnetism on the other hand.


As a result, we will find that on the positive pole of an electric pocket battery (in principle the shorter blade), the pendulum performs a positive gyration (as it would on positive bodies). It will spin negative to the negative pole (long blade). Be wary of manufactures of foreign batteries including the:; polarities are sometimes different from those of French manufacture . Note that, on the center of a pile placed upright, the pendulum oscillates between the two poles.

The menie remark is to be made on a magnet (magnetism). auil either horseshoe or straight. The pendulum rotates positively or negatively depending on the poles. Note also that. on the center of the bar magnet, or in the hanger of the horseshoe magnet , the pendulum swings.

These observations are confirmed when, having directed our index finger with the free hand, towards a cloud, for example, we find that the pendulum turns positive, when the cloud is charged with positive electricity, and turns negative, when it is charged with electricity. negative electricity.

Finally, to avoid any hesitation for the beginner, we will specify that, when, in the exercises which follow, he has to regulate his pendulum on electricity (on a pocket battery), he will present it between the two poles of the stack upright: the maximum oscillation will indicate the perfect setting. For adjustment on, the magnetism (on a horseshoe magnet), the pendulum will be presented in the hanger; and in the case of a bar magnet, it will be presented on the center: in these last two cases, the maximum oscillation will indicate a perfect adjustment.

The setting to electricity or magnetism may



as well be done respectively on the poles (ends) of a pocket battery, or of a bar or horseshoe magnet. but then the pendulum will manifest itself by oscillations, which, according to the pole, will be positive or negative.

Exercises:

8. Set the pendulum on several of the bodies mentioned above and note that it does turn in the direction specified above.

9. If it is otherwise, use the polarity indicator (used pocket battery or magnet; see “**Physical Method Dowsing**”, page 50).

Repeat the exercise as long as necessary for the pendulum to turn automatically in the normal direction.

Impose, whatever the polarity of the observed body, to always take very precise experimental settings.

‘TROISIEME LEÇON

GYRATION OF THE PENDULUM BY THE INTERVENTION OF THE FREE HAND IN ANTENNA OR IN CONTACT WITH THE BODY

(Refer to “ Physical Method Dowsing”, tolls 54 and 55) A. —

SUMMARY AND PRECISE DOCUMENTATION

Index in antenna : If one directs the index of the free hand. (the one which does not suspend the pendulum) towards an object, as figure 37 indicates to you, the pendulum regulated beforehand on this

'TROISIEME LEÇON

objet, mais écarté de cet objet, se met à tourner.

Que se passe-t-il ?




Fig. 37

Rayon nerveux digital

The **nerve ray**, whose range is practically unlimited, comes out of the index finger and strikes the object. Then, in return, it conveys towards the operator, by serving him as a support, the ray ES or MS which serves as a vehicle for the specific rays of the object. It brings back, therefore, these last rays to the operator.

The human receiver is thus instantly activated . Experience proves, in fact, that an EM wave can perfectly serve as a support for another EM wave which propagates in the opposite direction by the same route of the free index finger.

The range of the nervous ray being unlimited, it then becomes possible to identify the nature of an object situated at a considerable distance.

Exercises :

1.O. The beginner will practice adjusting each time, **experimentally** , his pendulum, by the **nervous ray of his free index finger**, on numerous objects located first at . : small distance, then at greater and greater distance, finally at very great distance. However, **he** will only seek to identify the nature of a very distant object when he has fully grasped the substance of the present lesson.

11. He will first practice exercises more within his reach as a beginner, including this one: He will adjust his pendulum to the vertical of any body (a silver coin, for example, which is a positive body). Then, instead of continuing to present its pendulum on this vertical, it will present in its place, on this vertical, the left index in antenna. Note that the pendulum adjusted for this body and held with the right hand, far from this body, but at the height of the left index finger, rotates as if it were presented on the vertical of this body.

12. **12.** Identify, after adjustment on electricity, the polarity, positive or negative, with which a cloud is charged, while avoiding, however, to make this experiment in stormy weather.

13. Get used to adjusting the pendulum experimentally by taking the previous position, that is to say by looking for

singing the precise point of adjustment, by trial and error, the pendulum remaining away from the body studied, the index. left still on air.

Cover this silver coin with different substances: paper, leather, wood, cardboard, porcelain, sheet of zinc, copper, aluminum foil, silk, rubber, wool, various fabrics, etc... Note the permeability or not permeability of these various substances to vertical rays. Note that these rays being electrically supported, the electrical insulators do not allow them to pass.

14. The previous exercise having been completed, it will be possible to check whether it has been successful, remembering that the real electrical **insulators** are glass, porcelain, slate, dry paper, dry cardboard, natural silk, rubber, etc... As for the **bad conductors** of electricity, they are wood (unvarnished), carbon, earthenware, cotton, artificial silk, marble, etc...

In the previous operation, it is advisable to remain physically and mentally neutral. Never look at the studied object, nor look at the pendulum, in order to prevent that this one turns under the action of the specific rays exported by the **visual ray**, which is produced under the same conditions as the **ray of antenna**. This visual ray, which we will study later, is caused by the meeting of the rays of a body with the nervous rays which start from each of the two eyes of the operator.

15. the coin covered (by a third party who will put himself in a state of perfect neutrality) with the aforementioned electrical insulating screens, asking this third party to note down his successive findings. Indeed, the perception in this exercise must only be carried out with the finger in antenna.

B. — SUMMARY AND ACCURATE DOCUMENTATION


Index in contact: It goes without saying that if we bring the index in antenna closer and closer to the object, until it is almost **in contact** with it, the pendulum, regulated

on this object, will turn, but on one condition, however, and that is that it is held away from the object. (Fig. 38).

Let us put, in fact, the pendulum, regulated, on the vertical of the object, it will turn. Leaving it in this position, let us touch with the index finger of the free hand the upper part of the object, that is to say always on the vertical rays, . the instrument will stop.

The reason for this is very simple :

Captured by the pendulum (held on the one hand, for example, by **the** positive hand), arrive at the



, **Figs. 38** Nerve Centers of the Operator, • the Specific Support ES— Rays of the Object. Through the free hand, on the other hand, identical rays will arrive. These two wave trains propagating in opposite directions, but of the same nature, since both emanate from vertical rays, will meet in the body of the operator and will produce an interference

there. From then on, his nerve centers no longer being impressed, the reflex will be extinguished and the pendulum will stop.

On the other hand, suppose that the experiment is carried out on the **first** cylinder represented by figure 39. If, instead of. place the index at **a**, in the middle of the web of vertical specific rays (on which the pendulum is suspended), we place this index at **b**, point of emission of specific rays **at 45° to support MS**, the meeting of two waves propagating in the opposite direction, but of a different nature (vertical rays on the one hand, ray at **45°** to MS support on the other), will not produce any interference and the pendulum will continue to turn.

Exercises:

16. The beginner will also practice the experimental adjustment a great number of times, the index finger of the free hand being in contact with the upper part of the object studied, that is to say with the vertical rays, but taking care, contrari-




Fig. 39

C — apparent horizontal contour line of the body

Unlike the previous exercises, to hold your pendulum **outside** the vertical layer of this object. In this case, **the** pendulum will turn.

17. He will find that if, while maintaining contact, he places his pendulum on the vertical of the object, the instrument will stop.

18. It will turn if the index, instead of being placed on the vertical of the object, is put in contact with the zone of emission of the rays at **45°**, the pendulum remaining distant **from** the object.

QUATRIEME LEÇON

GYRATION OF THE PENDULUM BY THE INTERVENTION OF THE VISUAL RAY

(Refer to “ Dowsing Physical Method”, pages 55 and 56)

^SUMMARY AND PRECISE CUMENTATION

The ray which issues from each of our eyes is a **nervous ray** absolutely identical to that which escapes from our fingers. Previously called capital radius. we have chosen the denomination **visual ray**, to avoid any **misunderstanding**, as will be explained later.




Fig. 40 Visual ray

If we hold in the positive hand, for example, **a** pendulum, previously adjusted to an object, but distant from this object, and if we look attentively at this object, the pendulum will begin to rotate by a mechanism identical to that determined by the action of the digital nerve ray of the preceding lesson. The visual nervous ray will meet the experienced object and will bring back towards the brain of the operator the **ES**, **MS** or **RS** rays according to the cases which serve as a support for its specific rays, and the pendulum will turn. (Fig. **40**).

Let's cover an object with a transparent drinking glass, spill it , and present. above the glass, and without looking at it, ■ a pendulum previously adjusted S on this object. Glass being an electrical insulator, the specific vertical rays with support **ES** **ES**— will not pass and the *pendulum* will remain motionless. Re , g let's keep the object, the ner- ray

'l ----- -- want visual, breaking through the glass,
Fig. 41 the instrument will start rotating .

(Fig. **41**) .

Instead of placing the object in a transparent cage, place it in an opaque cardboard box. Since the vertical rays are stopped by the cardboard, which is an electrical insulator, the pendulum set on the object and placed above the box will not turn. •

Let us walk our gaze over the box and try to make it penetrate into the interior, invisible to us, of this box. The moment the nerve ray from our eye meets the object, the pendulum will turn. Our gaze therefore acts not by a phenomenon of visibility, but solely by the nervous ray which it directs.

.. The visual nerve ray can be used to .discover hidden objects and to identify the nature of ' very distant objects, because its range seems, practically, unlimited.

Proceed in the same way by replacing the opaque cardboard box with a porcelain container: Note that the pendulum, adjusted, suspended above the opaque porcelain container does not turn. the vertical rays being stopped by

porcelain, also an electrical insulator, and the visual ray. unable **to** cross the porcelain, therefore **to** join the covered object, will not be able **to** make the pendulum turn.

Exercises:

19. Set the pendulum on an object of a determined substance and note that it turns if we look at an object of the same substance far from us and from our pendulum.

20. Experiment on an object. of copper. nickel. silver, aluminum.

21. Practice **adjusting the** pendulum **using** the visual ray by looking at objects of different substance, verifying that the adjustment is correct by placing the pendulum thus adjusted on the vertical of these objects.

22. For the study of distant objects. you can use • a binocular or a telescope held in your free hand.

FIFTH LESSON

WAXING OF THE PENDULUM BY THE INTERVENTION OF THE MENTAL RAY

(Refer to “ Physical Method Dowsing ”, page 57)

SUMMARY AND PRECISE DOCUMENTATION

As we shall see further on in Chapter V, the **mental ray** is an electromagnetic wave quite different from the visual nerve ray.

When we think **of** a concrete object, our thought, by a mechanism which will always escape us, provokes a movement of the brain, and this movement triggers an electromagnetic wave which is none other than the **mental ray**.

As soon as it encounters the object, this ray, serving as a support for the specific rays which this object emits, brings them to the human nervous centers, and, if the operator holds **in** his hand a pendulum regulated on the substance of the object , this pendulum begins **to** turn, without it being above this object.

The mental ray, being a magnetic wave like the others, can be stopped by suitable screens; among these screens is porcelain.

Let us place a piece of silver on a table; let's set our pendulum on money and with our pendulum turn our backs **to** the table. If we think **of** this piece of silver, our pendulum, far from it, will begin **to** turn. While continuing **to** think of this room, let us turn our backs **to** his table and ask-

give an assistant to cover it with a porcelain plate, the gyration of the instrument will stop after two

or three seconds.

The length of the mental ray being unlimited, gives man immense possibilities for his research, and constitutes for him who knows how to use it, an incalculable advantage of which very few people suspect.

Mental tuning : As we have just explained, if we think of an object of copper, for example, our mental ray goes to meet this object and bring back to the human receiver specific rays of copper. From then on, it will become possible to regulate our pendulum on the copper and it will be **mental adjustment**.

Let us hold, in the positive hand, for example, a pendulum, away from any copper object, but let us **think** of copper, and while rolling between our fingers the stick of the instrument. lengthen or shorten the suspensory thread. There will come a time when the **pendulum** will begin to swing . Let us perfect the length of the wire until the gyration is done following a very regular circle and that its speed is maximum. At the moment. we will have carried out a **mental adjustment**, which we will be able to realize by presenting our pendulum thus regulated above a copper object and by noting its gyration.

Mental tuning, when done with all due care , is, generally. very precise.

Under the same name of **capital ray**, Abbé Mermet, who did not possess the means we have today to identify and distinguish between them the electromagnetic waves, confused the **visual ray** and the **mental ray** which, obviously, come out both **from** the observer's head.

To avoid this misunderstanding. we definitively renounce in this work the expression capital ray by which we have hitherto distinguished the **visual ray**. We therefore designate this ray by the denomination which suits it and which **can** not give rise to any misunderstanding.

Exercises:

23. Set the pendulum over a silver object. With the pendulum, turn your back on this object and think about it. Note, if the setting has been taken correctly, that the pendulum begins to turn each time you think of this object and, on the contrary, its gyration stops each time you stop thinking about it .

24. Repeat the experiment a large number of times with objects of very different substances.

25. Let us place, or rather have a third party (not hostile to dowsing) place it and without him telling us at the time, on a coin, a screen of glass, paper, cardboard or silk, etc... This screen will have no effect on the phenomenon: the pendulum set on the metal of the coin, properly set, will continue to turn.

26. But, let's place the coin. under a porcelain plate, however much we think about it, the pendulum will not turn. If we even prolong the experience, we will soon feel a headache determined by the prolonged reflection of the mind ray, which has failed to achieve its objective. Not to be confused with earthenware. which allows the mental ray to pass.

27. To practice mental tuning on numerous substances and to note, each time, that the tuning has been well taken, by then presenting the instrument on an identical substance and by noting its perfect gyration.

Selection of the waves : The preceding exercises have also taught us by mentally designating in our mind any body, among many others, to select the waves by precise mental adjustment of the pendulum on the specific waves of the body studied. We will recall however, that in physical method, the **selection** is done by the addition to the pendulum, of a “witness” (of which it will be question. page 118): thus the pendulum does not react any more but in the presence of • a body identical to this witness.

However, it is necessary to attach great importance to knowing, precisely what one is looking for, with or without rimentalization, is témoin. Si par exemple, ayant une main positive, nous avons pensé que la pièce d'argent sur laquelle nous réglons expé

an alloy of various metals where silver predominates, however, this piece will be presented under the pendulum as impure silver, and the pendulum will turn negative. It is therefore necessary to consider in this coin only the pure silver which it contains, by regulating on pure silver, and the pendulum will turn positive.

The same great care must be taken in the designation of the witness whose composition must be well known to avoid **any** error of interpretation of the exact movement of the pendulum.

SIXIÈME LEÇON
TIP -ANTENNA
REFLECTION — REFRACTION

(Refer to “ Physical Method Dowsing ”,
pages, 29, 30, 31 , 77 and 7 8)

SUMMARY AND PRECISE DOCUMENTATION

- **antenna work** : The proposed exercises consist in collecting on a screen the trace of the limits of the




Fig. 42

Delineation of ascending vertical rays. Vertical holding of the tip -antenna.

contour of a beam or a field, using the tip-antenna.

Case of ascending vertical rays, coming to strike the screen: when operating, as in the case of FIG. 42 on a beam of vertical rays, electrically supported, therefore coming from bottom to top, it is necessary to prohibit the use of the sheet of paper to record the limits of the **emission** zone of the object studied which is located in below. Paper being an electrical insulator, would not let itself be crossed. But use a sheet of zinc, or even operate by placing the object under the wood of the table (the latter acting as a screen), either under the marble of the table, under a wooden board or a metal plate placed on this Exercise Table. These different substances will serve as a screen to receive the trace of the contour of the beam or of the field, because these substances are good or bad conductors of electricity . tricity, but not insulating.

Case . descending vertical rays striking the screen: if the object is, on the contrary, above the exercise table, for example on a higher floor, we can perfectly trace, on the lower floor, its horizontal projection downward zontal on the upper surface of a sheet of white paper placed on the table. The paper, electrical insulator , will not allow the beam to pass through; it will be easy to collect the outline trace of this beam coming from above. This projection will only be somewhat displaced and, no doubt, also slightly distorted due to the refraction of the descending sheet through the floor.

Use of the tip-antenna : To trace with precision the limits of a zone of perception, let us extend our free left hand by a copper or steel rod, with a fine point (an old lady's hat pin, for example).), or even by a simple finely sharpened pencil, acting as a “ neutral tip-antenna ”. Always hold this tip very vertically.

To operate well at the antenna point, it is essential that the pendulum be held outside the surface on which this point moves, and as much as possible at its level.

The eye of the operator must always avoid looking at the pendulum, but he must watch its movements from the side, on the sly. The gaze must, on the contrary, fix the end of the tip-antenna which, held vertically, must be in contact with the surface of the screen. .

Each time the tip-antenna is inside the trace and in contact with the screen, the pendulum, adjusted to the object studied, will turn. When she comes out of it, it will oscillate.

The choice of a new sheet of white paper as a screen, and a well-pointed pencil as an antenna tip, moved very slowly, makes it possible to mark the line with a succession of small dots in the lead of the pencil. On paper.

' Exercises:

28. Determine the contour of any object placed under a sheet of zinc, then under the top of a wooden table, a marble table, a wooden board placed on the table, a metal plate placed on this table.

29. Use in the same conditions a sheet of dry paper (electric insulation) and note the impossibility of perception.

30. 30. Moisten this sheet of paper and note the perception.

31. Determine the outline of any object placed on the ceiling of the experiment room (metal or crystal suspension, for example), or placed on the floor above, on a sheet of paper placed on a work table . Note the deformation due to refraction.

32. Reflection : Place a copper coin in a small rectangular cardboard box, the lid of which will be • held (discarded of its front return), tilted at 30° , as shown in Figure 43 below:

Let us collect in front of the box, on a sheet of dry white paper forming a screen, the trace of the vertical tablecloth reflected




Fig. 43

**Réflexion d'une pièce de monnaie
sur un écran papier**

through the lid. This is generally not perfectly flat, but more or less convex or concave, we will obtain an elongated or shortened and deformed image of your coin.

33. To stop the rays at 45° , reflect on the lid of the box. •place an iron plate horizontally on the open box (Fig.- 43 bis).




Fig. 43a

Stop of this reflection by an iron plate

34. **Refraction** : Let us place on a stool placed below the table (of wood or marble) on which we are working, any object, copper for example. We will collect on the wooden or marble surface of this table, the trace of the vertical sheet which will affect the shape of the projection of the object on a horizontal plane, and that of the beam at 45° slightly refracted by the thickness of Table.

35. Above the object, place two or three cement tiles inclined at 30° or 40° . Note that the two images have moved and have been slightly distorted by refraction.

LESSON SEVEN

PRECISION SPHERICAL NEUTRAL PENDULUM ADJUSTMENT

(Refer to “Physical Method Dowsing”,
pages 27, 28 and 29)

SUMMARY AND PRECISE DOCUMENTATION

The adjustment of any pendulum, whatever it is (neutral or active) is based on this very curious property of the human receiver, .that the length of wire of an adjusted pendulum is proportional **to** the wavelength of the radiation **0** to collect , and this for all operators.

The human receiver is therefore comparable **to** a TSF receiver and its sensitivity in a given case depends, like the clarity of hearing, on the precision of the adjustment.

According to M. Voillaume, “by the constant use of a precise adjustment one develops one's own analytical sensitivity and one very quickly manages **to** no longer feel a wave if the pendulum is disturbed in an increasingly weak proportion. It is **on** this condition that we manage **to** discriminate the waves from each other and **to** identify them in a precise manner. »

As some people require proof, we describe below for their benefit the series of experiments by which **Mr. Voillaume losquelfos** arrived at **this** conclusion. •

Having adjusted his pendulum on pieces of cardboard of the following colors: blue, green, **yellow** , red, he found that the lengths of the suspending **wire** were almost proportional to the wavelengths of these colors.

Il construisit alors un pendule très précis (bâtonnet très mince, terminé par une pointe très fine). Sur une ligne droite, en partant d'une origine, il porta, après un nouveau réglage, les longueurs de




Fig. 44 longueur d'onde du
pendule en centièmes de microns
sur les couleurs

pendule (Fig. 44). obtenues en O.B. pour le bleu, O.V. pour le vert, O.J. pour le jaune, O.R. pour le ..rouge. Fixant à ces divers points la longueur d'onde moyenne de ces couleurs donnée par l'Annuaire du Bureau des Longitudes, il gradua les intervalles en centièmes de m. (microns) • '

. Sachant que la longueur d'onde moyenne du violet est **0 u 41**, il mesura sur l'échelle une longueur de pendule correspondant à la graduation **0,41** et le pendule tourna immédiatement en donnant la série du violet (voir § 6, Séries). Même succès pour l'indigo **0,44** et l'orangé **0,60**.

Pour serrer la question de plus près, M. Voillaume voulut opérer non plus sur les couleurs, qui constituent un élément insuffisamment précis, mais sur les

lines of the spectrum furnished by simple bodies in the gaseous state. He sought the wavelength of these radiations in the spectroscopic tables which give their value, and

brought the figures to a sketch, beginning with the D line of sodium (**0 J. 589**).

He noted that by presenting his pendulum stopped **at** this length ¹ above a salt of sodium, this pendulum rotated quickly giving the series of sodium.

Having repeated his experiment a third time and under even more precise conditions, he found a new confirmation of all his previous conclusions and was able to formulate this law:

There is a simple relationship between the lengths of pendulum wire corresponding to very precise adjustments to a body or radiation and the wavelengths of this body or this radiation as they appear on the scale of a spectroscope. This relationship, which is of the first degree, is that of proportionality.

This discovery prompted M. Voillaume to suppose that, in the art of the dowser, one was in the presence of phenomena having clearly as their basis the undulatory movements of the radiations of the bodies **and** that it was appropriate, consequently, to continue the verification of this hypothesis, which would probably lead to the observation of

phenomena of interference and polarization, likely to explain .biefJ,-- of s P.Oints mysterious; .

. To arrive at the énpncé of these conclusions, M.:. Voillaull'O,\$1vqit dN to construct a drawing of which it is reproduced here. . 0 the scale ' '1/2, a fragment' qtJi en' gives urte ^A ideè {Fig. 45). It is this blueprint that *allowed him* to draw up the table of 'adjustment' by 'wavelengths' in question. on page 96. •. — . 'u! • ■ - -•'

Continuing the series of his experiments, Mr. Voillaume, demonstrated experimentally: “•

1° That in the same way as Hertzian, caloric, light, radio-active waves, etc., the **specific radiation** of a body is really an electromagnetic wave and possesses all its properties; . you

2. That this radiation is not simple, but composed of a **beam of radiations** which it has been able to disperse and whose wavelengths correspond to each of the optically determined spectral lines of the body.

These invisible radiations, the existence of which is revealed by the reaction of the human receiver, are the same as those which act on physical apparatus. The radiation of sodium which, on the spectroscope, gives the well-known line D = 5.890, is no other than that which makes a pendulum turn on the sodium. There is therefore equality of wavelength between that of the yellow color of sodium and that of the metalloid corresponding to the line. '

3° That the pendulum, held by a positive hand,. rotating positively on all positive bodies. and negatively on negative bodies, the direction of gyration constituted a specific, very real property of each body, which he called **the polarity of the body**.

4. That specific rays are reflected and refracted like light waves. Bodies therefore have indices of refraction for these waves, like bodies transparent to light.

5. That, like light rays, they give rise to phenomena of interference and diffraction . .

To carry out these experiments, Mr. Voillaumo built a pendulum whose suspension is ensured by a **very** solid wire and the oscillating mass by a sphere **31** millimeters in diameter and weighing 40 grams. The neutral spherical pendulum "Scientific Progress" is built on the same bases, as advised by M. Voillaume in his remarkable work : "Essay on the Radiation of Man and Living Beings" (I).

” precision spherical neutral pendulum includes, to facilitate adjustment, a bayonet

(1) Scientific Library of Paul Lcchevalier, !2, rue de Tournon, Parts.

thin. finished at a peak. This tip also facilitates taking the wavelength on a simple double-deimeter. With the **neutral** pendulum c P. S.:» **the** length of the suspension wire , for the **1st** adjustment in millimeters, \$e counts **from** the axis of **the** rod, held horizontally, and gripped between the **two** fingers **suspenseurs**. (Voir Fig. 46).




Fig. 45

Wavelength of the pendulum in hundredths of a micron *without* the lines of the spectrum provided by simple bodies and the gaseous state

With this instrument one **can** obtain for **your** current prospection, the precision of half a millimeter, which **is** sufficient.

MÉTHODE PHYSIQUE

It is important to be convinced that *precise* adjustment **is** essential **to** operate **methodically** . **With** loose settings there are plenty of secondary phenomena that can confuse the beginner and confuse *you*. discourage (polarity reversals, interference, odd serial numbers). With precise settings on the contrary, **your** results are generally simple and clear.

Exercises:

36. **silver** coin, the **presence of each** of the three elements : **silver** , copper **and** **aluminium** .

37. Later, when you are familiar with the 8th and 9th Lessons, look for the " particular number" and the " fundamental angle " of these three substances, so as **to** identify them without possible error.

38. Note the presence of copper and aluminum in a piece of aluminum bronze, used in the manufacture of household utensils (the bronze of coins is an alloy of copper, tin and zinc) ; note the presence of carbon in the wood; calcium in marble; phosphorus in ivory.

39. Check **electrical settings**. (electric battery or incandescent lamp on, electrical conductor in circuit, etc.), and on **magnetism** (horseshoe or rectilinear magnet). Take into account, in passing, the peculiarities of polarity of electricity and magnetism, indicated on page 69. .

40. Check the settings specific to the **water**, **by operating on a glass** filled with water .

41. Check that settings on **oxygen** or **hydrogen** give gyration above water (H₂O).

42. Try the various settings on the **trough** (air, oxygen,

*)_..

a1.0te), .positioning .above . of a cellar/, of a, -floor-.C\1 of a known underground.

43 • Try the re **g fages** on the colors. A nuance of color **has** only one true setting, which corresponds to its own wavelength: "But you can set a **pendulum on** the different harmonics" of this wavelength, which are sometimes easier to take than the true setting, because they **are more** powerful^ 1

Millimetric adjustment of the precision neutral spherical pendulum, either Voillaume or " P. S.". The precise adjustment of these two identical pendulums; essential for the identification of radiations. It is necessary, in fact, - the "success of 3 or 4 lengths of known millimetric adjustments, of the " particular number" and of the "fundamental angle" of a radiation, to distinguish it, without possible error, from other radiations or waves. Knowledge of these three elements enabled Voillaume to draw up an identity card for many simple bodies and usual radiations. A table of these characteristics can be found **on** page **96** . .

• As soon as the 'beginner' will have acquired the rapid habit of experimental adjustment, that is to say by personal trial and error, which experienced dowers perform instantaneously , by playing themselves, he will be able to afford the use of these ' lazy, automatic, millimetric adjustments, established in this Table, as desired, in advance and definitively for the most common bodies. The convenience and rapidity that this Table provides, thanks to the use of a millimeter ruler (or a simple double decimeter) also allow a technique of adjustment of a rigorous punctuality. .

But, let us repeat, this Table, useful to initiates to save them time, is not recommended for beginners . Before using it, you must personally overcome the initial difficulty of the experimental setting.

Manner of automatically adjusting the neutral spherical pendulum PS and the Schumfell developer by millimeter adjustment. Figures **46** and **51** give, better than by explanation. how to take a measurement, millimetric, according to the Voillaume Adjustment Table, using a draftsman's double decimeter, in the absence of a special ruler.

The millimeter lengths in this Table have been established for an operator of average sensitivity. But to obtain the best gyration, there is an important advantage in taking into account the personal equation of each operator, checking (by a very precise experimental adjustment on several different bodies), once and for all, if it is advisable to increase them or reduce them by a fraction of a millimeter,"

or even by a millimeter in the. most disadvantageous case.




Fig. 46

Example of automatic wire length adjustment on 99 millimeters
(Sympathetic wave)
according to the wave identification table, page 96

**CHARACTERISTIC NUMBER SETTING TABLE FOR SIMPLE BODIES
AND THE MOST USUAL RADIATIONS**
allowing them to establish their identity card using the neutral spherical pendulum dS precision
PS,

0:0

or Voillaume, or Schumfell.
From table by C. Voillaume, ECP Engineer
(Between N and n must exist the relationship : $N=2n+1$)

Name of Corps or Radiations	Polarity and Serial Number (for positive hand)	Particular number N $N = 2n+1$	Successive millimetric settings of the most advantageous suspension wire	Fundamental Angle to Pendulum PS Add 90° for Schumfell _ _ _ _ _ s	Characteri stic - number n $N-1n$ ----- 6
. 1	2	3	4	.	.
Aluminum	Al + 3	3	196.90—164.40—101.25—62.60	128°	1
Antimony.	Sb -- 3	3	186.40— 154.85—117.25 — 84	30°40	1
Silver	Ag + 3	3	148.65—133—101.60 — 76.10	90° (East)	1
Arsenic...	Ace—3	3	192.20— 160— 119.30 — 70	229°O	1
Nitrogen.....	N+1	3	219.50— 161 — 131.20— 108.25 —	293°	■1
Barium ...	Ba + 7	7	212.80— 190.40— 152.85—116.40 — 95.20	326°20	3
Bismuth . . .	Bi —3	3	189.65—164—133—104.60	58°30	1
Bromine	Br —3	3	203.60— 156.20— 140.50—131.50— 110	313°10	1
-Boron	R+ 2	3	204.50- 164.45—153.50 — 98.60	345°20	■1
Cadmium ..	CD + 7	7	209.15 — 143.35 — 125.50 — 109 — 102.10	196°30 .	
Calcium...	Ca + 3	3	212.65 — 209.20 — 189.15 — 156.70 — 136.40	22°30	1
Carbon ...	C+3	3	218 — 154.50 — 129.50 — 97.75	336°45	1
Cerium ...	Ce - 7	7	179—154.70—141.75—137—121.60.	156°50	
Chrome...	Cr+3	3	1 67.80 — 1 4 5.25 — 1 3 2 .9 0 — 97.10	227°	1
Chlorine	Cl—3	3	147.30- 133.50- 125- 1 10- 108.60	290°S0	1

PHYSICAL METHOD

Cobalt....	Co—7	7	141.75— III .20—107.85 — 93.90	218°.	
Copper	Cu + 5	5	167.90- 163—133.50— 126.65— 100.60	41°32	2
Tin	Sn-2	3	61.25-94-140- 156.25- 168 — 210	21o	1
Iron	Fe + 2	3	95 — 122.50 — 138 — 192.50 — 203.50	180o(South)	1
Fluorine.....	F-3	3	71.50- 86.25- 115.50- 140.25- 155	286°	1
Helium	He - 1	3	116 - 121.50 - 173 (Adjustment of cor- puscu l ai res a)	289,,20	1
Hydrogen	H+1	3	66.25-89-112—134.50	Oh (North)	1
Iodine	I - 3	3	76 50- 109 50- 147- 163 50- 178	320050	1
Magnesium	Mg + 3	3	66.80 — 90.75-104-131	331°	1
Manganese.	Min—3	3	76.80 — 86.75-106- 110- 119—148—183	233°	5
Mercury...	Hg+II	He	81.50-118-148.50—167	315°40	
Nickel....	Ni — 3	3	192.60- 174.75- 149.30- 125.15- 112.60	2490	1
Gold -----	To + He	11	199-171.30- 160.30— 134.25— 108.50	270° (vaci)	1
Oxygen ..	O —1	3	236.40- 191.40 — 146.85 — 84.80	142035	1
Phosphorus.	P -.3	3	183—146.15—135.50-97.45	324040	1
Platinum.....	Dt + 2	3	171.75- 149.20- 138.60- 123.80	115°30	1
Lead	Pb-13	13	223 — 181.50- 157.35- 143 — 120.50	50°	6
Potassium.	K+ 7	7	239-188.85-169.15—142.19-110.75	l(o	3
...	Ra+5	5	169.85—160.50—110.40—102.35	161ois	2
Radon.....	Rd + 3	3	82.75-118.25-142.25—162.10-179.75		1
Silicon ...	If' + 5	5	204.80- 178.90- 123.75 — 96.40	37o50	2
Sodium	Na + 3	3	191.60- 174.50- 162.20- 129.55- 119.25	16°30	1
Sulfur	S-3	3	201.65 - 160.50 - 157.15 - 146.35	7803Q	J
Tellurium	Te — 3	3	209.10—179.90-159.75-147.55	25o	1
Thorium ..	Th + 3	3	210.30- 186.90- 159.20 — 110—95.30	214°J5	1
Thoron...	+ 1	3	72-119.25—145—195.50		•1
Tungsten.	W —3	3	220.70— 182.25 — 151.55— 134— 111.35	140o40	1:
Zinc.....	Zn +3	3	204.45- 187.90- 174.80- 115.20- 109.50	36005	1!

Name of Corps or Radiations	Polarity and Serial Number (for positive hand)	Particular number 1 st N N = 2n + 1	Successive millimeter adjustments of the suspension wire (1) . the most advantageous	Fundamental Angle to the Pendulum PS • Add 90° for Schumfell	Characteris tic number $N \frac{n}{2} =$
1	2	3	4	5	0
Zirconium. .	Zr+ 5		190.40- 141.60- 109.80- 105.50- 102.50	307°IO	
Air (Hollow) .	+ 2	2	99- 120- 161 - 178- 198	352°	
Water.	H2Q + 2	2	52-106-132-151-190-221	312°30	
Crude oil	- 3	3	63.25-94.75-117-148.60-154.75-192.40		
Electricity ... (ES waves)	EI ± 5	5	233.10-197.20-161.75-127	90° OR 270°	2
Magnetism. (MS waves)	Mag + 5	5	211.20-177.20-144.25-112	None	2
Radioactivity RS waves)	Route + 7	7	115.35- 131.85-149- 172	62°30	3
Rays e ...	Epsilon + 3	3	117.50- 142.10- 155.75- 171.80	198°50 •	1
Rays 'i ...	Nude -19	19	97.50-125.50-170-195	42°	9
Spokes 0...	Theta + 13	13	101 - 120.75-138- 180	219°3Q	6
Rays C ...	Dzeta - 1	3	112-130.50-160.50-197.	202°10	1
Rays w...	Omega+ 17	17	100-136.50-167.50-191.50	12°30	8
Rays o...	Omic. + 7	7	115-158.50- 172- 198	49°	3
Rays ;; ...		He	72.50- 126.40- 142-163		5
X -rays ...	Pi +II				
Rays p...	Kappa • Ro				
Rays -r ...	Tau				

Rayons G ...	Sigma + 5	5	105—148,50—168,75—203,50	297°10	2
Rayons VO .	— 3	13	104,80—127,50-138,75—166	339°	6
Rayons CL . .	— 3	3	103,75—128,75—145,25—176,50	39°	1
(Calorifiques) Rayons CH .	• + • 1	3	124,50—146—161,30	57°30	1
(Chimiques)					
Couleurs :					
Rouge	— 13		181,45		
Orangé ..●	— 12		213		
Jaune	' — 11		169		
Vert.....	— 10		132,42		
Bleu	— 9		103,36		
Indigo	— 8		86,60		
Violet	- 7		69,65		

NOTE. — The indication + before a serial number means that the ray has a polarity either 4- or —.

When a wave is dual polarity, column 5 of the table shows only the fundamental angle of the positive wave. To have that of the negative wave, it is necessary to add to the first 180°.

Thus: Rays **o** + = 49o Rays **o** — = 229o

Rays **0** + = 219o3Q Rays **0** - = 39o30

The rays RS, **B** +, v, £, are the **y -rays** of physicists.

Rays **and** and **£**, corpuscular rays (atoms of Helium and electrons) are perceptible for the dowser only by the associated wave (expression of Louis de Broglie) which accompanies them: specific rays of Helium for the former; **ES-** waves for the latter.

If we rule over a compound body, the series of this body must be the sum of the series of the simple component bodies, each of them taken only once, without consideration of the sign (Sel m ^{er} i ^{er} = N ^{er} 4Q — 3 + 3 — 6) — (Hydrated lime=CaO, HO=Ca+O + H=3+ 1 + 1 =5).

HUITIEME LEÇON

PENDULUM SELF-RECLOCK

(Refer.il c Dowsing Physical Method ”, pages 32 to 35)

SUMMARY AND PRECISE DOCUMENTATION

A pendulum. whatever it is, being a compound of various materials, necessarily emits the specific radiations of the substances which compose it. Thus, a wooden sphere pendulum emits carbon and hydrogen radiation. The Voillaume pendulum, like the “ PS ” pendulum. ball sandstone, emit radiation or silicon. If they are set to silicon, they will run continuously. They are said to be **self-adjusting**. If, thus self-adjusting and constantly rotating , we present one or the other above a body composed of silicon (glass or sand), it will stop rotating and begin to oscillate.

A neutral pendulum which has turned for a long time on a radiation ends up being impregnated with it, and becoming self-adjusting, turns permanently. This is why, to avoid errors, it is important to give to the suspended pendulum, a light shock, by its base, on the work table, each time one changes radiation. **The not very tenacious impregnation** vanishes immediately.

The colored pendulum : A solid, **colored pendulum** is self - adjusted for the color of the same shade as that of the paint with which it is coated. If in the search or the identification of a body we use, without realizing it, a setting very close to that of this nuance, we risk making a mistake. (Example: orange, setting — 181.45 and lead,

setting +181.50). This is one **of the** reasons why it is recommended to use only black coated pendulums, as neutral as possible. **Indeed , black is not** a color from the physical point of view; white, on the other hand, represents all the colors together. Thus the white pendulum would be self-adjusting for all colors.

The hollow pendulum : There is sometimes an advantage for certain researches on a substance to use a self-adjusting pendulum on this substance. Take a hollow container, for example a jar of pharmacist's ointment. Provide it with a suspensory thread attached to a stick. Place inside a fragment of the substance concerned. Modify the length of the wire experimentally until the instrument begins to rotate continuously. It will be self-adjusted for the substance introduced into the hollow of the pendulum if, presented above an identical substance, it begins to oscillate.

Autotuning is interesting to use, for example, by a particular ore prospector: the mass of his pendulum will then be made of this agglomerated ore. Many professional "dowsers" have built a pendulum whose mass is made of a small corked white glass bottle, previously filled with the precise water they are looking for: drinking water, mineral water at such and such a degree, arsenical water, hydrofluoric water, etc...

Exercises:

44. Check the self-adjustment on the silicon of a pendulum with a mass of sandstone (the neutral spherical pendulum "PS") or of glass; that of a wooden sphere pendulum for carbon and hydrogen ; of a rubber ball pendulum for the . chlorine, etc ..

Note that this pendulum oscillates above the bodies of the substance for which it is self-adjusted.

45. Find that the rubber ball pendulum is self-adjusting for negative electricity.

you
" , " " . < • t . • • • < • •

46. Look for bodies whose settings **u** are very close to . to. ce|ui. to , the .puance considered; hey. note **c.ezquj** .

se. Pa.sse. ;

*
' *
' *
' *
' *

. 47.. Constater. qu'un pendule. qui. tourne. longtemps sur:'

a .radiation **ends** up _by, self-adjusting. on. this radiation and to. constantly spinning. . . .

48. When the phenomenon has occurred, remove the impregnation from the pendulum by gently knocking it on the exercise table, at the base of the suspended pendulum •between the fingers. -

49. Do the . search for the hollow with an empty bottle. All setting on air,. oxygen. or .nitrogen succeeds on the hollow cavity,. fault, underground gallery, pipe ?empty water. But • the confined air is radioactive, it is better to try first on the trough. **the** various - settings on .the radio-. activity, according to the settings indicated in the Measurements Table. millimeters, page 96. This will allow a first verification. Then, we will use, in counter-verification, the self-adjusting hollow pendulum:. small bottle or small stoneware pot, empty, filled with air, plugged for the cavities supposedly closed, or not for the cavities having a relationship with the outside air. Don't be surprised by an old dip. who. later was filled. '

The substance which constitutes the hollow pendulum is irrelevant . It constitutes a simple support of the body for which. it is self-adjusting.

NEUVIEME LEÇON

THE SPOKES, AT 45 ° AND AT 135°

(Refer to “ Physical Method Dowsing” , pages 68 to 70) . .

SUMMARY AND PRECISE DOCUMENTATION

»

Under the action of the terrestrial **electric charge** , any body emits two thick sheets of specific vertical rays (**some** with support ES-directed towards the sky, others with support ES+ directed towards the earth).

Under the action of the terrestrial **magnetic charge** , this arbitrary body emits two cones provided with inclined rays, the. some at 45°, the others at • 135° (with MS+ support).

The .45° rays are very useful for raising existence. of a buried body and appreciate its depth. .

•

See figures 8 **and** 16, pages 7, 14, 62 and 76.

Exercises:

50. Check **the emission at 45°** on bodies of simple shape : cylinder, sphere, vertical or inclined prisms, cube, parallelepiped, body of any shape, by looking for their **apparent horizontal contour line**. (See Fig. 39, page 76.

The setting of the pendulum can, for the aforementioned purpose, be carried out on the one hand on the substance under study itself, by a prior setting on the vertical rays. of this substance. Thus adjusted, place either the pendulum, or better still the index finger of the free left hand, on the path of the rays at 45°, the starting point of which will be found on the line of the contour appearing.

horizontal rent of the object. The eyes will be fixed on this path and in the second case, the pendulum will be separated **0** to the right of the object and monitored from time to time **to** know the direction of its movement. The meeting of the pendulum (or of the left index, according to the accepted process Fig. 39) with the • radius **at** 45° or **at** 135° will cause the pendular reaction.

51. For more precise search of the beam path; at 45° and **0** 135° the index finger of the free left hand can be replaced by a very fine point, held between the fingers: point formed preferably with the substance of a body that is a poor conductor, but without an electrical insulator. Use for example an unvarnished wooden needle, tapered **at** its end, or an unvarnished pencil, well pointed: a steel knitting needle , etc... The points of origin of the various emissions will thus be recognized with more than • precision.

52. Verify by the test (settings and series) of electricity and magnetism that these rays at 45° are **magnetically supported** and that, consequently, they will pass through bodies that insulate electricity: glass, paper, cardboard, silk , wool, etc. .., impermeable to vertical rays, which are only electric.

53. Check that only an iron blade (kitchen knife) is capable of stopping the rays **at** 45° and **135** °.

54. Bury an object in the ground, of a noted depth, and recognize the emergence of the rays **at** 45° .

DIXIEME LEÇON

THE SERIES PHENOMENON THE PARTICULAR NUMBER OF A WAVE THE CHARACTERISTIC NUMBER

(Refer to “ Physical Method Dowsing”, pages 57 to 64)

SUMMARY AND PRECISE DOCUMENTATION

The phenomenon of series was discovered by the Abbé Mermet, who from the start attributed to it an objective origin . Since then, other dowsers have denied this objectivity , but the presentation which will follow demonstrates that they are in error.

Process for determining the serial number by shaking:

This process is ideal for the study of a hidden and distant body to be identified, but whose direction has already been found, or else which is found at a great depth under the feet of the operator. Currently we will continue our exercises on visible and known bodies.

The pendulum being regulated with precision for a determined simple body, if one presents it above this body, it puts itself. **to** turn (in the positive direction, if this body is positive, in the negative direction, if it is negative).

If we give a jerk to the pendulum, or if we simply raise it vertically, it stops its gyration for a temporary swing towards the operator's chest.

(effect of the shaking), then starts to turn again. At the end of a certain number of interruptions of gyrations, the number of which is constant for each body, it ceases to turn and begins to oscillate in a fixed direction which is that of the fundamental radius particular to the body. very near and a little above this body.

The **number** of interruptions or **jerk**s which determines the final arrest of the gyration is called the **series number of the body**. The series is positive if the pendulum rotates in the positive direction, negative in the opposite direction.

Process for determining the serial number “by finger”:

Before passing from the preceding operation to the following one, it is advisable, to avoid errors resulting from the possible impregnation of the pendulum, to de-impregnate it, by knocking it lightly, by its base, on the table of experimentation , and to disimpregnate oneself by vigorously rubbing the two palms . Adopt this habit before and after each operation. .

The so-called “finger ” **process** is only suitable for exercises on visiole bodies, because it requires the possibility of putting the index finger of the left hand in successive contact with this body.

First check the rigorous adjustment of the pendulum on a silver object, taken as an example.

Place the index finger of the left hand, without contact, above the silver object. Hang the pendulum over this object. : it will rotate positively, silver being a positive body.

Then lower the index of the left hand by putting it in contact with the money: the pendulum will oscillate.

Raise the finger, then lower it so as to obtain an alternation of gyrations and oscillations. Note that when the number of gyrations has reached the serial number, in the circumstance 3 for the silver, the pendulum will refuse to turn, will stop and will oscillate in the fundamental plane in particular of the body, very close and a little above.

Process for determining the number of series “colored squares of paper”:

Let's always use a silver coin as an example. Let us rigorously regulate our pendulum on this metal: it will turn.

Very close to this money. place a square of paper or cardboard one centimeter on a side. red: the pendulum will continue to turn. Let's add. by superimposing them on the first square, a second, then a third square of red paper.

Exceeding the silver serial number by one. which is 3. let's add a fourth square. Note then that the vertical radiation - of the money has stopped and that the pendulum oscillates.

Let us add a fifth and a sixth square, the pendulum will continue its oscillation.

But let us add another seventh, that is to say one more than the double of the serial number of silver, the radiance of silver will be reborn and the pendulum will begin to turn again.

The series number of compound bodies : It is advisable to verify, by their own series, the nature of the various elements constituting an object. A • silver coin, taken into study, contains, we know, money . copper and aluminum. It will be necessary for each element constituting this piece to adjust the pendulum beforehand, and successively, on each of these elements. (See example on page 99: setting on a compound body).

Experimental determination of the serial number : The - experimental determination of the serial number requires absolutely perfect adjustment. A slight misadjustment, still giving on the body, elliptical gyrations, leads to obtaining either multiples of the serial number, or numbers which seriously deviate from it.

One will find in the second column of the Table on page 96 the number of series obtained by these methods for the simple bodies and usual radiations.

It will be noted for simple **bodies** that each time it is equal to or greater than 3, it coincides exactly with the corresponding objective figure which we have called : **particular number** of radiation.

He can't, and, for good reason, to exist with a particular number less than 3.

number . Studying, a few years **ago** , the families of images produced under various influences by a simple body, **J. Correnson** noticed that these families include a number of individuals equal to the experimental number of series, each time this number is equal or superior there 3.

He drew the following conclusions :

1° That the number of series and the number of individuals of the families of images were determined by an identical cause still unknown;

2° That each time the serial number was less than 3, it was purely subjective, **which does not prevent it** from being used as such in research and experiments carried out with precision adjustment.

Determination of the particular number of a radiation : **J. Correnson** observed that if next to a source of emission of radiation whose serial number is equal to or greater than 3, a silver coin, for example , superimpose cardboard squares of a magnetic color (white, red, orange, yellow or green). in a number greater than one unit, at the series chmre of this radiation, the regulated pendulum revolving around this source begins to oscillate.

. Instead of placing the colored cards next to the radiant source, they can be held in the free hand. As long as they are held loosely, they have no effect; but if they are squeezed between the thumb and the index finger, they produce on the pendulum held in the other hand, the expected effect.

If **n** is the number of cards which determine the oscillation, **n — I** will be the **particular number** of the radiation.

Under these conditions, instead of seeking an entirely subjective figure **by empirical procedures, it is infinitely simpler to determine the particular number**

itself of an emission source, by superimposing one **A** one, **next** to this source are squares of colored cardboard. If **n** cardboards transform the gyration of the regulated pendulum into oscillation, **n** — I will be the particular number of the radiation observed.

Note that the swing of the pendulum will occur in the direction of **the fundamental plane** of this radiation.

Exercises:

55. Choose from the table on page 96 different substances and radiations; take the series experimentally, that is to say without taking note of the indications in the column of the table “Polarity and Number of series”.

56. Using the neutral spherical pendulum “PS” of precision, note that we obtain the figures of this second column of the table. Note the same result with the Schumfell.

1.1. Then take the particular number of these waves by the method of colored cards.

58. Small cardboard boxes of one centimeter on each side are suitable for laboratory operations; but if you want to use the method for external operations, extinction of harmful waves, for example, you can use small plywood boards of magnetic color (white, red, orange, or very light yellow).

The characteristic number. We will see later, chapter VI . page **183**, that a sheet of fluidic images is composed of a **central image**, which can be magnetic, electro-magnetic, electro-radio-magnetic, or electro-radioactive, surrounded by **two identical groups of** still radioactive images. It will be understood that what is interesting for the researcher is not so much the number **N** of the members of the family, but the number of individuals of each of the .
* . . .

identical groups that surround the central image. It is this number **n** that we call the **characteristic number** of the EM wave. Between **the particular number N** of a radiation and its characteristic **number n**, there exists the relation $N = 2n + 1$.

59. If one places beside a source of emission a number of colored cards **equal to 0** its characteristic number, this emission is extinguished.

What does the characteristic number physically correspond to? Mys tery, until now. Perhaps it will be possible, **one** day, to formulate a hypothesis about it that we will have the joy of seeing verified by the facts !

For the moment, it is not implausible to suppose that this figure is determined by the ratio of intensity which exists between the electric vector of the EM wave and the corresponding magnetic vector .

ONZIEME LEÇON

FUNDAMENTAL ANGLE THE FUNDAMENTAL PLAN

(Refer to “ **Physical Method Dowsing** ”, pages 81 to 85)

SUMMARIZED AND PRECISE DOCUMENTATION

The fundamental angle is none other than the angle made with the magnetic North-South direction by the plane of magnetic vibration of an EM wave when it propagates towards the sky in a vertical direction. This orientation is caused for each EM wave, and consequently for each simple body, by the action of the earth's magnetic field. This action has the effect of provoking for each simple body an emission of specific rays **with** an MS support in a theoretically unlimited vertical plane, called **the fundamental plane**, and around which is formed, as we will see later, a field of reduced dimensions, limited by horizontal and vertical planes.

This field is called **the field of the fundamental plane** and the specific rays which propagate **in it, of** very limited length, moreover, **fundamental rays** of the radiation or of the substance.

These rays were discovered by Abbé Mermet.

The direction of the fundamental rays is fixed and characterizes each wave and each substance; it is that according to which the pendulum oscillates after each series.

The direction of **the fundamental plane of the iron** being that of the magnetic south , **there** was every reason to think that the phenomenon was due **to** the action of ferrestro magnetism . Mr. Engineer Voillaumo **has** verified the accuracy of this hypothesis by observing the displacements undergone by the fundamental plane when the earth's magnetic field applied **to** this body is methodically

varied .

Having determined the direction of this plane for a large number of bodies under conditions of a fixed degree of illumination, he summarized. the result of his experiments by the establishment of a **map of fundamental rays** (whose angles are given in column 5, page 96) having for axis the magnetic North-South direction. By taking the zero in the direction of the North, and by turning in the direction of the needles of a clock, he was able to observe thus, that the diamagnetic bodies (which are not attracted by a magnet) had their fundamental angle located in the North semi-circle and the magnetic or para- magnetic bodies, in the South semi-circle of the Map.

Pursuing research on the same question, J. Correnson was able to realize that the action of the terrestrial electromagnetic field (EM) was exerted, from this point of view, no longer on the substance itself, but on its emission, and that the fundamental angle existed not only for the specific radiations of the bodies, but for any polarized EM radiation. will explain below, convinced that it readers.

Angle fondamental : L'orientation C'est cette théorie que nous du ohm fondamental (ou ongle fondamental) d'une onde électro-magnétique plane, n'est autre que celui du plan de son champ magnétique lorsque cette onde se propage dans une direction rigoureusement verticale. Ce plan, **Pf** bien entendu, ost alors vertical.




Fig. 47

Rayon vertical
effectivement co
temps, déjà, que
olle est blanche,

On conçoit que si cette définition est exacte, il doit exister, en corrélation avec le **Pf** ct, dans une direction qui lui soit perpendiculaire, un autre plan caractéristique correspondant au champ électrique de l'onde verticale. C'est bien, en effet, ce que révèlent les réactions du pendule. Nous appellerons donc ce second plan : le **Plan électro-fondamental (PEf)**.

, Si **Cl.** est l'orientation du **Pf** par rapport au Nord magnétique pris pour origine, .la direction

du **PEf** sera **Cl.** —
Cl. + 90° si olle est

L'orientation du
l'action du champ
conçoit quo l'action
tique puisse la faire
même de celle d'un champ électrique influençant celui de

900 si l'onde est positive ;
négative.

Pf étant déterminée par
magnétique terrestre, on d'un
autre champ magné- dévier et
qu'il on soit de,

l'onde, puisque les deux plans sont solidaires et
invariablement liés l'un à l'autre par la condition do
perpendicularité. C'est bien qui se passe puisqu'il a **6t6**
constaté, depuis long- le **Pf** ost dévié par l'action do la
lumière qui, si intervient par l'action des rayons électriques

ou

magnetic waves which accompany the light waves, and if it is monochromatic, depending on the case, either by one or by the other.

When, therefore, one wants to determine **with precision the fundamental angle** of an EM wave or of a substance, **one** must work in pronounced penumbra.

PEf having only a purely scientific interest, we will stop there as far as it is concerned and we will deal only with the Pf whose practical interest is considerable.

The validity of our hypothesis on its origin and mechanism is confirmed by the following two facts:

I° Non-polarized EM waves have no Pf, because they actually have an infinite number, which is the case with natural light waves :




Fig. 48

**Field of the Fundamental Plane of a copper disk
placed on a horizontal table**

2° Two identical EM waves, but of opposite polarity, have their Pf oriented respectively at **180°** from each other.

Fundamental plane of an emission source : The Pf of an emission source (electricity, magnetism/radioactivity, rays (° human rays, etc.) is a **vertical plane** carrying this source in a single direction and which, in all its extent, emits the radiation of this source (waves ES, MS, RS, <•), vital ray, mental ray, for example). Its dimensions, both vertically and horizontally, are theoretically unlimited.

Fundamental plan of a substance : The specific beam of a simple body is composed of several radiations of different wavelength, but which all have the same objective number of series (therefore same " particular number", same " characteristic number ").), and also, even “ fundamental angle”. We find all these radiations carried by the support MS, in the fundamental plane.

Field of the fundamental plane : Along this plane, the active part of which seems practically unlimited, a kind of box is formed **with** vertical walls having as its axis this plane, for thickness that of the radiating object and limited, in height by horizontal planes, **ot** in length by a vertical plane perpendicular **to** its direction. It is the **field of the fundamental plane** whose main property is to be a space in which circulate specific waves of the object **with** magnetic support (MS) and a filiform emission of rays **a.** , of limited length, which several dowers claim to propagate under a strictly constant angle for a determined substance. (Fig. 48).

Fundamental rays : The specific rays which propagate in the field of the fundamental plane and which have been called **fundamental rays** have a support MS ot of limited length. Experiments are currently underway to determine their direction. Are they parallel to each other or oblique ? Vertical or Horizontal ? **Straight** or curved trajectory ? We are looking for it.

Find 'the direction of the fundamental Ray of an object : We recall that the fundamental Ray can be determined exactly, either using the neutral spherical pendulum PS, or using the " Schumfell" pendulum, which is an active pendulum . However, the Schumfell, developing a magnetic field, the normal orientation obtained by its use will support a deviation of 90° .

Research experimentally, therefore by trial and error, the fundamental radius of a body. — For this purpose, slowly move the regulated pendulum around and




Fig. 49

Conduire le pendule, tout autour, très près de l'objet. Dès que le pendule rencontrera le champ Pf, la giration se transformera en oscillations dans ce plan.

aussi près que possible, d'un objet, par exemple une pièce de monnaie d'argent. Regarder l'objet, surveiller le pendule du coin de l'œil. Dès que le pendule pénétrera dans le champ du plan fondamental, l'instrument qui tournait sur les rayons obliques (entre autres à 45°), émis par l'objet, se mettra à osciller fortement dans la direction du Pf. Constaté et noter le degré d'angle fondamental de la direction fixe obtenue. (Celle-ci sera

different for each different substance or different radiation).

Compare the result obtained with the figure in the Table of Characteristics of Usual Substances, page 96. In the event of a discrepancy, seek to **ascertain** the reasons which

pu le déterminer (réglage imprécis, influences étrangères,

have

heat, light, magnetism, electric fields, etc ..., nearby).

Another process : Walk around the body, or the coin taken for study, a pendulum regulated on MS rays (that is to say on Magnetism: on magnet). This pendulum will swing. As soon as it enters the field of the fundamental plane of the body, it will begin to turn. This observation confirms to us that the fundamental ray does indeed have a magnetic support. Note the direction in which this gyration occurs.

Note that the pendulum, self-regulated for a body, will oscillate if we move it around and very close to this body under the same conditions as before; but that on the contrary it will begin to turn as soon as it enters the field of the fundamental plane.

The fundamental rays have MS support . the

pendulum, settled on substance, oscillating strongly in the plane of the fundamental field, let us interpose between it and the object an iron blade (knife blade) which serves as a screen for the MS rays. we will

find that the strong oscillation stops instantly.

Determine with precision the fundamental angle of a new radiation : Let us obtain, or build ourselves, a graduated circle of 360° , about 0.10 centimeters in radius. (Fig. 50). After placing ourselves in the half-light (room darkened by the measured closing of the




Fig. 50

flaps). so as to eliminate any influence coming from the light, let us orient the long side of our rectangular sheet of paper, in the magnetic North-South direction (approximately 340°). so that the zero of the graduation coincides with the direction of this magnetic North.

In the center of the circle, place as small a fragment as possible of the substance to be studied; and, with a pendulum set on this substance, let us revolve around and very close to this point, operating very slowly. At some point • the gyration of the pendulum will turn into very powerful oscillations.

Let us plan the vertical plane passing through the suspension thread of the pendulum and the center of the circle and note the point where this plane intersects the graduation. We will thus obtain the **fundamental angle** of the substance, which we will then carefully measure using a protractor.

Let us make sure, with a pendulum adjusted to MS rays (Magnetism, magnet), that we have indeed operated in the Pf and that we have not measured the orientation of the Pef.

Let us also make sure that at the place where we operated, there is no electric or magnetic

influence, noting that if we operate on silver we obtain an angle of 90°.

If, instead of the fundamental angle of a **substance**, we want to determine that of a **new radiation**, we will operate as follows: We will determine on a sheet of new white paper, (or better still, a new white blotter in the case of liquid impregnation), an impregnation of this radiation. It will suffice to have placed a body on the above-mentioned sheet of paper for some five minutes for the radiation to persist after having then withdrawn this body. In the part where this body will have made contact with the sheet of paper, therefore in this **impregnated part**, or **impregnation**, we will cut out a fragment the size of the head of a small nail, and we will operate as before, placing it at the center of the circle. But, remembering that we operated on an identical wave, but of a polarity opposite to that of the wave studied (as is the case for any impregnation of the first degree), we will deduct 180° from the angle obtained, so as to place us in the extension of the orientation found.

Exercises:

60. carefully adjusted precision spherical “ PS ” **neutral pendulum** the serial number of different substances, and note the direction in which it oscillates after the last jerk. •
61. Seek as it was exposed on page 113 the **direction of the Pf** of the same substances and note that the two directions coincide in each case if both. operations were well done.
62. Find the **fundamental radius** of a body by turning around this body. by not looking at this body. Turn on the 45°, these tilt **at** the place of Pf.
63. Compare the result obtained with the orientation indicated by the Table, page^ 96. In case of discrepancy. seek to realize the reasons which have determined this difference (foreign influences, heat, light, magnetism, etc.) but also imprecise adjustment.
64. Note that the fundamental ray has a **magnetic support** and that, for this reason, it enjoys from the point of view of the screens the same properties as the rays **at 45°**, stopped by an iron blade.
65. Also try to find the Pf on **MS setting**.
66. Find that the pendulum **self -tunes** for a body. which oscillates above this body. **turns on the contrary** in its **fundamental**

plane.

LESSON TWELFTH

WITNESSES

(Refer to “ **Physical Method Dowsing** ”, p*age 31)

SUMMARY AND PRECISE DOCUMENTATION

We call a **witness** a body identical **to** the one we are looking for and which we use to help **us find** it . This is the simplest and most accurate method of all.

The rod provided with a witness ceases to **react**, except in the presence of a body of the same nature as the witness.

It is the same with the pendulum, regulated with a length of wire suitable for the witness, and provided with a witness held by the same hand (hand which suspends the pendulum), which now turns only on a substance identical to the witness.

This therefore plays a role analogous **to** that of the devices which, in wireless receiver apparatus, only allow waves of a determined length to pass. It 's a **radiation filter** .

If, instead of being held by the same hand as the pendulum, the witness is seized by the free **hand, the left hand**, the reactions are the opposite and analogous **to** those of a self-adjusting hollow pendulum . The suspended mass then turns permanently, and passes to oscillation, as soon as it is on the vertical of the body identical to the control.

For the stick, the witness can be placed either on the fork, or in one of the hands. Thus, if one slips the ring of a bunch of keys into the thumb of the right hand, the stick will rise each time one places the foot above an iron joist of a floor. .

. Exercises:

67. Check this cookie filter property in four different ways:

The witness is held with the right **hand** pressed with the thumb against the suspension stick. Adjusted with a length suitable for the witness, the pendulum turns on the vertical of an identical body, but gives no reaction on another body, even if this body has, with the witness, an almost common adjustment length.

Check this phenomenon with the length of **133** mm. common to copper and silver. • ,

68. Check it with different bodies: tin, copper, aluminium, nickel, lead, etc., by placing the regulated pendulum equipped with a witness on the vertical of these bodies.

69. Check it by placing the index finger of the left hand in antenna or in contact with these bodies.

70. If the witness is held with the **left hand**, the phenomenon is different. The pendulum set for this witness, far from the object, rotates permanently. It oscillates if it is represented above an identical body.

71. Its reactions are the same as those of a self-adjusting hollow pendulum . Thus a hollow pendulum, containing a fragment of gold, suspended vertically from a gold object, rotates. It will oscillate if one holds a gold witness in the left hand.

72. Note that when the pendulum rotates, if we take in the hand that does not carry the pendulum, an object of the same nature as the one we are looking for, the pendulum will continue. to turn, but in the opposite direction, if one is really in the presence of a body of the same substance as the witness. This is excellent control. And the pendulum will pass to the oscillation as soon as it is on the vertical of the body identical to the witness.

It is advisable not to look at the pendulum, because then the action of the visual ray would put the operator **in the same situation as if he presented the pendulum above the object.**

73. Let's take a silver coin in the left hand. Set to silver, the pendulum will always turn.

74. Presented above another silver coin, it will oscillate. .

LESSON THIRTEENTH

INSTRUCTIONS FOR USE OF THE " SCHUMFELL " PRECISION electro-radio-magnetic DEVELOPER, ANTI-IMAGES

SUMMARY AND PRECISE DOCUMENTATION

The essential characteristics of the Schumfell Revealer have been set forth on page 46 of this Course.

The previous exercises familiarized us with the use of the black neutral spherical pendulum PS A few exercises will suffice to quickly initiate the constant use of the Schumfell Revealer, anti-images, Patented SGD

amplifying pendulum . very sensitive, with ample and deep movements, has particular qualities of precision and security, recognized by the most renowned professional dowsers, whether it is a matter of dowsing or teledowsing (Physical or Mental).

Radio-magnetic position : When the median plane of the ring is oriented in any plane, other than that of the North-South magnetic meridian, the developer functions like a magnetic pendulum, specific to the **Mental Method** ; this is how many dowsers, operating according to the mental method, use it continuously (1). That is to say, under these conditions. it gives no indication to the prospector who operates by the exclusively physical method, the one that is the subject of this course.

0) See page 229 : “ P.rœocédés of •the *Mental Method* .

Position électro-radio-magnétique : Par contre, quand le plan médian vertical de l'anneau coïncide à peu




Fig. 51

1^{er} TEMPS : Réglage millimétrique de la longueur du fil de suspension.

North-South magnetic meridian , and that. the device is made to oscillate in this plane, the winding is traversed by an electric current • induced by terrestrial magnetism (Faraday's experiment); thus the Schumfell momentarily becomes < [?] .-electric, and is suitable for ne —re Physical method .

Its operation is the same as that of the black neutral spherical pendulum PS; the same way of suspending it, of adjusting it experimentally or according to the Table of automatic millimeter measurements, pages **96** to **99**. However, it also requires, during perception, the constant maintenance of its ring in the direction Magnetic North-South ; this is called “orienting h Schumfell ”.

Before orienting the Schuinfell by means of its central compass, in the magnetic North-South direction , the operator will check that there are no large metallic masses in the vicinity (iron or u 'i magnetic field,

such as a magnet, or even too close to an electric field constituted by a closed electric circuit). liable to disturbingly influence the needle of the compass of the instrument.- This is easily realized by moving to several places in the same room ; at each check-




Fig. 52

2^{me} TIME: Slight swing of the pendulum in the orientation
Magnetic North-South do the ring.

cation, the compass needle must take exactly the same orientation. This precaution is indicated as an indication, because in practice, the case rarely arises.

We will then correctly orient the Schumfell on the magnetic North-South, that is to say that we will place the blued point of the compass needle at about 3-4()0 from the dial, i.e. about 20° to the left of the letter N (North), which is at 3600, indicating the geographic N,Qrd.




Fig. 53

3^{mo} TIME: Positive gyration of the pendulum, on the silver, maintaining the magnetic North-South orientation of the ring.

Thus oriented magnetic North-South, and regulated, for example, on a silver coin, the Schumfell will turn positive, if the operator's hand is positive.

If he takes too long to set himself in motion, help him slightly by making him swing imperceptibly in the median plane of the ring, i.e. in the direction A, B, C, D, of figure 52, which will be the 3400/1600 direction. Then, while maintaining this magnetic North-South orientation of

the next , it will slowly at its ease, take its normal direction of

gyration, in a more or less short period of time, depending on the sensitivity of the operator, and when the latter has obtained the exact measurement of adjustment some thread.

The use of the automatic millimeter settings of the Identification table is recommendable before the suspension of the Schumfell. Well adjusted beforehand, it immediately takes its normal gyration.

With regard to the precision given by the Table of automatic millimeter measurements, it is advisable, as we recommended for the Spherical neutral pendulum PS, to always take into account the **personal equation** of each operator. If the free and wide rotation of the Schum fell were to be delayed, it would be necessary to increase or decrease, at most by one millimeter, the figure indicated, until one has obtained a maximum gyration. •

CHAPTER IV

SOME MICROPHYSICAL PHENOMENA . of PRACTICAL USEFULNESS

LESSON FOURTEENTH

LICNES AND FLUID PLANS

(Refer to " Physical Method Dowsing)),
pages, 97, 98, 103, 110 and 113)

*

We substitute the expression **fluidic lines and planes for that of dowsing** lines and planes . used by many authors, because we consider this new designation more appropriate.

SUMMARY AND PRECISE DOCUMENTATION

1° **Fluidic planes of the magnetic cross.** The action of de la terrestrial magnetic charge causes in each the formation of two objet orthogonal vertical planes

oriented N.-S. and

N.
magnétique
•
•
E.-O. magnets . which intersect at a point
of this object which does not appear to
be its center of gravity and which it will
be interesting to determine one day.
These planes which are traversed by
specific rays of the MS - supported
object are sensitive to a certain distance
: S
magnétique

Fig. 54

both horizontally and vertically. These are the plans of the Magnetic Cross. The polarity of the MS rays varies in the various branches of the Cross; it depends on the polarity of the body and the position of its ground plane .

These maps can be used after sunset to search for a lost object whose approximate location is known.

Where the compass is deflected by local conditions, the Magnetic Cross undergoes a strictly equal deviation.

2° Fluidic planes of a magnet or of a magnetized body. The action of the terrestrial magnetic charge develops on 'a rectilinear magnet'. a vertical plane which, if the magnetic charge of the magnet is strong enough, can




Fig. 55

be sensitive at very long distances: 30, 60, 120 kilometres. The specific rays involved have a **w** support .

It is the property of this plane that has been put to good use in Le Percepteur Brard et Gorceix. This plane passes through each of the poles of the magnet.

If the object, instead of being rectilinear, is of a more complicated shape, the plane may be double. Example: Horseshoe magnet, nickel pot.

3° Fundamental plane; We will not return to the fundamental Plan on which we sufficiently extended in the 1st Lesson, page **I 1** O. It is a fluidic plane with one direction.

4° Plans photogenic. (See “Physical Method Dowsing”, pages II 0 to 113). Under the electric and magnetic action of light, a body C illuminated by a source S gives rise to the formation of three fluidic planes, one in alignment with the light source and the object, the other two oblique. These planes emit specific rays from the object with various support ES and MS, and RS. The central plane or luminous plane (solar plane, if the source used is the sun) • is much more sensitive than the lateral planes.




Fig. 56

Exercises:

75. Look for the Magnetic Cross of various objects, and of various shapes. For example, let's place any object on a table. Let's set our pendulum to the specific radii of this object: a ^{silver} coin. Let us seek using the compass the North-South magnetic meridian (3400 approximately), passing by this body, and note the existence of the four orthogonal fields of the Magnetic Cross (of different polarity). It will be remembered that these four fields are directed towards the four magnetic cardinal points North, South, East and West.

76. Note that these fields extend very far.

77. Keeping the previous arrangement on a wooden table, which is an excellent conductive medium, let us note, with the aid of the antenna point, that the radiesthesic task of the object prevents us from capturing its images at the level. Note also that it suffices to raise the tip-antenna by a few millimeters to encounter these images.

78. Remember that if you want to easily trace the outline of this spot, you should place the object on a glass plate or a porcelain plate, both of which are poor conductors. For more precision, always use the antenna tip to follow the emission layers.

79. Bury an object in a flowerpot and find that its Magnetic Cross still exists.
80. Have an object buried by an assistant in a restricted space of 4 to 5 m², and find its location using the Magnetic Cross.
81. *M*-ray setting, that the fluidic plane of a rectilinear magnet extends over a very long distance.
82. Look for the fluidic planes of a magnetic body, a nickel coffee pot. for example.'
83. Remember that **solar radiation reaches** all bodies in a straight line. with a **light**, electric and magnetic support. It is also known that when the sun is hidden. the support loses its luminous property, but exerts its electric and magnetic properties, a kind of dark rays sufficient to allow prospecting

-r. Sun




Fig. 57

Perception of solar radiation between the sun and any object

84. In a first process, put a silver coin on a table, so that you can turn your back to the sun on the one hand, and have the coin facing you, at three meters for example. Then set your pendulum on silver, adding a second piece of silver as a “witness”. Try to perceive the ray

ment existing between the room and the sun You **will get there** as soon as your pendulum turns, without looking at the room.

85. This alignment acquired, follow its direction by searching - from time to time if, on the perpendicular line, you do not encounter the CM and CN radiations, whose routings meet on the coin. See that the pendulum will swing when you perceive them as well.

86. By a **second process**, repeat the same experiment, by placing the silver coin between the sun and your regulated pendulum and including “witness”. Note that the swing of the pendulum will occur when you are exactly in this extension.

87. Let's create the “Photogenic Ray” device and **verify** the existence of the **three** Jans fields which propagate the specific rays of coros C.




Fig. 58

Les champs du rayon photogenique

FIFTEENTH LESSON

FLUID FIELDS

(Refer to “ Physical Method Dowsing ”,
pages 85, 98, 99, 104

SUMMARY AND PRECISE DOCUMENTATION

Fluidic fields are spaces limited by defined surfaces in which propagate, following paths which are not always rectilinear, waves serving as a support for the specific radiations of the bodies around which they are formed.

Depending on the nature of their support, we will classify these fields as ES+ or ES— fields, MS+ or MS— fields, fields (JJ . fields with luminous support, fields with nervous support, etc. .. The same phenomenon often giving rise to to the simultaneous production of fields of several species, we will study successively, not all the known fields of a given species, but, on the contrary, all the phenomena giving rise to the production of fields of which we will indicate each time the species and Particularity However, the fields of the same category presenting common characteristics, we are going, first of all, to examine these common characteristics, before proceeding to the successive description of the phenomena themselves.

ES fields . : The specific electromagnetic rays of bodies propagating in an ES field are stopped by an electrical insulator (glass, porcelain, slate, dry paper, etc.) on which they are reflected to take a new direction. This depends both on the form

reflective surface of the surf .

MS fields. : The rays circulating in an MS field appear to be extinguished from their point of emission if, on their course, an iron plate is interposed. No reflection phenomenon is observed. However, the extinguishing action of the iron screen is only temporary, and ends as soon as the plate is sufficiently saturated with magnetism, which happens fairly quickly.




Fig. 59

Magnetic cross from a copper disc

This adjustment was made on the support, that is to say by adjustment on magnetism.

If the setting had been made to the specific spoke, the polarity would have been, for copper or silver, positive. It would have been negative for nickel. Thus, nickel would have given: South+, West+, East—, North—.

Fields w : The rays propagating in a field to are stopped, but, from this screen only, by a barrier of cellophane paper on which they are reflected.

Let us quote again:

1^{oh} The Field of the Magnetic Cross. This field limited by parallel horizontal and vertical planes develops under the action of light around the planes of the Magnetic Cross of an illuminated body. It disappears when the body falls into darkness at the same time as the families of images that form there.

It will be noticed that the branches of the Cross of this field are unequal.

2° The Field of the fundamental Plan. indicated here for the record. (See **1^{lm} Lesson**) .

3 ° "The fluidic field of a rectilinear magnet. It is a cone or a pyramid with a very slightly open nail which develops around the fluidic lines of the magnet. This field is permanent.

C' rays . which escape from each end of a bar magnet, determine, on the side of each pole, the formation of a ^{field} **C(<)** long distance (**approximately** 1 kilometer for a magnet 4 millimeters in section and 0.10 cm in length). limited by the contour of a quadrangular truncated pyramid of very small opening, having as its base the cross section of the magnet. To these two fields of great length (**a** and **b** of figure 60) are added two other small very short fields, **e** and **d**, of the same strength, but more open, from 0.03 to 0.04 centimeters in length only .

In each of these extended fields (**a** and **b**), the **<'J** rays export **MS** rays to which they serve as carrier waves, **MS + rays** on the positive pole side, **MS —** on the negative pole side, and, in each lateral fields (**e** and **d**) of **MS** rays of different polarity whose respective sign seems to depend on the orientation of the magnet.

These **MS** rays, **in** turn, serve as a support for the specific waves of any body included in one of the fields and whose radiation can thus be felt at any point not only of this field, but also of the three others. .

We conceive the use that it is possible to draw from this **property** for long distance perception.

The range of the extended field of a magnet is a hundred times that of its restricted field, which will be discussed in 5°.

Magnetized bodies of a more complicated form than that of the bar magnet give rise to the formation around their fluidic lines of analogous fields. .

4° The fields of the Ray of Union : Indicated here for the record. (See At the 16^{me} Lesson).

5° The restricted fluid fields of a bar magnet : There exists, on the side of each of the poles of a rectilinear magnet, a zone having the shape of a truncated elliptical paraboloid flattened in the horizontal direction, and which we will name **zone of sensibility**, in which propagate the specific rays of any body such as A which enters it (Fig. 62).

This paraboloid is all the more elongated as the magnet **has** more range. It stops abruptly **at** a certain fixed distance **SQ** from the obturated pole (Fig. 62). This field exists for each of the two poles (Fig. 61}. and its common parts constitute a neutral region in which the pendulum swings.

Extended field and rays (d

exporting MS+ rays which, in turn, support the specifi c waves of any body included in one of the fields.

+ ÜN KS! d Lateral field.

! 6111 Extended field and rays (»> exporting
Fig. 60 rays MS—.

Extended and lateral fields of a bar magnet.

The simultaneous formation around a magnet of two distinct radiesthesic fields, **extended field** and **field r** is re i nt encroaching on one another, seemed improbable for a long time; and the different perception of one or the other, according to the observers, was, first of all, attributed **to** entirely subjective reasons (1). It is perfectly explained, today, by the fact that the MS waves, which circulate in the two fields, play there only the secondary role of waves carried, and that the waves-supports which involve them, rays <o in **the** extended field, rays **o** in the limited field, are absolutely different. For an operator who works with a pendulum set on MS rays, the distinction is difficult to **observe** , but if he works successively with a pendulum set first on **tt.** > **rays** , then on **o**, he will obtain with the greatest facility the successive delimitation of the two fields .

This established, it will be understood that if we want, **with** the help of a magnet, to expiate a given direction, that towards which the end corresponding to its Mord pole is oriented , we will have to remove the possibility of an indication from from any other direction (rear, sides, top or bottom of magnet) . We will therefore close the South Pole end with a nickel cap (' . , q. 63) and we will surround the middle of the magnet with a double belt of copper (positive metal) and nickel (negative metal), in order to annihilate the action of fields **c** and **d** (Fig. 60).




Fig. 61

The two paraboloid trunks of sensitivity

d'un aimant droit

(1) The existence of both sides; being known, one can, with a pendulum adjusted to the waves of the prospected body ct. by operating by 6é .:ecillon mental, to eliminate indications which come from one of them.

On the occasion of this study on the bar magnet, let us recall that, on the one hand, the poles of opposite names attract each other and those of similar names repel each other: on the other hand, that it is agreed to call the North Pole of the compass or a bar magnet, the end that is attracted to the Magnetic North Pole of the Earth. This end emits MS + rays.




Fig. 62

Sensitivity paraboloid trunk of a bar magnet




Fig. 63

End capping and middle coating of a bar magnet

But precisely because it attracts the North Pole of the compass, the Boreal Pole is negative and emits MS—rays. Likewise, the south pole of the compass or of a bar magnet, which emits MS— rays, is attracted towards the south of the earth. For the same reason, the Southern pole of the Earth is positive.

It follows that the south pole of the Earth is positive; but the south pole of the magnet or the compass is negative.

We can verify the exactness of the polarities of the extremities

of the needle of a compass, or of the extremities of a magnet, by adjusting the pendulum to MS rays.

In case of failure, carry out the check which can be done using the "polarity indicator" provided by the pocket battery, even when used. The positive pole of a pocket battery is, generally, for French-made batteries, on

la lame la plus courte.




Fig. 64




Fig. 65




Fig. 66

Le pendule étant réglé sur les rayons MS et suspendu sur

the presumed North pole of the right magnet, touch with the middle finger of the left hand the positive pole of the pocket battery. If the pendulum is indeed on the North pole of the magnet, it will continue to turn; if, on the contrary, we touch the negative pole of the battery, it will begin to oscillate.

Since the middle finger is the only neutral finger from the point of view of the emission of the sympathetic wave, which in the other fingers is alternately positive and negative, it is well suited to use it for the polarity check.

6° Aureole of a magnetized body. Indicated **here** for the record. (See Chapter 1, § 3 J.

i" Field of the Photogenic Triangle. Indicated **here** for the record. (See Chapter VI, ERM Images).

Exercises:

88. Place yourself, in the evening, in a room lighted **by** electricity and search with the aid of a pendulum adjusted to the specific rays of a chosen object, placed on a table, the limits of the field of the Magnetic Cross of this object.

89. Note that if the light is turned off the field vanishes.

90. Do the same experiment on the horizontal trace of the fundamental plane of this object.

91. Note that this field is permanent and subsists in darkness.

92. Observe with the neutral spherical pendulum PS exclusively (because it is not magnetic), set to MS, the polarity of the two poles of a straight or horseshoe magnet. (See page 69 : Adjustment to magnetism).

93. Check with the same pendulum, adjusted to ES, the **polarity** of the two poles of a pocket battery, even used. (See page 69 : Electricity setting).

94. Check the correctness of the polarities of the ends of the needle of a compass, or the ends of a magnet using the "polarity indicator" which is the battery to check the magnet, and which is the magnet to check the battery.

95. **Rechorchor** the limits of the sixfold photogenic field of an **illuminated object**.

LESSON SIXTEENTH

THE RAY OF UNION

SUMMARY AND PRECISE DOCUMENTATION

By the fact of a reciprocal action, two bodies of the same substance emit towards each other specific interfering radiations (1) at a particular point on the straight line which joins them. (Fig. 67).

This very important point is called the neutral point. If we suppose these two bodies of very small dimensions, reduced for example, to material points A and B, and that we walk without looking at these bodies a pendulum adjusted along the straight line AB, we find that it turns at each of the points of this straight line, except at point N where it oscillates normally in its direction. This phenomenon is called the ray of union.

$$\frac{A(D)^n}{(M)} = \frac{B(d)^n}{(m)} \quad \text{Fig. 67}$$

If the masses of the two bodies are equal, the neutral point is strictly in the middle of their distance. If they are unequal, it is rejected towards the smallest. Between the masses. M and m of the points A and B, and the distances D and d which separate them from the neutral point, there is no proportionality, but a more complicated relationship represented graphically by a logarithmic curve. This relation, moreover, is independent.

■ ----- i .
(H We give the hypothesis of the interference which conveniently explains the phenomenon, but it is far from being verified.)
dante (I) of the substance of the bodies and depends only on the ratio of their masses.

The ray of union has a very great practical importance in dowsing.

It is used for the identification of bodies, the search for hidden objects, for medical and veterinary diagnosis, and in particular microbial diagnosis, with the pendulum balance for precision dosages and with the BG collector for searches at distance.

But if we examine it in detail, we find that this phenomenon is somewhat more complex than a cursory examination might lead one to suppose. An adjustment on the electricity allows us to observe, for example, that the pendulum rotates positively from A to N. and negatively from N to B.

We are therefore in the presence of two ES fields, of different polarity, having the neutral plane as the common tangent plane.

Figure 39, page 101 of “Radiesthésie Méthode Physique”, indicates for two coins placed on a horizontal table the shape of the vertical 1/2 section of these fields.

The shape of the horizontal section is analogous.

In short, these fields have the shape of two pears joined at their base.

The neutral plane (2) is vertical and its horizontal trace makes with the straight line AB an angle all the more acute as the ratio of the masses differs more from unity.


The specific rays of bodies A and B seem to propagate in each of the two fields in two independent movements and without going beyond the neutral plane.

In addition to the double field ES, there is produced between the two bodies a field MS limited horizontally by vertical planes and of low height. This field consists of a double neutral point.

(1> There is nothing surprising in this:!!Ult, the ray of alone, as we shall see, was, above all, a phenomenon of an electric and magnetic order.

(21 It should be observed that the neutral plane is vertical, whatever ^A it the inclination.>on d:::a right t e AB If this line is itself vertical. becomes horizontal.


If between the two bodies A and B we interpose an electrical insulator, the ES fields are extinguished, but the M.S field which, alone, still allows us to perceive the ray of union at a low height.



ES fields in forming a union ray



Increased range of union radius
by magnet



Union radius 19 long of a conductor

The fields of the radius of union have the shape of two pears side by side and separated by the **neutral plane**. Figure 68 represents the **horizontal** section of these fields; the vertical is analogous.

Exercises:

96. Establish the **radius of union** between two silver coins of the same mass, that is to say of the same weight, and seek the **neutral point**, the pendulum being in the circumstance regulated on the silver.

97. Look for the **neutral point** between two pieces of my silver naie. sizes, therefore of different weights. Observe then **where** the neutral point is. Notice that it comes close to the smallest piece, which represents the smallest weight.

98. To note by means of regulation on electricity (on pocket battery) (1) the existence and! A different polarity of the electric fields (ES), separated by the single neutral plan.

99. Note, after adjustment on the magnetism (on •magnet), the existence of the magnetic field (MS) connecting the two bodies.

100. Look for the two neutral points of this magnetic field.

101. Note that the interposition of a vertical electrical insulating screen (glass for example) suppresses the two ES fields, but leaves the MS field with a double neutral point.

102. Note that the interposition of a metal screen (copper plate for example) leaves the double electric field and its unique neutral point, but that no specific ray of the bodies studied is propagated there (silver in the circumstances).

.11 See! .r paG e 69 '...:s particularités co^rnant Célctric^ and ;e U&me.

103. Note that if this metallic copper screen is replaced by an iron plate, the MS field disappears.

104. • If necessary, attach two silver coins of the same weight to a flat draftsman's ruler (the middle of which will have been indicated by a transverse trait **i**) by means of laundress's wooden clamps. Note that the pendulum will oscillate on the line, that is to say on the neutral plane. Then put this trait **r** on the vertical wedge of a silver object (3rd silver coin of the same weight), in other words slide this third silver object under the flat ruler, at the place of this trait. Note that the pendulum, set to silver, will turn at this point, instead of oscillating, the neutral point disappearing as a result of the **phenomenon of the triangle of union**. This feature brings great interest in prospecting.

105. Let us take the same flat ruler, always having the pieces of the same weight at their extremity. On the middle of this ruler, the pendulum set on silver, will swing, as we have seen it. Let's raise it vertically until it is at the level of a 3rd object, for example copper (copper coin). Note that when the horizontal line meets the object, the neutral point will reappear and the pendulum will begin to oscillate. The main thing is that the 3rd object be of a different substance from the two similar substances at the two ends of the wound ruler. This other particularity constitutes another great interest. practice.

106. Let us place two nickel coins in two identical aluminum boxes (aluminum boxes for travel toilet soap, for example) and make with a cotton (or copper) thread the device **ray of union along a conductor**, and seek the neutral point along this wire which can be arranged in a snake.

107. Using the pendulum, draw the horizontal trace of the fields of the radius of union of two nickel coins of different sizes, placed on a table, at the two ends of a large sheet of white paper.

1 08. Find on the same sheet of paper, the limits of the magnetic field of the ray of union at double neutral point using a pendulum adjusted to MS rays (magnetism). This control has the shape of a rectangle. He is of low height. (Fig. 71) .




Fig. 71

MS fields in union ray formation

To make, so to speak, more visible, the layers of emission and the fields thus perceived either with the pendulum, or with the finger in antenna, it is advisable to become familiar with the use of a **point-antenna**. It is, as we have already said, simply a rod most often metallic, very finely pointed, which one walks by the tip, with the left free hand, very vertically, in the fields in question. . The pendulum, still suspended by the right hand, operates its reactions a little apart, its mass having to be 10 in the body of the operator. and maintained at the height of the tip of the tip-antenna. Thus the beginner will get a more exact account of the modalities of the various phenomena and will be forced to get into the habit of precision.

LESSON SEVENTEENTH

THE RADIO-ACTIVE RADIATION OF BODIES THE SPECIFIC AXIS OF SIMPLE BODIES

(Refer to “ Physical Method Dowsing” pages 92 and 95)

SUMMARY AND PRECISE DOCUMENTATION

As we recalled in Chapter 1, § 5, page 15, radioactive bodies emit three kinds of rays, corpuscular rays α (atoms of helium), corpuscular rays β (electrons). electromagnetic waves γ . The first two are of limited length. The rays γ can be recognized using the pendulum

set to helium (wave associated with the corpuscle).- and for the same reason the rays β using the pendulum set on rays ES (wave associated with the electron).

Among the rays α, β, γ only the rays RS are the product of the disintegration of bodies: rays α, β, γ are the emanation of negative electric field that occurs at the bottom of any radioactive body.

Delivered by itself, the beam of these three radiations presents itself very variably. Months if we act, on radioactive body, a negative magnetic field, separate them. three write-offs

The rays α fail in the fundamental plane substance radioactive, deflected by the magnetic field: the β rays give the corresponding PFE (see 11th Lesson).

The RS rays are analogous to heat (CL rays): up to a certain intensity, they are beneficial. It is only above a certain number that they are harmful. They are emitted by living beings and accompany the vital wave, which seems to prove that they are produced by cellular work. They play an important role in plant life and health.

Emitted by a negative electric field, α rays of dual polarity should be negative. The fact that they are positive, α^+ , constitutes an anomaly, the reason for which has not yet been discovered.

Non-radioactive bodies. As physicists have pointed out for a long time, the so-called non-radioactive bodies are so **to an infinitesimal degree**. They also emanate rays **a**, **y**, but at such a weak intensity that the pendulum **o** has difficulty in grasping them. To separate these rays, there is no need for an extraneous negative magnetic field. The earth's negative magnetic field is sufficient for this.

Under the action of this field, the rays **:J** escape into the fundamental plane of the body, the 2 rays into the PFE. For our readers with a scientific background, we would add that it is in the plane extending the PFE that the electrons set **in** motion by the photoelectric phenomenon escape under the action of light.

The rays **Cl**, of limited length, escape in a single filiform bundle which we provisionally call the **specific axis of the body**. From some experiments it seems to result that this axis, directed towards the sky for the positive bodies, and towards the earth for the negative bodies; makes, for each substance, **a** constant angle with the vertical. But the number of experiments made by us is still too small for us to be able to deduce a law and the identity of this axis with the **Baradat axis**.

Radioactive radiation plays an essential role in the production of AR images under the action of light.

Exercises:

109. Silver has its fundamental plane (Pf) oriented to the magnetic West. Its fundamental angle is therefore 900. Consequently, its PFE is oriented to the north.

110. Observe with a pendulum set to helium that a silver coin, for example, emits rays **a** in the west direction magnetic. With the pendulum

set to ES rays, that it emits these rays in the direction of the North, and if it is illuminated, also in the direction of South (photoelectric phenomenon] (Fig. 72). ■




Fig. 72

Champ du Pf d'une pièce d'argent

de rayons $\hat{U} + j'1, \mathcal{E}_s$ ce qui fait comme nocifs.

111. Add that the rays **j** (**J** are always accompanied **.*** quo physicists consider them ' >

LESSON EIGHTEENTH

AMPLIFICATION BY THE ACTION OF RADIO-ACTIVITY

(Refer to “ Physical Method Dowsing n , pages 92 and 95)

SUMMARY AND PRECISE DOCUMENTATION

If, on a silver coin, for example, lying on a table. we place a fragment of any radioactive body (small piece of pitchblende). we note that the specific vertical rays of silver, emitted by the coin, exceed it by a lot and that. from this point of view the silver coin seems to have taken on the dimensions of a plate.




Fig. 73

Silver coin topped with a piece of pitchblende ore

It is that, through contact with a body. radioactive. the piece of silver has itself become radioactive; it is a phenomenon of radioactivity by influence, or better. by induction.

This experiment shows that reciprocally when an object rests on a radioactive ground, its weight, from the dowsing point of view gets along, is multiplied and it seems much more important than it is in reality. The same is true of its dimensions.

Thus, in practical prospecting, the perimeter of a phosphate deposit located in radioactive ground would be considerably enlarged, and, to avoid miscalculation, the prospector will always

have an interest in making his first soundings in the center rather than 'on the outskirts.

We know that the air enclosed in a closed enclosure ionizes and becomes radioactive. Any object enclosed in such an enclosure becomes radioactive and is amplified (underground cave, deep cavity).

By carrying out the experiment on the same object enclosed successively in boxes of different dimensions, one will note this rather curious fact that the amplification is identical, whatever the dimensions of the enclosure.

For experiments on radioactivity, use a small piece of pitchblende, ore whose study led P. Curie, Mme Curie and G. Bémont to the discovery of radium. This ore is sometimes called péchurane. If this ore is not available, use **betafite**, another radium ore, or granulites and serpentines from Limousin, or titrated solutions of radium salts.

In the absence of one of these radioactive minerals, an empty bottle that has been corked for some time will suffice. The void or hollow is, indeed, radioactive. It is the same for a pocket electric battery. If you use the Schumfell Revealer for the pendulum, it should be remembered that the lead stops the radioactive rays, and constant checks should be made on the settings of the thread adopted. See the Identification table, ion, page 98, the adjustments to the Neutral Pendulum PS and to the Schumfell Revealer, on radioactivity (RS rays). These always positive rays belong to the group of physicists' **y -rays**.

Exercises:

112. Place a piece of pitchblende ore (or péchurane, or betatite, or granulites or serpentines from Limousin, or titrated solutions of radium salts, etc.) on a silver coin and note the amplification of the specific radii of the silver coin, which normally should be limited to the perimeter of that coin.

112 *IJts.* The same result can be achieved as follows: Place the silver coin, which is of positive polarity, on a large copper plate (thick or thin, it doesn't matter), about 30 X 30 centimeters square preferably, and place your left middle finger on the positive side of a magnet, or on the positive side of a pocket battery. • If the body studied (a silver coin, in the circumstances) was of negative polarity, the left middle finger would, on the contrary, be placed on the negative branch of a magnet, or on the negative pole of a pocket battery, even used.

113. The vacuum or hollow being radioactive, observe by enclosing the silver coin in a hermetically sealed cardboard box for a few minutes, that the specific rays of this coin are also considerably enlarged, widened.

114. Do the same experiment with several similar boxes of different sizes, to see that in each case, the amplification remains the same.

SHAPE WAVES

DOCUMENTATION

If we are not mistaken, the existence of **waves of form** was announced for the first time to the public in the work of MH Chrétien : " The Invisible World of Waves ". The question was then extensively developed in the work of MM. Chaumery and Belizal : " The Radiesthesic Physics ".

Each object of a well determined geometric shape emits a special wave, always the same, whatever the substance of which it is composed. It is **the wave of. shape**, attached to the geometric shape.

Thus, a ball, whether it is made of wood, iron, sandstone or ivory, will emit the sphere-shaped wave or SPH rays. It will be the same for the cube, the rectangular parallelepiped , the ellipsoid, the egg, etc ...

The form wave has the particularity that it is generally very powerful, but very short and most often harmful to health. The Egyptians used this last property to defend the tomb of the Pharaohs: the emotive objects having become deformed, the waves lost part of their harmfulness over the centuries.

The shape wave of the plane (flattened rectangular parallelepiped) has this particularity that it differs according to whether the plane is colored with an electric color or with a magnetic color (1). The properties of the latter were used to find the **particular number of substances** .

(1) The electric colors are blue , violet . _ _ _ ^ Magnetic colors , M>nt Je b: a n c , red, orange or taune, green .

One may wonder why a spherical mass pendulum , like our black neutral spherical pendulum PS, is not self-adjusting. The reason is very simple, it is that the PSH wave is too short (7 to 8 millimeters) to reach **the** human receiver. This wave is damped after a few millimetres.

The plane figures, of well-determined regular shape, also give rise to the emission of form waves, such as **the** Pa-Koua and its Yn-Yang.



Fig. 74
The Yn-Yang

It is the same when these figures are applied to non-flat surfaces.

The shape wave of the disk (ray Dq) has the property of extinguishing certain images on condition of being very close to them. (See Chapter VI).

We give below, by way of curiosity, some characteristics of the SPH, Dq and Yn-Yang rays.

SPH rays. Series: —11. $N=I \ln=S$.

Settings: 83.75 — 103.75 — 141.25 — 104.50.

Rays Dq. Series: —1. $N=3 \ n=1$.

Settings: 108 — 121.50.

Rays of the Yn-Yang.' Series: +S.

Settings : 88.50 — 106.75 — 144.50 — 180.50.

The question presenting little practical interest for the beginner, we will not expand further on this subject.

Exercises:

115. An interesting study of the radiation of a wave form on the surface can be made on a drawing which was probably the foundation of ancient Chinese dowsing : the Yn-Yang.

Even if you don't have this drawing handy, you can reproduce it from memory • on anything, even in pencil. It represents like two fish joined together, one black and the other white, each with an eye.

Fine-tune experimentally on the white fish eye above and see the negative gyration.

Then pass over the eye of the black fish and note the oscillation .

,

HARMFUL EARTH RAYS

(Refer to “ Physical Method Dowsing”.
pages 141 to 151, 96 and 97)

SUMMARY AND PRECISE DOCUMENTATION

Since the publication of the work “The Dowsing explained and practiced by the Physical Method”, in 1938, it occurred on the subject of the telluric rays a discovery of great importance. Mr. Francois Orce! found. that underground streams fall into two distinct categories: **arsenical streams** and **hydrofluoric streams**. With few exceptions, arsenic and fluorine are in both cases present at an infinitesimal dose, which chemical analysis is unable to reveal. The subject can be found developed in the works of M. Orcel.

After remembering Pasteur's warning: "With water we drink 90 percent of our illnesses ." note that the vertical radiation of the former (arsenicals) is favorable to the health of living beings. That of the second (hydrofluoric) when it is prolonged for years, brings about very serious disorders in the most resistant organisms. Arsenic streams are closest to the ground; hydrofluoric streams generally flow to a fairly great depth (80 to 120 meters at most) . They are extremely widespread; they are found everywhere. They flow from East to West.

In the present state of science, the harmful telluric rays are therefore of two kinds, the **r rays** generated by radioactive media and the **hydrofluoric rays**, pro-

coming from fluoride-laden streams; both being dangerous only when they reach a certain intensity.

Rays γ (i.e. RS, J , v and **C rays**) can come from:

1° Soil containing radioactive minerals, such as certain marls or ochres, such as radium, uranium or thorium salts.

We must be careful not to build on such land which also emits **$r:J$ rays**.

2° Faults or natural underground cavities. The confined air of these cavities becoming radioactive by ionization, these cavities emit rays α (and in places of the rays **'1.**).

If their walls or bottom contain salts of radium **or** thorium, they also emit rays **?** and r , also harmful. •

3° Man-made underground passages, galleries or rooms ; these underground passages only emit rays γ (J , α , β ;)•

4° Peyré rectangles; same show. (See page 19).

5° Of radioactive underground streams. These are streams that have crossed radioactive ground and are impregnated with their emission. As for 2°, these rays can emit in addition to the v rays. rays $i e > -;$.

The blood of persons having stayed in an irradiated zone being impregnated with its emission, one will be informed of their existence by presenting above their hand successively a pendulum

regulated on rays **(1)** or on fluorine.

If the test is positive, we will search for the area, trying to capture the emission from the area with the free hand in the antenna and a pendulum adjusted to that of the two supposed radiations (**J** or fluorine). Once the location has been discovered, it will be a question of eliminating its emission at least at the most dangerous points.

There is always an interest in installing the elimination devices

harmful rays in a cellar or basement, because there they usually cannot disturb anyone. er protection benefits all floors of the house.

The elimination processes are, to our knowledge, three in number: the tip, the antenna and the screens.

1 ° The vertical point directed towards the ground. It has the effect of capping, within a certain radius and up to its limit of vertical action, all the radiations it encounters and to eliminate them through his head.




Fig. 75

**Pièce de mon-
naie surmon-
tée d'un bou-
chon et d'un
clou**

If above a silver coin (Fig. 75) we place a nail driven into a cork so long as to keep it vertical, we find that the silver coin does not emits more rays specific to silver and that these escape entirely through the head of the nail. If, instead of harmless rays, the point captures harmful rays, it is advisable to direct them towards the ground by ent)uring the head of the nail with a piece of iron wire passing above this head and whose extremities are directed towards the ground. This avoids the harmful effect of the numerous images that the head of the nail emits while it is not buried in a dark place (cellar).

If it is a question of protecting a room in an apartment and when the circumstances do not allow the nail to be driven into the ground, which preserves the upper floors, one can contain to plant it in a small square board, 0.15 x 0.15 centimeters.

2° The antenna. This is a metal rod surrounded by a wire wound in a spiral, the two ends of which should be directed towards the ground.

The attractive power of the antenna acts up to the limit of the horizontal rays which it emits. It is for this reason that the horizontal antenna seems more advantageous than

the vertical antenna, contrary to **what** is said on page 46 of " Dowsing Method Physics". (See also in "Physical Method Dowsing", page 67, Fig. 22 **and** pages 96 and 97).




Fig. 76

3° The screens. The screens are, depending on the case, either reflectors or radiation extinguishers. Thus, black bodies extinguish all light rays; opaque bodies reflect them .

Amont	Protected Part	.nails
_____	— - -	• Downstream

Fig. 77

Protective dam against the effects of a harmful stream

RS rays are stopped by lead and the color blue or yellow.

Rays **0** by the colors green or black.

The rays v by an electrical insulator and the colors . •
storm or black.

The rays \ by an electrical insulator and the colors yellow, green or black.

Rays **a** by lead , pure aluminum (or by a salt of this metal, such as a terracotta tile).

Clay is known to be a hydrated alumina silicate. The same applies to cement and, consequently, to cement paving.

We have not yet found a screen for the **j** and - rays. .

In each particular case, the operator will choose the most practical elimination process.

La Pointe is suitable for small areas (Peyré Rectangles, •hydrofluoric or radioactive streams).

On the vertical of a harmful stream, drive spikes (large nails) 0.15 centimeters in length, 0.30 centimeters from each other. The dam thus formed. protects a zone AB, upstream of the device, but not **downstream** ; it is very important to know. Beyond points A and B, the unwanted radiation continues (Fig. 77). If the stream is less than a meter wide, a single spike will usually suffice in both cases.

In other cases, the antenna that can be extended as desired may be preferable. It will have to be oriented N.-E., S.-W. or N.-W., S.-E. to give it its maximum effect. In the N.-S. and E.-O. its attractive power is nil, because it does not emit rays **c.)** in **these** directions.

In the case of a room in a house built on radioactive ground, it will be necessary to pave the cement floor without joints, to stop the a-rays , then cover it with a carpet of linoleum or balatum, electrical insulation, which will stop the radiation. ' / and X rays . Under this mat lay a layer of blue paper to block RS rays, surmounted by a layer of green paper to block **0 rays**.

The effectiveness of the process proves that a colored body is not only a body which eliminates certain colors of white light to give our retina the impression of nuance, but that it also has other properties which persist. in the dark, and this is not one of the least surprises that the study of microphysics has in store for physicists.

It should not be believed that the suppression of the cause suffices to suppress its effects. These, when **they**

prolonged for several years, are profound and confound **doctors** .

Their recovery requires long and continuous care.

First of all, the body seems to resist and the discomforts are infrequent and go unnoticed. After 7 or 8 years, serious ^{A disorders appear for the first time;} the following year they become more serious, then, all the organs being affected, there is a rapid breakdown in a few weeks and the doctor sometimes finds himself face to face with unexpected problems.

When the above will be known and applied for several years, three quarters of the diseases which afflict our poor humanity will disappear, ' the average life will be prolonged and this is undoubtedly the compensation that Providence reserves for the future to the trials of the present time.

In support of what we have just written, let us quote some cases that we certify authentic, of people victims of harmful rays.

M. A ..., a civil servant in a city in the Paris region, occupied a room **there** in 1923, in one of the corners of which was a Peyré rectangle. Bad luck would have it that he installed his bed precisely in this angle: so that each night he slept on a pillow traversed by rays ; • During the first years, he did not feel any apparent trouble, at least to his recollection; but from 1926 he had to undergo regular surgery every two years for a fairly large growth in his left ear, which was qualified as an epidermal plug by the specialist.

In 1930, having gone to the South, he experienced each time he i
exposed himself to the August sun with serious disturbances, dizziness, fainting, '
nausea, etc., to the point that he had to compel himself to remain perpetually ;
in the shade In 1931 the disorders of the brain became very serious, at the j
point of causing serious alarm to the attending physician. This one, |
after various analyses, diagnoses the disease including the symptoms |
they came closest to those he observed, and performed a cerebrospinal fluid puncture on the patient, which saved him.

Finally, in 1934, Mr. A having made the acquaintance of a dowser f
who informed him of the real cause of his illness, changed his bed from place to place. But I
the disease in the left ear had continued to so I
develop, and in 1936 he had to **be** operated on for mastoiditis. Since then *his* health has left k
nothing to be desired.

Another example. M ¹¹ 'B. ., annuitant in a small town in the South ,! was enjoying excellent health when she had the unfortunate idea of * changing rooms. The one she adopted was swept away by . oblique harmful rays coming from a nearby underground cavity. After a few years she died of cancer in the left breast, which she had refused to have operated on.

M^{TM°} C ..., a grocer in a village in the South-East, had, what she did not suspect, her counter crossed by the vertical of an underground stream carrying J-rays .After a few years, she was affected of phlebitis. Almost at the same time, his son, living on the floor above, whose bed was crossed longitudinally by the vertical of the same stream, was attacked by the same disease. The attending physician found the case so extraordinary that he thought he should report it in a report **to** the Academy of Medicine.

A dowser having heard of the case, extinguished the pernicious radiation by the method of nails driven vertically into the ground. Eight months later, the two patients were definitively cured and the illness did not reappear.

Mr. D ..., a farmer, had received the kick of a horse in the stomach, the effects of which were completely cured by intelligent and assiduous care. He had completely forgotten this annoying adventure several years ago when he changed farms. Unfortunately, in the room he chose in his new home, his bed longitudinally smelled **of an** underground stream impregnated with **y - rays**. excruciating suffering.

In a small town in the Vaucluse, two families succeeded each other in the same apartment, which the first occupied for three years and the second for six years. This apartment was bordered by the vertical of a hydrofluoric stream 0,120 **meters** deep . In the composition of each of the two families was a little girl, one ten, the other eleven. Each of the two children suffered from growth disorders which the doctor was unable to **control** . These troubles only ended each time when each family left the apartment.

Among other misdeeds, the vertical radiation of fluoride, produces in the persons it affects for several years, the destruction of red corpuscles, anemia, muscular weakness, extraordinary weight loss, hepatic troubles and the weakening of the heart. These ravages are then very long **0** repair. Fortunately, nothing is easier than to protect oneself from it on **the** condition of going **there in 0** time.

Exercises:

1 i 6. To search for the presence of harmful rays, the easiest way is to adjust the pendulum to radius **U** (the most convenient adjustment is that of 138 mm.).

If the pendulum swings over the suspected area, chances are it is emitting harmful rays. To check it, try the '*i spokes*' (adjustment : 125 mm. 50) and the ζ SPOKES (adjustment : 160 mm. 50). Finally, try Fluor radiation (adjustment : 155 mm.).

117. Research on people: Present above one of their hands the pendulum set successively on rays 0, ν , ε , as well as on the radiation Fluor. If one of these people has stayed in a zone of harmful rays, his blood being impregnated with them, the pendulum will turn on one of these four radiations.

Fig. 80




Fig. 78

Tracé sur le terrain
d'un roctangle Peyré




Fig. 79

118. Research on harmful places: Repeat the experiment on the ground or the room to be studied. With the pendulum set to one of the supposed harmful radiations (**J** or Fluor), search the area by picking up the harmful emission with the free hand in the antenna.

Trace the limits of this area, marking on the ground the points where the pendulum stops turning. If it is a Peyré Rectangle, the union of all these points will form a rectangle. (Fig. 78).

119. Elimination: Neutralize a harmful Peyré wave by driving a large carpenter's nail (or placing a large nail in a stabilizing plug) towards the center of the drawn rectangle.

120. If the harmful area is close to a place where one is used to parking, it is advisable to avoid that the " images " of the harmful rays, which concentrate on the head

of the nail, cannot also inconvenience (images of the Magnetic Cross, of the solar ray of the nail, etc...). To do this, wrap a wire under the head of the nail, pass it over the head of this nail, and bury it in the ground. This wire will catch the harmful rays escaping from the head of the nail and conduct them to the ground.

121. If it is a stream impregnated with harmful rays, recognized according to one of the aforementioned settings, operate as described on page 157 , remembering that the defense is only valid for the part of the stream upstream from the nails. Protect against the " pictures" of the head of these nails, as previously mentioned.

122. If it is a large area, cavity, fault,. underground, operate with a horizontal antenna that is moved so that the irradiation is completely captured.

LESSON TWENTY-FIRST

THE SEARCH FOR AN UNDERGROUND RIVER

(Refer to " Physical Method Dowsing " , page 160)

SUMMARIZED AND PRECISE DOCUMENTATION

Notions of geology-hydrology : To better understand the reactions of his pendulum, which can only be jodic and all natural, the operator will first benefit from realizing **the external aspect** of the region to be prospected. A few notions of • geology-hydrology can only be useful with regard to the succession and permeability of soil layers.

The origin of water currents is most often rain. or the consequence of melting snow. These waters seeping into the earth become subterranean. As soon as they encounter an impermeable layer (limestone or clay), they accumulate in pockets or layers, or else, sliding on this impermeable layer, flow in numerous more or less important currents. Stagnant pockets of water are difficult to capture. On the other hand, **moving** groundwater emits its waves with greater intensity, and escapes in **springs** on the surface of the ground. It is these moving waters that particularly interest the dowser.

In certain coastal regions of southern France, it should be noted, for example, that **surface waters**, i.e. those close to the surface of the ground, follow a course from the northwest to the southeast . East. It is then sufficient **to cross-** cut these invisible currents perpendicularly, that is,

that is to say to walk South-West-North-East, or vice versa, to raise the rod or turn the pendulum, either on the great first parallel (rays at **45°**) or vertically of these waters (current properly said). The compass attached to the Schumfell Revealer will inform the operator of the position of the cardinal points.

As for the **deep waters**, the best, we will find that their currents follow the slightest resistance . As a result, they do not necessarily follow the slope. çJu soil; this is why they do not meet each other, but often cross paths, either by mixing or by avoiding each other, at different levels.

In any case, the real inexhaustible and copious sources come mainly from the mountains, not from the valleys. There are springs in **the free state** (real underground water currents), sometimes considerable, springing from the sides or from the top of the mountains. These are the ones that should preferably be captured, by channeling them, if necessary, to homes.

We will prospect less, if we can, the valleys, the foot of the hills, where we will most often find only water in **the pellicular state**, forming wetlands, not present. as low runoff or inconsistent water.

It will also be remembered that:

1 °A **synclinal** terrain . (see “ **Dowsing Physical Method**”, page 161) is formed by the collapse of geological layers inclined towards the bottom of a valley. There is the greatest chance of finding water in this valley, especially on the surface; however, this water will generally be of poorer quality than in an anticline.

2° An **anticline** is formed by a sort of convex fold in the surface of the ground, the geological layers of which diverge, drawing the water received away from the bottom of the valley. However, at a certain depth, an anticline can provide interesting siphoning sources by their abundance and their quality of very filtered potability.

3° In a valley where the layers are inclined on only one side, **there** can be water only on the plunging side.

4° Four meters is the minimum depth that can be required of a spring.

there

From what precedes, we can find ourselves in the presence: either of an uninteresting capillarity, or of an underground watercourse (stream or river) of more or less important flow, or of siphoning water (griffon or natural artesian well), or a crossing of two superimposed streams.

And in the most interesting cases, the watercourse chosen will have to be considered from its points of view: drinkability (drinking, watering) or harmfulness (polluted water, colibacillary, etc.), or even mineral points of view, thermal, etc... Also draw inspiration from recent observations, revealed by **M. Orcel, with regard to dowsing** arsenical or hydrofluoric streams (page 23).

apparent watercourses , on the surface of the ground, even. with an exactly tuned instrument. Indeed, he would quickly be discouraged: he would find that in the open air the vertical rays and those at **45°** of these streams and even of these strong rivers are hardly perceptible for an operator who is still little trained . On the contrary, streams or underground currents **emanate much** more

powerful waves, which with a little practice, he will perceive more and more clearly.

.

Exercises:

123. detection in the field: Set the . Schumfell or the rod on the quality of water sought: drinking water, for example, by presenting the chosen instrument on the vertical of any container, filled with the same quality of water as that desired.

Provide the instrument with a **witness**, also with the same water. It is on the witness that the adjustment will be made most exactly.

There can be no question of traversing square meter by square meter, a field of several hectares, pendulum or

wand in hand. Therefore, based on the known • geological information, first estimate whether there may be water in the direction where the discovery of such a water current could be most useful. Walk slowly in this direction, but in such a way as to cut this current perpendicularly at several points. Then, recognize the direction of the current (downstream being the side towards which the stream descends ; upstream, the side from which it descends); perceive the • underground banks by delimiting them over a certain length; finally, determine the points of greatest flow.

But the operator may be deprived of any geographical indication •. In this case, he will go to the limit of the land to be explored, at a place from which he can encompass, at the best angle, the entire useful surface to be excavated. From this place, he will be able, by pivoting on the spot, to study successively different angular strips of ground •, taking the starting point at his feet and moving away 25 meters in 25 meters. He will be put down marks by an assistant, with each reaction of his detector. Thus, quickly, the dowser will give himself a knowledge of the general direction of the various underground currents present.

Some dowsers, skilled with a rod, consider it preferable to first roughen the ground with it, even if it means then using the Schumfell for a more precise study of the water points found.

Be that as it may, this roughing out obtained, it seems to them necessary to check the results obtained with the stick at the anti-image Schumfell, to eliminate possible errors due to possible images.


- This control will be carried out as follows: Help yourself:
 - of the digLal ray (index of the left hand in antenna);
 - the setting of the magnetic North-South oriented ôchumfell ;
 - of the witness (small glass tube stoppered, filled • with drinking water, attached parallel to the hanging rod of the pendulum).

A second check will consist in carrying out cross-checks on the best water point found. Moves- •. Successively in two different places, we will operate two other similar "aimings" on this water point. The intersection of these three cross-checks will mark in a certain way the position of the water point in question.

' **Detection on the water point itself** : The dowser, having then moved vertically to this water point, will determine whether it is a current or a griffin.

In the case of a current, he will first delimit the shores by staking them out over a distance of 50 or 100 meters •.

To better perceive the vertical rays of each bank, it is advisable to bend down or even to squat, the base of the Schumfell at approximately 5 centimeters from the ground, the index



watercourse

Fig. 80a _

with the free hand in antenna, searching by touching the ground: you can still, for more convenience, use a cane-antenna to avoid bending down.

The banks being marked out, the recognition of the direction of the current will be a confirmation of it; the Schumfell will turn downstream, it will oscillate upstream. The wand will raise upstream and lower downstream.

In many cases, the delimitation of the banks can be facilitated by the preliminary recognition of the parallels, more or less perceptible, and especially by the two large parallels, whose emissions are more powerful, when the currents perceived are of some importance. It can happen that the great parallels prove to be more powerful than the

current itself. Sometimes, it will be noticed that the great parallels and especially the parallels are not always perceptible when it comes to small underground streams; only the vertical rays of the banks, and of the current, the being. Check the accuracy of the various reactions thus obtained by walking in front of you, by renewing them by walking backwards and with your eyes closed when operating with a stick.

A water point being known, and the dowser having gone there, he will step back 20 to 30 steps or more, and return, pendulum in hand, towards this point, walking slowly and perpendicular to the current, swinging the Schumfell in the plane of the North-South magnetic meridian, and flowing towards the current. The pendulum will stop abruptly as it passes the large **parallel** and perhaps also of the six other parallels, resuming its swing between each of them, to end in the vertical rays of the first bank, on which the pendulum will swing parallel. On the current itself, the pendulum will turn with more force, reacting in the same way as before on the second bank and the parallels and great parallel on the other side of the current.

This will then be the time to check the series number of the water, which must be +2. according to the fine-tuned shaking method.

All these operations must have been carried out continuously with the witness. After each reaction of the rod or the pendulum, desire to impregnate these instruments, as well as the hands. (See page 40).

On reading the above procedure, the beginner may believe in a certain complexity, but, in practice, this process becomes child's play after several tries.

We have just indicated, for the good rule, the details of the process of research on the ground, we must however add that any practiced dowser, simplifies and considerably accelerates these operations of prospecting, whereas it is on the ground. ground, by resorting to work on plan (teleradiesthesia). It is, in fact, much easier and faster to scan, with the antenna tip, a fraction of a sheet of paper, placed on a small board.

countryside, than to slowly survey the land, of which it represents the reduced dimensions and layout. A quick sketch, even in pencil, as long as it keeps the proportions and is magnetic North-South oriented, will allow you to search and re-search all the nooks and crannies several times, avoiding • any fatigue by walking if you are in the countryside, and without moving a piece of furniture if the search is done in a living room.

' Flow

124. Having proceeded as stated above, the dowser will be certain of being able, in complete safety, to have an excellent well drilled. However, before having this work rectified, he must seek to obtain it at the lowest cost, by knowing the greatest flow and the least depth.

He will therefore have to wonder if this found place has a sufficient flow first, and if he could, next, or a little further, along the watercourse thus delimited over a certain length, locate a place of higher flow. It is by going down the direction of the current that he will go

account of the greater or lesser intensity of gyration of its pendulum, informing it, by the physical method, of the successive points of greatest flow. .

Yet the process of " mental calculation " is too commonly used not to recall it. It consists simply of a preliminary mental convention of measurement, made by the pendulum, with the subconscious, which one questions. Having agreed that each gyration on the vertical of the water point will represent a liter, or a decalitre, or even a hectolitre, you will raise vertically even pendulum in gyration. It will start to oscillate first, then, to turn. You will then count how many gyrations will be made until the stop. This figure obtained will give you the flow.

The griffin. — This question of flow and pressure leads us to consider the case of the griffin, or the natural artesian well. When underground water is under pressure, this means that it receives by communicating vessels, weights which push it to rise. But it is prevented from doing so by the impermeable roof of clay, rock, etc., which compresses it.

The pendulum signals ¹ this state in the form of a column of force, called a griffin, which rises vertically above the real water, and rises to the level where the water will rise, as soon as a drilling has opened a passage for him and set him free.

Capturing siphoning water is tricky and often discouraging; it is better to

~~focus on finding more or less horizontal water currents.~~

The depth

125. Over a length of 50 meters, we have staked out an underground water current. But the water level of the latter can be at very different heights, depending on the impermeable layers of ground followed by this current. To drill at the lowest cost, it is therefore necessary to determine the place on these 50 meters, where the catchable water will be closest to the ground.

The dowser will have, for this purpose, to assess the depth of the water every 5 or 10 meters along the course. But the slightest depth thus found may still be only approximate, and he will follow the good advice of Abbé Bouly, who suggests giving the well-digger a measurement only as follows: "You will dig to a depth of 40 meters, plus the thickness of the "compact soils ". If, therefore, the well digger encounters a few meters of clay during his work, he must add these to the 40 meters indicated by the dowser.

Prospecting in general, whether for a logical mineral vein, a lost object, or any other radiation, being done, except for a few details, in the same way as that for a stream of water, we We will not dwell here on these different subjects, inviting the reader to refer, for particular details, to the corresponding chapters of the work of " Radies.hesie Method Physics".

CHAPTER V

THE RADIATION OF ^{A.}AND LIVING RES

(Refer to “Physical Method Dowsing”, pages 203 to 211)

SUMMARY AND PRECISE DOCUMENTATION

For a long time it was believed that the human body only emits radiation of a single species and Mr. Henri attributed a wavelength of 10 microns to what he called the human wave. It is the engineer Voillaume who deserves the honor of having revealed that radiation is not unique, but multiple, and that several of these elements are common to animals and plants. He even specified its characteristics.

Among these **human rays**, some are of double polarity, as well in the man as in the woman; others are positive in males (H +) and negative in females (F—). It is by these signs that they are designated in the table below of their characteristics.

(See pages 65 and 66 for details on Human Polarity).

CHARACTERISTICS OF HUMAN WAVES

allowing them to establish their identity card using the precision neutral spherical pendulum PS
or the Voillaume pendulum, or the Schumfell pendulum

Designation of the wave I	Has and file n 1	Polarity and Digit d:series O	Special Numbers N	Number of Characteristics n S	Length of wave in microns »	Successive millimeter adjustments of the suspension wire 7	Fundamental Angle (D)
Nerve ray... (digital, visual) vital wave.....	H+ F-	+5 _	5	2	0.790	204.80-173.90-144-114.50-87.10	142°50
Sexual wave....		+H	He	5	10,515	201 - 184.50 - 168 - 152 - 135.90	326°50
Human wave. . ;		+ 3	3	1	9.954	179.75- 149.10- 134-! 18.85	26°00
Species wave...	H+ F-	+ 19	19	9	9.934	184.80-165.50-146.60-123	311°30
Mind ray....		+ 7	7	3	10,360	To be established experimentally, on each subject	28°00
sympathetic ray		±19	19	9	9.915	195--178.75-162.75-147	273°30
Personal wave.	H+ F-	Features vary on each wounds depending on the person. To be established experimental- subject					■

(1) The fundamental angle can only be measured exactly with neutral pendulums: pherlquCfi PS . or Voillaume. alone, at the cxc'.us'on of the active pendulum Schumfccll. The sign ± \"eould mean plus or minus: 11 applies to ondCfi with double polarlU. sometimes positive, sometimes negative.

VINGT-DEUXIEME LEÇON

NERVE

RAYS VITAL WAVE — SEXUAL WAVE

HUMAN WAVE

SPECIES WAVES

(Refer to " Physical Method Dowsing >>
pages 204 to 205)

SUMMARY AND PRECISE DOCUMENTATION

Nerve rays : Nerve rays escape from the ends of the nerve filaments (digital rays, visual rays). They spread over a very long distance. Some are positive, others negative. They are identical in animals. They are accompanied by rays (•). The **digital rays** provoke the emission of the **antenna ray** and serve as their support.

Their intensity varies with individuals.

From the electrical point of view, the five fingers of the same hand are of the same polarity. From the magnetic point of view, they are all negative, whatever the hand, and that as well in the man as in the woman.

But from the point of view of the emission of the **nervous waves** and the **sympathetic** waves (which essentially interest the student), their polarity varies, as we shall see, **and** sometimes, in a different way, according to the individuals.

Indeed , the fingers of the hand are the point of departure of both **nervous and sympathetic** rays of different polarity. The former extend very far; the latter are short.

If, in a man with a positive operative hand (the one holding the pendulum), we present a pendulum, regulated on the sympathetic wave, successively above each one. of the five fingers, we generally notice the following polarities and intensities of movement:

Positive male hands: Thumb	/• .	+ 50%
Index	—25	%
Middle finger	—50	%
Annular	+50	%
Auricular . . .	—25 /"/"	
Negative male hand: Thumb.	—50	%
Index	+25	%
Middle finger	+50 /"/"	
Ring finger	—50	%
Auricular . . .	+25	%

It follows that, theoretically, to obtain the maximum force in the maximum gyration, it would be advisable, generally , to hold the pendulum by two fingers of the same polarity, controlled, that is, in the example above, thumb and annular.

However, in practice, we have found that many students have more certainty in the movement of the pendulum. by adopting, for the positive operating hand, either the thumb and the index finger, or the thumb and the middle finger. Each student will make a personal study in this respect, as soon as he is somewhat exercised in the practice of the pendulum.

It is recommended, in the search for polarity of the fingers, to avoid the phenomenon of impregnation by taking care, in principle, after each operation, to give a jerk to the pendulum.

Vi<a'e wave : The vital wave, always positive, is identical in man, animals and plants. It is only emitted by healthy organs. (This is what often misleads people who look for deaths on photographs : their pendulum, not adjusted to the vital wave, rotating positively, they deduce life from it, hence the error. For research on photographs or writing, tune to the nerve wave, the human wave, or the personal wave, discussed later) . The vital wave is extinguished with apparent death. Its intensity varies with the vitality of individuals.

Sexual wave : The sexual wave, with double polarity, is identical in animals and plants. It escapes from the reproductive organs, positive in the male, negative in the female. In being. human, its emission is made in two very slightly open cones, one almost horizontal, the second whose axis falls

0.25 centimeters, 0.35 centimeters in front of the feet. In man, both cones are full; in women they are hollow.

Human wave: The human wave is a species wave, which characterizes the human species. It persists after death, even in the bones and ashes of cremated corpses.

Species wave: In animals, the species wave corresponds to the human wave. It is common to all animals of the same species: canine wave, porcine, galline, etc... Animals, such as the mule, the hen-turkey, emit the waves of the double species to which they belong .

Exercises :

126. Digital nerve rays : (Antenna radius): <
 Note the different polarity of the waves coming out of the . I
 extremities of the five fingers of the left free hand, hori
 zontal
 zontally elongated, the pendulum being set to the radius j
 nervous. you

127. Place the pendulum (adjusted to the nerve ray) in the path of an animal's gaze and observe that the pendulum rotates. Preferably, put your hand on the air on this path.

128. Let us place ourselves 5 or 6 meters from a copper object, holding in the right hand a 'Pendulum regulated on this metal.

Let us extend the left hand towards the object, the pendulum will turn. Let's fold the arm, the gyration will stop.

129. Let us extend the arm a few meters above the object in order to meet its vertical tablecloth, the pendulum will turn.

130. Likewise, at a certain distance in the path of the rays at 45°.

131. us ask an assistant to reflect the vertical tablecloth by means of a porcelain plate, let us extend our arm in the direction of the reflected tablecloth, the pendulum will turn.

132. Let's set the pendulum on electricity and reach out for a cloud. If this cloud is charged with positive electricity, the pendulum will turn in the direction + : if it is charged with negative electricity, it will turn in the direction .

133. Visual nervous rays : Let us place ourselves in the same position as before and look at the object, the pendulum will turn: let us stop looking at it, it will oscillate.

But it will also rotate, if we look at its vertical line some distance above it, and at the **45° rays** coming from it. Let us close our eyes, the gyration will cease; let's open them, she'll start again.

134. Vital wave : Present above a plant the pendulum set on the viial wave, and note that it turns.

135. Sex wave : Find the sex of the eggs using the pendulum set to the sex wave and a “polarity indicator”. Take care to de-impregnate the pendulum frequently after each operation.

136. Note that there are unfertilized eggs.

137. Note the emission of the sexual wave by the flower.

138. Waves of species : Search by mental selection, on a second degree impregnation on paper, the characteristics of the wave of species of the various animals.

139. Note that the mule emits both the wave of the donkey and that of the horse.

TWENTY-THIRD LESSON

THE MENTAL RAY

(Refer to “ Physical Method Dowsing ”, pages 56 and 204)

SUMMARY AND PRECISE DOCUMENTATION

Mental ray : How can the soul, an immaterial principle, act on the brain, a material organ ? This is a problem that will no doubt be hidden from man for a long time. Let's not try to unravel this mystery. But what **is**, of certain. it is that thought causes a movement of the brain, a physiological phenomenon, and that this movement triggers an EM wave, a physical phenomenon. This wave is the mental ray. The mental ray is identical in man and in the higher thinking animals.

If we collect it by impregnation on paper placed opposite the temples of an animal, we find that it is identical and in all the animal species capable of thinking: higher animals and insects.

When two ants meet and put their 'antennae in contact, the mental ray of one enters the brain of the other and there determines the movement which, by a mechanism inverse to that which we have mentioned. provoke thought.

The mental ray extends **to** considerable distances. We only know of one screen: porcelain.

When we think **of** an object, the electromagnetic wave resulting from the movement of the brain produced by the action of the intelligence is automatically directed towards this object and brings back the specific waves.

Let us, for example, set our pendulum on silver, and place a coin of this metal on a table. No.re pendulum in hand, so away from the coin, let's turn our backs on this table and close our eyes. We will find that each time we think of the piece, the pendulum turns, and that on the contrary, it oscillates, before stopping, as soon as we think of another object.

M. Voillaume, who has specially studied this ray, has determined the starting points which vary according to the position of the body studied in relation to that of the operator and one of which is, in most cases, a special point from his temple closest to this body.

He was able to reflect, refract and polarize this ray, whose properties are no longer in doubt today.

If we think of an object enclosed in an enclosure impermeable to specific rays, but accessible to infra-red rays, the pendulum set on this object will turn. If, on the contrary, we mask the object with a porcelain screen, the mental ray not being able to reach it, the pendulum will oscillate and then stop. .

The use of the mental ray. is the principle on which rests the dowsing by the Mental Method and the Tele dowsing.

Mental rays are always accompanied by (•) rays. .

Exercises:

140. Mental ray: Let us place a silver coin for example on a table and turn our back to the table. Then let's set our pendulum on money. Then think of the room, the pendulum will swing. Stop thinking about it, it will oscillate before stopping.

141. Let's place a screen of glass, paper, cardboard, natural silk, etc. on the piece. This screen will have no effect **on** the phenomenon.

142. Let's place the coin under a porcelain plate , however **much** we think about it, the pendulum will not turn .And if we prolong the experience, we **soon** feel a headache determined by the prolonged reflection of the mental ray, which will not have been able **to reach** its objective.

143. mental wave selection exercises . On a sphere, for example, distinguish, if necessary by study on an impregnation on paper, to the second degree, the wave form of the various specific waves emitted by the various constituent elements of this sphere, which can be wood, marble , of a metal alloy.

- 144.** Look for the characteristics of the wave form.

LESSON TWENTY-FOUR

THE SYMPATHETIC RAY THE PERSONAL WAVE

(Refer to "Dowsing" Physical Method,, iVages 204 to 206)

SUMMARY AND PRECISE DOCUMENTATION

Sympathetic ray : Refer to the work "**Radies thesis Physical Method**" : Food and Remedies, pages 217 to 219.

Personal wave : Each human individual emits a special radiation which is personal and characterizes him. Its wavelength, of the order of ten microns, varies with each person, thus making it necessary to find the setting for each one. This experimental adjustment is obtained, therefore by trial and error on this radiation, by presenting the pendulum above, either this person, or objects having been in contact with him for a certain time (clothing, bedding, pen holder , etc...) .

By carrying the pendulum thus adjusted above a letter written by a person, we find that this letter emits the same radiation, the paper having been impregnated with it for a certain time by the contact of his hands.

Because of its persistence, the personal wave can be used to search for missing persons, to identify objects that belonged to them, and to find, either by physical means or by the aid of mental orientation, objects that they lost.

The personal wave seems to be the result of the action on the human body of the terrestrial electric field. Its characteristics

teristics may not be absolutely constant and vary at different ages of the. life. Its polarity differs according to the sex of the person being examined. .

An impregnated object emits not only the personal wave of the person who usually uses it, but also all the other radiations of his radiation which may have encountered it : nervous wave, sympathetic wave, human wave, sexual wave, wave vital, disease waves, electricity, magnetism, radioactivity, eic. .. It is therefore only by a mental selection, in which the will intervenes, that one can. to discriminate a personal wave from the other radiations which accompany it.

Positive .at the man, the personal wave is negative at the woman.

For animals, the individual wave corresponds to our personal wave. »

• As for the plants, these also have a personal wave. This is why, if we experimentally regulate our pendulum on a plant, put _{our} finger in the antenna or in contact with a diseased part of the plant, the pendulum begins to oscillate.

The personal wave has a fundamental plane (Pf) and in this fundamental plane emits a specific axis that we will call the individual axis of the person. The point of ; departure from this axis appears to be a point at the top of the bone; iliac (hip protrusion bone). I

Is it the axis of Baradat to which astrologers attach! important ? We don't know yet; affirm it. . ;

For the rest of this study, we ask you to refer to pages 207 to 211 and 217 to 219 of the book “Dowsing Method Physics”. ;

Exercises

145. Sympathetic wave : Let's adopt one of the 1
ges indicated for this wave in the Settings Table and :
identification of bodies **and** radiation, pages 1 71. With the ;
chosen setting, the pendulum rotates positive (for a hand “

positive) above the back, or negative on the hollow. with the left hand.

On the other hand, thus regulated on the sympathetic wave, when one places the left index finger in front of or in contact with a healthy part of the human body, the pendulum will also turn positive. If, on the contrary, the left index finger is in contact with a diseased part. which therefore emits radiations for which the pendulum is not regulated. it will oscillate or stop. Let's do the experiment on an abrasion, a known lesion of the lung, a sprain. sore throat, conrusion, neuralgia, bronchitis. an effusion of synovium, etc. ...and we will see the oscillation or stoppage.

ⁱ 146. Let us experimentally set the pendulum on a remedy or a food. Its positive or negative reaction, on the back of our left hand (free hand which does not suspend the pendulum), will then tell us whether these elements suit us or not.

The concordance of the directions of gyration will fix us on the suitability and, on the contrary / on the contraindication, or harmfulness. In the event of a persistent oscillation, it is indifference that must be concluded.

147. Make the counter-proof by setting the pendulum on the sympathetic wave (positive) of the back of the left hand, and then present the pendulum above the food or the remedy to be examined. Note the direction of reaction. Positive gyration: good; negative gyration: bad; oscillation: indifferent.

148. Let us operate in the same way on the water, to know if it is drinkable (positive rif) or polluted (negative). First, let's experimentally set the pendulum on water deemed to be drinkable (positive). For polluted, colibacillary water, the pendulum will turn in the opposite, harmful direction.

149. Personal wave : Search on a writing page and •by mental selection for the characteristics of a person's personal wave. ■

150. Note that the written sheet of paper emits all the above radiations ..but with reversed polarity. (Refer to “ **Physical Method Dowsing** ”, pages 205 and 206).

IMAGES IN RADIESTHESIA _

LESSON TWENTY-FIFTH

THE VARIOUS CATEGORIES OF IMAGES

(Refer to “ **Physical Method Dowsing** ”,
pages 114 to 140)

SUMMARY AND PRECISE DOCUMENTATION

The image constitutes for the researcher a most formidable obstacle. Through the illusion it gives him of the presence of a real body, at a point from which the latter is sometimes very far, it misleads him and diverts him from the path on which **he** would find the crowning achievement of his efforts.

It follows that the search for the location of a hidden body usually presents itself in very arduous conditions. Each object, in fact, determines the formation of a large number of images, themselves mothers of other secondary images. It is therefore necessary to eliminate these multiple images in order to arrive at the real object that they mask.

By combining the various anti-image processes described in the following last lessons, it will be possible to methodically eliminate the images that will present themselves. But this will often be tackling a delicate and very long job. It is therefore preferable from all points of view to use the Schumfell Electro-Magnetic Developer, precision pendulum , **perfect anti-image**. designed primarily for the purpose .

to eliminate the images, whether by working in the field, or by teleradiesthesia, and at any time of the day or night. It is the ideal device for the practitioner.

However, theoreticians of dowsing will not fail to find in lessons 25, 26, 27, 28 and 29, on images, interest and even practical benefit. It's up to them. intention that we have established the three synoptic tables on the images, page 216 bis.

*
**

One could not better imagine the incalculable number of radiant foci which can be produced indirectly by the radiation proper to an object, than by comparing **it** with the considerable quantity of sound images which can be determined on receivers provided with the energy necessary **to** their training and adjusted **to** the proper wavelength, the voice of one person transformed and propagated by a powerful transmitter.

Just as the vibrations of these sound images are generally more powerful than that of the original voice, in the same way, more often than not, the radiation of certain images is more intense than that of the object which gave them birth, so that , fascinated by the emission of a powerful image very close to the radiant body that we are looking for. we sometimes pass **by** him without feeling him.

It is therefore of prime importance to be able to recognize images and to know how to eliminate them.

The existence of the images has been reported by the most expert dowsers and several of them have proposed several processes to detect them and protect themselves from their disconcerting action. To claim that we have never encountered any in our research is to 'ingenuously admit that we have never been able to distinguish their radiation from that of a real object and that, consequently, we have committed without to doubt many diagnostic errors...

It is to the research and experiments of Colonels Correnson and Camps pursued with tenacity during these last five years that we owe the acquisition of data. specifics on this fascinating and essential question. .

Picture species

"The images that one encounters in dowsing are of two very different kinds. They were, first of all, distinguished from one another by the denominations of " physical " images ;**q** and "dowsing" images ; but these qualifiers having been for many reasons recognized as unsuitable, it is advisable to substitute them **with** more adequate expressions. We shall therefore call direct images those which are formed with the very prolongation of the radiation of the radiant source, and indirect images those which are only connected **with** this radiation **and** whose whole radiant vigor is supplied by an external energy.

Direct images

>

Direct images are the result of a partial or total accidental deviation of the inherent radiation of an object or an emission source. They can arise **following** an accident absolutely identical to one of those which determine **the** formation of luminous images (reflection, refraction, dispersion, for example) and, consequently, obey the same laws. But they can also be the effect of a cause for which we find no analogy in optics. We will quote for this last case, the quadruple images of an object located close to the top of a dihedral angle (direct images of the phenomenon of the corners); the quadruple images of an object suspended by a **thread** or by a chain, etc., etc. .

Indirect images

These images are generated and maintained by the intervention of one **or** more **energies** external **to the** generating **body or to the radiant source**, and absolutely independent of them. These forces thus create new sources of emission possessing their **own energy** but attached **to** that of the generating source, the radiation of which is thus multiplied and, depending on the case, reinforced, reduced or modified.

It is these external forces: magnetism, electricity, light, which determine the emission of Electro Magnetic waves (EM) particular part which serve as a framework **for** the image and support for its specific radiation which they convey. When this support is multiple, it suffices to extinguish a single one, whatever it may be, of the radiations which compose it, to cause the image to vanish.

The different nature of the media generates different categories of images with different physical properties. It is for this reason that we have divided the **indirect images** (formerly called dowsing images) into two classes: the daytime **images** generated and maintained by the action of light (solar or **artificial**) and the **permanent images** generated and maintained by one of the three causes that we have cited (magnetism, electricity, light) and, in particular, by the earth's electromagnetic field.

Diurnal indirect images. — There are three categories of known daytime images: **magnetic images** (imago M) **with** MS support, **electro-radio-magnetic images** (**ERM images**) **with** triple support ES, MS, RS and **radioactive images** (**RA images**) **with** simple RS support.

Permanent indirect images. — There are three categories of permanently known images: **electro-magnetic images** (**EM images**) **with** dual support ES and MS, **electro-radioactive images** (**ERA images**) **with** dual support ES and RS **and ascending impregnation** (IM image) with triple support ES, MS, and <•>.

Families (1) of images. — Diurnal and permanent indirect images are never isolated, but united in family groups. The **remanence**, which is an RA image, and **impregnation**, which is an EM image, are linked to groups of families whose exact location can easily be determined **using** the pendulum. Each family consists of a **central mother image** (which may be M, ERM, EM or ERA, or IM), flanked by two equal groups of RA images, so that the number of individuals in a family is always odd and equal or serial objective figure of the radiation (Fig. 81).




Fig. 81

Le nombre d'individus que comprend chacun des deux groupes jumelés d'images R.A. qui flanquent l'image centrale dépend d'une propriété particulière de l'onde de la source radiante et est appelé **nombre caractéristique** de ceHe onde. Dans chaque groupe de familles, l'image centrale de la famille la plus proche de l'objet générateur, est directement liée à cet objet. Celle de la famille suivante est reliée à cette première image-mère centrale, ct ainsi de suite.

Si on détruit une image-mère centrale d'un groupe de familles, toutes los images-mères qui lui sont subordonnées disparaissent en même temps. Dans chaque famille c'est l'image-mère centrale qui donne naissance **aux** deux images radioactives qui lui sont contiguës; celles-ci engendrent les suivantes et ainsi de suite, suivant la forme d'un chandelier. Quand, dans une famille, on détruit une image radioactive, sa symétrie subsiste ; mais toutes celles qui sont subordonnées à l'image détruite disparaissent avec elle.

L'image-mère centrale, qui est toujours la plus brillante du groupé familial, peut être, suivant les cas, de même polarité que le corps générateur. ou de polarité inverse ; les images radioactives sont toujours d'une polarité inverse de celle de l'imago centrale.

(1) C'est **M. Charles ne',pon de Vaux** qui a observé le premier que **le nombre d:s individus d'une famille** d'images était toujours impair **et** que l'image centrale, beaucoup plus **intense** que ses compagnes, jouissait **de** propriétés physiques **très différentes**.

at sunset or when artificial light goes out. **It** follows that when the central imago is permanent, the family grouping is reduced, in the dark, to a simple populating of central images. If the central image is itself diurnal, the settlement vanishes completely. .

...O-⁰ 0... 0 .

Fig. 82

copper bigamist image

Polygamous pictures. — The same central image can belong **to** several family groups. In this case, she begets in each of the distinct groups the same number of radioactive daughters as she would have had if she had belonged to only one group (Fig. 82).

NOT
I
O).
: Radioactive images
0

• "I roage .bodyguard"
0 .. 0 • 0 -- + .. "' - • .. O 'O' 1
:
•
0
o .
1
s

Fig. 83

Radioactive footage *has* around a copper object

Radio-active images around an object. — An object generates around itself the same number of radioactive images as a central imago-mother from **which** it would have given birth. An object producing groups of radioactive images in each of the four branches of **the** magnetic cross can, **from** this point of view, be likened **to** a polygamous image (Fig. 83).

The four **electro-radioactive images** which are called **>:C bodyguards of the object** and which are much smaller than it, are intercalated between this object and the first radioactive image **to** which it gives rise.

An illuminated object also provides six groups of radioactive images; **one** in each of the six planes of its photogenic radiation.

Image radiation

The images emit, either with the same polarity, or with an inverted polarity, the specific radiation of the generating object, which gives the illusion of its presence; but this radiation is conveyed by different supports. Three categories of images, **M** images , **EM** images , **RA**

images. radiate exclusively towards the sky and have no vertical or oblique rays **directed** towards **the** ground. On the contrary, the images of the **ERM** and **ERA categories** radiate like the generative body in both directions. ■
the six sketches in FIG. 84 compares ;
the radiance of an aluminum box with that of an image of !
each of the five different categories **to** which olle can give j
birth. □

Various categories of indirect rmagcs i

Indirect image categories are distinguished by the nature of <
supports that carry the specific rays of the image, supports >
shown in **figure n° 84** on page 189. (

1 r* Class : DAYTIME IMAGES j


There are three currently known categories of daytime images: the ! magnetic images (**M**).
electro-radio-magnetic (ERM) images , '
radioactive images (RA). i

Magnetic pictures. — Los magnetic images (imagos **M**) li]
simple support MS are' produced by the action of light on the ?
radiate from the magnetic charge proper to the generating body. For ,
demonstrate this, it suffices to artificially extinguish co radiation **ot** j
we note the immediate fading of all the images M that **it**
has formed. 5

The families of images **with a central mother M** meet principally
ment in each of the branches of the Magnetic Cross of an object -
and, in ascending and descending rows on the vertical do each *
object. ;

Nr

Electro-radio-magnetic images. — These images (ERM images) with triple support. ES, MS, RS are produced by the action of light on the radiation behind the electrical and magnetic fields proper to the



body as well as its radioactive radiation. If we extinguish just one of these three rays,

the ERM image vanishes.

Groups of ERM central mother image families **occur** in each of the six radiant fields of an illuminated object (radiation from the photogenic double triangle or from the solar plane).

In the two cases of images which precede, we noted, the intervention of the light in their formation, but the research made to determine with precision the mechanism of this action did not succeed yet.

Radio-active images : These images (RA images) on a single support RS, are generated and maintained by the **magnetic action** of light on the radioactive radiation specific to the generating body (or an image of one of the four other categories). If, in fact, we extinguish this radiation by artificial means or if we illuminate the body with **blue** light, the image vanishes.

These images which, during the day, are in considerable number, meet exclusively as immediate satellites of a body or of an image of one of the four other categories (images M, ERM, EM, ERA), mother images of which they are the girls. They present themselves on either side of this object or this parent image in a double group of a strictly constant number of units for the same substance or the same radiation. **This number is referred to as the characteristic number of the substance or of the**

, radiation. For instance, two groups of two RA images on either side of a copper imago, two groups of 5 RA IMAGES, on either side of a gold image, of 6 images on either side of a mother image of lead, substances whose characteristic numbers are respectively 2, 5, 6.

Afterglow is a radioactive image.

Common general properties of daytime images : Daytime images are the only ones. **0** like objects, produce complete magnetic crosses around them (with their 4 radiant fields and their four groups of families of images). They are also the only ones which give rise to **0** ascending and descending vertical columns of families of images.

They appear to be insensitive to antenna ray, visual ray and montai ray.

2nd Classe: PERMANENT IMAGES

There are two major classes of known permanent images. them **electromagnetic** images (**EM images**) and **electro-radio images active** (**ERA**

images).]

Electro-magnetic images : These images with double support ES and MS are generated and maintained by the action of an electric field **electric or magnetic** on the specific radiation of a body. If we **at 4** extinguishes the radiation proper to the electric or magnetic charge of a body, the image continues **to subsist** . It is therefore not these charges which come into play in the formation of the EM image, **but** an energy of the same external nature or generating body.

One finds families of imoges **with** a central mother EM near a metal bridge, a motor car, a stair railing, a sewing machine , and, in general, any magnetic mass or of any electrically charged body.

Earth's electromagnetic field can also produce EM images. Such is the cos of imoge-impregnation.

Electro-radio-active images : These images (**E^AA images**) with **dual** ES and RS

support. are generated and maintained by the action of the earth 's **EM field** on the radiation of the own electric charge or the radioactive radiation of a body. To demonstrate this, it suffices to extinguish one or the other of these two rays and one immediately sees the image disappear. **These images** are small .

ERA images are found at the four magnetic cardinal points of each object and very close to that object . In each of the four bronchi of the field of the Magnetic Cross this first ERA image is topped vertically by a double ascending and descending column of families of images **with** a central ERA mother. The polarity of these central images is none other than that of the corresponding bronchus of the Cross and their intensity is all the higher as they are vertically further away from the horizontal plane on which the object rests.

ERA images are the only images that persist in the Magnetic Cross fields of an object that has ceased to be illuminated.

If the ERA images are generated by the terrestrial EM field, it must be possible to produce them artificially by subjecting an object **to** the action of any EM field. This is indeed what is happening, in fact.

Beside a piece of silver P, place, as shown in figure **85**, a bar magnet. Above the negative part of this magnet, place a capacitor **(1)** .

We thus realize an EM chomp. From then on, it will be easy for us to note that, in the extension of the magnet, and very close to the piece, an ERA image has formed, the starting point of ascending and descending columns of families of mother images. central to this category.

If we place the object above the condenser, not one, but four vertical rows of images are formed; two gifts lo direction

(The capacitor used for this experiment is a small object that can be found at radio equipment stores . two blades of ct varnished sheets separated by tm celluloid or cardboard electrical insulation. THE three pieces are joined by shoe eyelets.





Fig. 85

of the magnet and two does the perpendicular direction. This experiment demonstrates that it is indeed the terrestrial electromagnetic field that is due to the formation of the "permanent bodyguards" of an object.



@ Irradiation

Fig. 86

Two particularly interesting types of images:

Remanence and Impregnation -

Remanence (1) : When a body has stayed a certain time in a given medium, it impresses the ether of this medium, and, after it has been removed, he leaves there a temporary imago which gives the illusion of his presence. This is the phenomenon of **remanence**.

If we move the table on which it was formed, the remanent imago will not follow the piece of furniture, but it will subsist at the points of space that the object had momentarily occupied.

(li) It was Colonel Camps who was the first to recognize the absolute identity of the afterglow with a radioactive image. It is also to him that we owe the discovery of the EM

The afterglow, generated and maintained by light, is a temporary radioactive image of an inverse polarity of cello of the generating object and of the same form as this object.

It vanishes momentarily if the object ceases to be illuminated, or if the object continues **to** be illuminated, it is placed itself in shadow or penumbra.

No afterglow occurs in a room lit **with** blue or violet light.

The duration of the radiation of an afterglow is exactly equal **to** that of the presence of the object at the points of space which it has impressed. If it is momentarily extinguished in darkness, the time of its eclipse does not count in that of its total radiation, the duration of which is always equal to the time that the object spent at the points where it was formed.

The remanence is not an isolated image, it belongs **to** a group of families **with a** central mother EM whose location is very easy to recognize.

Impregnation : The phenomenon of fluidic impregnation, which some mistakenly confuse **with** that of remanence, is absolutely distinct from it.

If we place a piece of silver on a scrap of paper laid out on a table, and if, after having traced or penciled the limits of the location which it occupies **there**, we remove the piece and withdraw the sheet of paper, we note that the surface of the paper on which the piece rested continues **to** emit the radiations of silver, as, moreover, those of other alloy metals.

We will have achieved a fluidic impregnation which is nothing other than an electromagnetic image of the same form as the impregnating body, but of inverse polarity.

This image is not, moreover, isolated, and it is very easy **to** find the family groups to which it is attached and which move at the same time as the search of paper.

Generated by the earth's electromagnetic field, the imago impregnation is permanent on certain substances such as paper or certain fabrics, as long as it has not been destroyed by a western cause or by suitable artificial means. On other substances, on the contrary, dry wood, stones, leather, metal, the **click** is absolutely ephemeral and goes out in a few seconds.

Mental impregnation : If, after having traced a closed contour on a sheet of paper and after having thought very strongly (of the copper powder for example, we suppose that the surface delimited by the **co** - trace is filled with this powder, we project **it inside the outline of the montai - supported** copper spokes .

The paper becomes impregnated with these radiations and ends up in the Omottro in a permanent way. This created a mental impregnation of copper on paper.

Mental impregnation is a permanent image **with** triple support

ES, MS and (•) **of the same polarity as the generative substance** : the extinction of only one of its three supports causes it to vanish. It is engendered and maintained by the action of the terrestrial EM field **on** the specific radiation proper to the impregnating substance and on its A - supporting CO radiation which accompanies the mental ray. It emits this last ray as can be seen with the help of a pendulum adjusted to this radiation.

It is a lifeless image which, temporarily vanished by the action of a point, instantly revives as soon as this action **has** ceased.

The phenomenon of mental impregnation sheds light on a fact which for a long time seemed inexplicable and which some have even described as absolutely fanciful: the radiance proper to a painted picture, and, in particular, that of a portrait.

It also allows us to understand how it is possible for an experienced rad;esthetist to distinguish an original painting from a copy. mental impregnation. not comprising second degree impregnation.

Mental impregnation is a quadrigamal image, playing the role of an object and **which does not reproduce itself**, it gives birth. in two perpendicular directions. with **four families of images with central mother EM**

These two directions are magnetic North-South and East-West. They remain invariable if the paper is moved.

DISPOSAL OF IMAGES

Some operators affirm that, just as it is possible for them to subtract their sense of smell **or** the perception of certain perfumes, likewise, they have the faculty, by the sole action of their will, to render themselves insensitive **to** the perception of an image. There is an application of the wave selection operation discussed on page 82. The specific radiation of an image differing from that of the generating object, not only by the nature of its supports, but still by that of the accessory radiations which accompany it, one conceives that it is possible **with** a very delicate perception to discern it. But we think that, except for particularly well gifted subjects, the process is not **to be** recommended, especially for beginners. We found, moreover, these results to be lacking in operators who claimed to be able to rely on them entirely.

It seems infinitely safer to resort **to** a physical process.

We will not recall here the processes of extinction enumerated on pages I 18 128 of the work "**Dowsing Method Physics**", and of which one will find a summary in the synoptic tables of pages 216 bis and following.

We will content ourselves with concluding that a first solution to the problem **was** given by M. Discry, in his work "**La science des dowsers c\l l a range de tous**". The form wave of the disc extinguishing certain images, the black color making others vanish, 'we will **theoretically make ourselves** insensitive **to** the perception of an image by holding in the positive hand a pendulum completed with a witness and by seizing the free negative hand, between the thumb **and** the index, the other fingers elongated in antenna, a disc of black color.

We said **theoretically**, because in practice there is a very big pitfall **to** avoid. It may be that when we present the pendulum over the former location of an erased image, we inadvertently think **it** the object which gave rise to it. The instrument will then begin **to** rotate no longer on the radiation of an image which no longer exists, but on the specific radiation of the object itself which the mental ray has brought back ^{to us}. It will therefore be important, when we proceed **with** our final control operations, to banish distraction and adopt an **entirely passive attitude**.

Another process, recommended by M. Chouteau, in his treatise on dowsing, consists **in** holding one's breath during the gyration of the regulated pendulum. If the pendulum rotates on a real body, its gyration will decrease in intensity, but will persist all the same: if it rotates on an indirect "image" of any category, its gyration **will** transform into an oscillation.

To put all the possible chances on our side, co-control will have to be done with Electro-rodio-magnetiquo Schumfell Revealer, anti- image. provided with the appropriate witness, against the suspensor stick.

TWENTY-SIXTH LESSON

DAYTIME IMAGES

DOCUMENTATION AND EXERCISES

1. — Magnetic images

1° Let us place at one of the corners of a rectangular table a small aluminum box; and, after setting a pendulum on the aluminum, let us bring it to the symmetrical point of the box with respect to the center of the table. The pendulum will start rotating in the positive direction. This is because an image of the aluminum box and an image of the same polarity have formed at this point. Between the image and the pendulum, let us interpose an object of a height greater than that of the box; the pendulum will stop. This proves to us that the specific rays of .llu-




Fig. 87

RA-

Images of an object or edge of a rectangular table

minium emitted by the image are it support MS. Let us remove the iron blade, successively present above the image a regulated pendulum on MS rays, then a pendulum. set on ES rays, finally a pendulum set on RS rays. We will find that, alone, the first pendulum will start spinning. It is therefore an image with an exclusively MS support, therefore **a magnetic image**.

2° Let's repeat the same experiment in the evening, in a lighted

room:

we will get the same reactions. Turn off the light **(1)** ; the swing of the pendulum will stop. The magnetic image is therefore generated by the action of light: it is, consequently, a **diurnal image**.

3. We have said, page 188, that the magnetic image is produced by the action of light on the radiation of the **magnetic charge proper to the body**. This radiation is none other than the MS ray whose serial objective number is 5. Let us place **0** side and very close to the box five small square cards of yellow or red color superimposed, we will extinguish the MS rays which it emits. Let us **refer to** the location of our magnetic image. We will find that she has fainted.

4° **Images M on the vertical of the bodies**. Let us note above and below each illuminated object, the existence on its vertical of a double ascending and descending column of families of images **with** a central mother **M**. Let us note, by counting them with




Fig. 88

Images sur la
verticale d'un
objet en argent.




Fig. 89 Pénétration avec
l'index des images RA et M

the index finger of the hand negative than the number of these families ost, in each column,

(1) It was **Mr.** Ch. De'pon de Vaux who was the first to point out the existence **on the** vertical of each illuminated object of a double column (ascending **and** descending) of twigs of images.

equal or **objective number of series** (or **specific number of the substance**) ie 3 for aluminium. (Fig. 88).

Lo **point**, destroying the images **RA** and the image M being insensitive or ray of antenna, we will be, if we want to feel them one and the other. in the obligation to **penetrate them with the index**. (Fig. 89).

distinguish, by holding a coin in the negative hand (squeeze it well) the M images which resist or radiation from the disc, from the **RA images** which this radiation extinguishes. Note that the images of the ascending column are straight and those of **the** descending column are reversed (base up, top down).

! o illord maynetique




Fig. 90

**The four fields of the Magnetic Cross
of an aluminum object
and, in each branch, a group of 3 families of images**

5° **Images of the four fields of h Magnetic cross.** Let us adjust our pendulum on the aluminum and note the presence, around our box, of each of the four fields of the Magnetic Cross, in which propagate rays of aluminum **with** MS support and of limited length (unequal in each of the four branches). Let us note that the polarity of the MS rays varies according to the orientation of the fields and endeavor to discover the law which governs this phenomenon.

To distinguish the specific radiation of the chomps from that of the imoges that form there, we are obliged to perform an operation.

of mental selection with which we will very quickly become familiar and this selection will be all the easier since the emission of an image differs on many points from that of a real object. (See table 2 and figure 84 pages 216 ter and 189).

The objective series number of aluminum being 3, let us note that there is formed in each branch of the cross a group of three families of images of aluminum ($4 \times 3' = 36$) à central mother M to which are added the four radioactive images generated directly by the object, ie 40 in all (if we neglect the four columns of electro-radioactive images which are inserted between the box and the first RA image which it generates in each branch).

6° Radiation of images M. — Let us observe, by placing the index finger of the negative hand 'below the image. that it does not radiate towards the ground either vertically or in divergent directions.

He. — Electro-radio-magnetic images

1° Let us seek in the direction of the shadow cast by an object or on its extension, one of the ERM images determined by its beam of illumination and mark it. It will be easy for us to




Fig. 91

Image ERM dans l'ombre portée d'un objet

que nous interposerons à hauteur
sur l'aluminium et l'imago soit

distinguish it from one of its radioactive daughters by the fact that we will determine its fainting by squeezing an electrical insulator (dry sheet of paper, for example) in our negative hand. Let us note the existence of the triple support of its specific radiation - ES, MS, RS, first with the aid of a pendulum successively regulated on each of these three radiations, then by the fact that each suitable time between a pendulum regulated an electrical insulator (slide of glass or sheet of paper) or a sheet of sheet metal, or a cloth or a blue cardboard, the gyration of the pendulum will stop.

2° Mode of formation. — Under the same conditions as for the image M, let us note by the affixing against the box of either five or seven superimposed cardboard boxes and the extinction which it causes, that the formation of the **ERM image** is properly due to the the action of light on the simultaneous radiation of electric and magnetic charges and of the object's own radioactivity.

3° Artificial production of an ERM image — Let us place beside our aluminum box, as indicated in the figure opposite, a right magnet and 0 side of it, a pocket battery, the

pôles de même nom jurlaposés, nous constaterons, si ces pôles sont de positifs, la formation d'un groupe à trois familles d'images d'aluminium mère contrôle ERM+.

Pocket right stack magnet




Fig. 92

We will distinguish the ERM image of one of its radioactive daughters by seizing a black disc with the negative hand, which will determine its fainting.





Fig. 93

We will observe the same fact if we replace the previous device with a simple **well-stoppered bottle**, the air of which, having become radio- active, will determine a double field El— and Mag—; but then the ERM image will be negative. If we turn off the light in our laboratory, we will find that the ERM image disappears, which proves that light is essential **for** its formation.

4° **Images of the sixfold photogenic field or images of the fields of the double photogenic triangle** : The beam of light (Continued on page 203).

The six radiant fields of Photogenic Radiation from an Aluminum Disc with their six groups of image families.



Direction of the
light source

-9-

Fig . _ 94

The hatched outlines represent the central images ERM— of the aluminum disc; the video contours represent the RA+ images of the same family.

Since the specific radiation of aluminum is positive, the images RA generated directly by the disk are negative.

When the light source is very close to the illuminated object, which is the case in a laboratory lit by an electric lamp fixed to the ceiling, the soft triangles formed by the axes of the six radiant fields are neither isosceles nor similar.

which illuminates an object determines around it the formation of a sixfold field in which the specific rays of the object of limited length are exported in the direction of the beam of illumination by a double support ES and MS and in the four other directions by a triple support ES, MS[^] and RS. (Fig. 94).

Under the same conditions as for the Magnetic Cross, we will see:

1° The different polarities of these fields.

2° The formation in each of them of a group of families of images **with** a central mother ERM and a polarity conforming **to** that of the corresponding field.

5° **Images by reflection** : When the beam of illumination from an object encounters a flat reflecting surface, it first determines

a direct quadrigamc image of this object **at** its point of incidence on this surface, then four do groups. families **with** a central parent image ERM of an inverse polarity of ccllo of the object in the direction of the reflected beam **ct** in that of its extension, ot, on the contrary; of the same polarity in each of the two directions perpendicular to this **direction** .

Figure 95 represents the case of an aluminum disk (positive body) producing by direct reflection on the mirror a positive diroclo image accompanied by its 4 negalivos RA images and 36 indirect images in the horizon plane taI of the disk. Each of these 40 images produces a Magnetic Cross with sos 40 **images and an ascending** and descending double vertical column with sos 20 images.

III. — Radioactive images

1° Let us refer **to** the experience of page 196. Let us place our aluminum box **at** one back corner of our rectangular table. Diagonally. of the object we shall find, as has been said, an imago M of the same polarity as this object; but they two other anqlos 'we will observe the formation of an image **RA** of opposite polarity, which we will be able, thus, to study _{SMS} fear of confusion avocado. first.

Let us first note that olle is **0** support RS. If we place a **lead** object near its center, or if we seize with the negative hand, either the same object, or a fabric of blue color, the gyration of our pendulum will stop. This support is unique, but if we present above the image a pendulum set on any other known support, it will experience no reaction.

It will be a memo if between him and the image we interpose a screen formed either of a lomo of lead or of a cardboard of blue color.

2° Let us note that, if our laboratory is illuminated with blue light, we indeed find, diagonally of the object, an image **M**: but that at the two other angles, the images RA of the preceding experiment l') are not formed. **It** is the same with violet, indigo, dark green lighting, that is to say with a light of a color **with** electrical accompaniment. ,

3° Let us note that the **number of individuals** of a family of images is indeed equal **to the** objective number of the series, or, better, to the particular **number** of the substance which gave rise to it, that is to say to **3** for silver, iron, aluminium, etc..., 5 for copper, 1 for gold, 13 for **lead** , etc..., etc... tableJ an object of each of these different substances and make the verification on the second family of images

of its Magnetic Cross of which we will first of all seek the central mother-image.

EM wave a particular element whose nature still escapes us and which intervenes in the determination of the number of individuals of which a family of substances is composed. images generated by one of its emission sources.

Let us observe that the **diurnal images** of all categories emit calorific rays CL and chemical rays CH, by noting the gyration of a pendulum, regulated on these radiations, at the moment when **one is** presented above these images.

...

TWENTY-SEVENTH LESSON

PERMANENT IMAGES

MATERIALS AND EXERCISES

1. — Electro-radio-magnetic (EM) images

1 ° Artificial production. — If we place in front of an object, in silver, for example, one of the sides of a pocket battery or **the** pole of a magnet, we will obtain in the extension of their direction a group of families of images of silver with central mother **EM** of a polarity corresponding **6** to that of the field which has impregnated the object.

RA- RA- Ert+ RA- **O** *go [IJ Fl;* *RA- E|1 +*
RA-

Y y Silver object

Pictures —

0 0-0

Pocket Pile

F i g . 96

Let us note by analogous processes **6** those previously described that the central images of these families are with ES and MS supports. To distinguish them from their radioactive daughters, it will be enough for us to shake in the negative hand a small object of lead which will make us insensitive to the perception of the **RA images**.

Let us note that the **EM image** is **permanent and** that light does not intervene in any way in its formation, by observing that this imago will always continue to **subsist** after we have turned off the lamp which illuminates our laboratory.

2 ° Elimination. — Note that this image is vclctilized by **a** point directed against it and placed at a suitable distance, that it is also extinguished by a very close disc as well as by an electric or magnetic field of the same polarity, whereas the field of opposite polarity is its action. Let us also note that we can make ourselves insensitive to its perception by seizing in the negative hand either an electrical insulator (glass or dry paper, ebonite object, etc.) or an iron blade, or a coin (disk).

• **3 Screens.** — Note that the radiation of this image is stopped by an electrical insulator (glass or sheet of paper) by a sheet of sheet metal: finally, that the EM image • does not radiate towards the ground.

He. — Electro-radioactive images (ERA images)

1° Description. — Let us note with the aid of the antenna tip, **ot** in the immediate vicinity of an object placed on a table, that there is very close to this object, and in each of the four directions of its magnetic cardinal points, an image of the same shape, but much smaller than him. Note that from each of these four images start two columns. one ascending, the other descending from families of images **with** a central mother ERA. Note that the images of the descending columns are reversed (base in the air, top in bos). Note that these images are intercalated between the object and the first radioactive image that it forms in each of the four branches of the Magnetic Cross. Finally, let us observe that the ERA images are permanent by noting that they will continue to subsist after we have extinguished the lamp of our laboratory and that they are, therefore, the only images of the Magnetic Cross which subsist in the darkness.

2° Radiation : Note that, like ERM images, ERA images radiate either vertically or obliquely both in the direction of the sky and in that of the ground.

3 ° Supports-Screens : To note by the processes previously described that the images ERA are with double support ES **ot** RS, that Day radiation is stopped, either by an electrical insulator, or by a sheet of lead, finally that one makes oneself insensitive their perception **by** grasping with the negative hand: either an electrical insulator, **or** a lead object, or a blue object.

4° Artificial production : Artificially producing ERA images par. the process described on page 192. (Fig. 85 and **S6**).

TWENTY-EIGHTH LESSON

REMANENCE AND IMPREGNATION

DOCUMENTATION AND EXERCISES

1. — Afterglow

1 ° **Formation** : At 10 o'clock, place an aluminum box on a table. At 10 a.m. 15, let us withdraw it and present to the site which this box had occupied, a pendulum regulated on aluminum. We will observe the gyration of this pendulum. Let's remove the table; it is possible that we find at the point where we deposited the box, an image-impregnation on wood which, quite **ephemeral**, will disappear very quickly. Let us return to the location which the box had occupied in space; we will see a gyration of the pendulum which will definitely stop at 10 o'clock. 30. We will have made a remanence **of a quarter of an hour**.

2 ° **The afterglow is a daytime image.** — At the moment when our pendulum turns on an afterglow, let us ask **an** assistant to close the switch of the illuminating lamp, we will note that **the** gyration has stopped. Ask him, then, to give us the light, and we will see the gyration resume almost immediately.

3 ° **The afterglow is a radioactive image (1).** — To realize this, it suffices to observe that the afterglow is volatilized by the action of a point placed **at a** suitable distance, that its radiation is stopped by lead or **by a** blue cloth or cardboard and **quo**, to make us insensitive **to** its perception, it suffices for us, the regulated pendulum being held by the positive hand, to seize, with the other hand, an object of lead or a blue colored object; finally that, in a laboratory lit **with** blue light, **the** phenomenon of remanence does not occur.

•

4° **Of remanence : the duration of an afterglow is limited, but governed by a law .** On the table of our laboratory, illuminated with electricity, let us place, once again, our aluminum box in **a determined point**

(1) This discovery is due **to** Colonel Camps.

where we will leave 1a from 8 p.m. at 8 p.m. 15. At 8 p.m. 15, let us withdraw it and observe the remanence that its former presence has determined. At 8 p.m. 20, let's turn off the light in our lab. At 8 p.m. 25, let's open the switch: we will see that the afterglow will only disappear at 8 p.m. 35. Interrupted for five minutes, which will not have counted in the time of its existence, it will have lasted exactly a quarter of an hour. We can compare the phenomenon of remanence to the unwinding of a reel which would occur in a time strictly exact to that of its winding, but which would stop momentarily in the dark.

5° **Afterglow is not an isolated image.** L It is part of a family attached to a set of four groups of families with a central mother EM and aligned along magnetic North-South and East-West directions passing through the former location of the object. It is, in short, a transformation of the old Magnetic Cross of the disappeared object.

11. — Fluidic impregnation

Do not confuse **physical impregnation** with **fluidic impregnation**. If we soak a piece of absorbent cotton or cardboard with saliva, urine or blood, this cotton and this cardboard will emit, in addition to their own radiation, that of the substance to which




Fig. 97

**Objet posé sur une
feuille de papier
horizontale**




Fig. 98

**Imprégnation
sur le papier
après enlèvement
de l'objet**




Fig. 99

**Position
de l'imprégnation
quand on incline
le papier**

they will have been asked to serve as a support, and there will be no difference between the radiation of this substance and that which it would have emitted if it had been placed in a dish. We will have carried out a **physical impregnation** whose radiation is that of a **real body** and which, consequently, does not present any particularity from the dowsing point of view.

1 ° **Production of the fluidic impregnation.** Let's place a sheet of paper on a table: in the middle, let's place our cylindrical box

of aluminum o circular section and trace in pencil the circumference that limits it. Let's leave it for a while, then take it out. As long as we do not move the sheet of paper, we will observe **inside** the circumference an image remanence RA— whose duration will be equal to or time of the stay of the box at its omplacement.

But, if we move the search of paper, we will find that the traced circle continues to impress a pendulum set on the aluminum and that there has been formed, **inside** the circle, an image which we will call image-impregnation, which seems firmly attached to the paper and which follows it wherever it goes.

2° The formation of the fluidic impregnation is almost instantaneous. —• We left our box in place for a certain time in order to better grasp the transition from the phenomenon of remanence **to** that of **impregnation** : but, to better understand this second phenomenon, let us draw on a sheet of paper a circumference of a diameter a little greater **than** that of the box. Let us pose the box for a quarter of a second inside this circumference and we will see that the circle radiates the aluminium. **The formation of the 'image-impregnation was instantaneous.**

3° The fluidic impregnation is an EM image whose polarity (first degree) is **inverse** to that of the generating body. Let us first note in our experience the formation of a negative image, by observing (or need with the oido of a polarity indicator) that, presented or above this circle and held by a positive hand, a pendulum set on aluminum rotates counter-clockwise. Let us then note that this image is an **EM** imago by observing that, if we direct a point towards this image, the pendulum stops, that it is the same if we place the circle in the negative field of a magnet, that it is. even if we approach the image a coin, finally that its radiation, stopped, either by an iron blade (sheet metal) or by an electrical insulator (glass blade or cotton sheet). is absolutely insensible to the action of lead. From the above, we will conclude that the search for remedies must be based on a **physical** impregnation : urine, blood, joists, hair, used handkerchief ^(I) **and** not on a first-degree fluidic impregnation , **which would** give polarity contraindications, but it can be done on a second-degree **fluidic impregnation** using the writing of the molado for example , or on a Chanoy imprint.

Let us note, about the polarity of the impregnation either of the first degree, or of the second degree, that an impregnation of the first degree on paper emits the radiation of a radiant source with a polarity

" • **The handkerchief which contains the patient 's mucositis and which has been frequently in contact with his nose contains both a physical impregnation and a physical impregnation; but because of the perpetual chills of which it is ' / object, this dcrmerc is very quickly .race - , Fluidic impregnation is , moreover . very little tenacious on the canvas .**

reversed. Under these conditions, we have the greatest interest in **transforming it into a second degree** impregnation whose emission is of direct polarity, which is easy by the simple affixing of a search of white paper, as new as possible, on primitive impregnation, that is to say of the first degree.

This is what we can do, especially when we want to study the radiation of a person 0 with the help of a **page of writing**, this constituting the impregnation of the first degree. Having affixed to this page of writing, or letter. another sheet of white paper, as white as possible, one no longer has to worry about inverting the results, and that is infinitely more convenient. •




Fig. 100

Image-impregnation of a writing page

Its geometric shape is always a little fuzzy, from 0.04 to 0.05 centimeters in height. This impregnation persists at night. Pages written for a century continue to emit the personal wave of their author.

4° **Let us note that this image is permanent**, by observing that, when it is hung on a suitable support and that it is not destroyed by artificial means (grinding, energetic friction, prolonged action of a point, a Electric or Magnetic field of the same polarity). this image which persists day and night is permanent and that its permanence is its known limit. We also note that, if this permanence is observed with paper **and** certain clothing fabrics, it is not the same for all supports. Let us note that on dry wood (not painted or varnished), on leather, **on** marble, **on** stone, on metal, on glass , **the image- impregnation is absolutely ephemeral, its detachment** occurring at the after only a few seconds. Let us note (by making the test with a pendulum adjusted, successively on each of them) that a **page of writing** constitutes an image-impregnation of **all the radiations** of the person who wrote it and in particular of his personal wave . If we have the possibility, let us note the permanence of the image-impregnation by examining manuscripts on paper dating back several hundred years. (Fig. 100).




Fig. 101



Figs. 102



Figs. 103

Note that **if** we make a dumpling of a page of writing, **it** will be nice to straighten it **with** the press. the impregnation-image which will have disappeared forever will not be made to reappear.

Note that after the burning of the paper, the ashes of a writing bag continue to radiate sound around them as long as they remain on an agglomerated block. Let's spray the block, we will see the radiation extinguish.

5° Liquid supports : Liquids can serve as substance-support **for** an imago-impregnation. Let us note that, if after having left it to remain for a few seconds in a glass of water, we withdraw a silver spoon, the water will continue to be 6 **mottro** los radiations of silver. Let us note by the processes recalled or second paragraph that it is indeed a question of an imago EM—of the spoon of orgon . Let us note that the image is quite persistent and that it is necessary to change the water or the glass several times before seeing it available. Let us note quo, for that there remains at the bottom of the glass a thin blade of liquid, the image is there. grip, whereas it would not hold on dry glass.

Let us note that, as a result of the innumerable diurnal images formed by an object of gold or lead, the sap of numerous essences of trees is impregnated with these radiations up to a considerable distance. Establish by the processes indicated and in particular by the tip that it is indeed an image-impregnation EM-. Note the same fact for the liquid of the pits of aisanco.

If the sap of trees is impregnated, it is the same with the blood of animals. While seated, let us place an aluminum box on one side of our thighs, we will notice after a short time that a pendulum set on the aluminum turns above our hand. From this experience we can deduce important consequences for metal therapy and also for the precautions that must be taken by people who deal with medical dowsing, even by the mental method. In order not to become one day victims of their devotion, these people have an interest, after each work session, to **disimpregnate** themselves, as will be said further on, with the **aid**

of an electric or magnetic field.

6° The impregnation image is not an isolated image. — It is part of a set of four groups of families oriented respectively along the magnetic North-South and East-West directions . It is, in short, the magnetic Cross of the image-impregnation which moves with the paper, but pivots around it. of this . image which serves as its axis of rotation, so as to remain invariably in its normal direction. But, created by the terrestrial EM field, .the mother-images which are formed there are permanent EM images, instead of being, as in the magnetic cross of an object, diurnal M-images.

7° Impregnations of images : If, at the site of an image, we carry a sheet of white paper, we observe that an image-impregnation of this image has formed at the impressed point. The fluidic impregnation of an image, even diurnal, is a permanent **EM image** of reverse polarity, except for ERA images.

The impregnation of an M+ image is an EM image—

Imprinting an ERM+ image is an EM image—

The permeation of an RA— image is an EM+ image

The impregnation of an ERA+ image is an EM+ image

The impregnation of an EM + image is an EM imago—

8° Impregnation in the second degree : If the impregnation of an image is an EM image, it must be the same for the impregnation of an image-impregnation. There must therefore be second-degree impregnations. This is indeed how things are going. If, above the circle, which we have traced to materialize our first degree impregnation of the aluminum box, we place a piece of white paper, we observe that, **d s *** imprints there a new image identical to the previous one. , of the same form, of the same intensity, but. of reverse polarity. This will be the second degree impregnation-image of the box, image II'. In the same way, we will obtain images-impregnations of higher degree He', He..., etc..., all of the same intensity, the terrestrial EM field taking care of maintaining the centers that we will have thus lit; the images of even degree being of the same polarity as the body, the images of odd degree of opposite polarity.

9° Impregnation of the Pendulum : A pendulum which has received for some time the specific radiation of a substance, emits it **0** its turn and rotates when it is present on any body. This is a fact of temporary self-adjustment by impregnation which can cause us to make many errors if we do not take care to de-impregnate the pendulum each time we change substance. The impregnation on a spherical surface being unstable, a slight shock on the table or a simple shaking will suffice to restore the instrument to all its initial qualities.

10° Mechanism of the transition from remanence to fluidic impregnation. —

Let us refer to the experiment of **1 ° on page 208**. On a sheet of white paper placed on our table, let us carry our aluminum box. Draw with a pencil the circumference which we limit the contour and leave, without touching anything, the device we place for two minutes. Then, let us remove the object, and without moving the place let us place above the circle the pendulum regulated on the aluminium. Held by a positive hand, it will begin to turn in the negative direction on the imago RA— which constitutes the afterglow. After two minutes, we will see the gyration stop. Then, after a few seconds of swinging, the pendulum will return to **0** to turn in the same direction, never to stop. This is because the RA— image has been displaced **and** has been replaced by

an EM— image, the central image of a family of polarity different from that to which the afterglow belongs. There is a displacement of the Magnetic Cross, since we know that two branches of this cross, located on the same alignment, are of different polarity. A work with blue light would make it possible to find the rigorously exact modalities of this phenomenon. Without wishing to fix them, let us content ourselves, for the moment, with noting that the phenomenon of the devidago had the effect of determining a momentary displacement of this cross, a displacement which ended with the arrest of the devidago.

1 1 ° Ch:trcy imprint : A device indicated by M. Chanoy, ingeniouf "des Mines, makes it possible to obtain directly on paper back fluidic impregnations of the same polarity as cello of the impregnating substance, practically equivalent, consequently, with back impregnations to the socond degree.

Sweet lamos of glass constituted, by oxomplø, by old clichés. The **9 X 12** photographic prints are separated by a very **fine cloth** (a muslin handkerchief folded into eight) under which the excavation of the paper to be fluidically impregnated is placed: the four elements are then tightened by rubber bands which keep them firmly fixed.

The device is placed on. the jet of the radiation to be studied will pass in such a way that it does not encounter the paper until after it has passed through **the handkerchief**, which, as can be seen by the following experiment, **has g i t give this case as a polarity reverser.**

Let us hold, indeed, in the positive hand, a pendulum adjusted to silver, in the dish of the other hand let us place a handkerchief folded into eight on which will be deposited a silver object. We will find that instead of rotating in the positive direction, the instrument will begin to rotate in the negative direction. It is therefore radiation of inverted polarity which impresses **the** paper.

TWENTY-NINTH LESSON

MENTAL IMPREGNATION

(IM picture) •

DOCUMENTATION AND' EXERCISES

. 1 ° **Produce a mental impregnation** : On a search of white paper, trace a closed contour. Think very strongly of **any metal** , **the powder** of which is supposed to be piled **up** inside **this** contour. Set the pendulum on the metal and observe its rotation as soon as it is placed on the vertical of the **traced surface**. Try **several** different metals and find that this is indeed a general phenomenon . Note **0** with the aid of a **pocket pilo** that the polarity **of the** IM image thus produced is the same as that of the **metal**.

2 ° **Supports of mental impregnation** : Set the **pendulum suc-**. cease on rays ES, M S .<•) and note its gyration on **the** contour formed. **Note** that if one extinguishes any one of these three rays, the pondul o set **on** the metal **g é ner a t eur** ceases ; o **ur n e r** . It is therefore a matter of support radiations and not of accompanying rays.

Observe while implementing the identification procedures indicated on **page 159** of the book “ **Dowsing Method Physics** ” . q u o l i m a g c 1 M emits the mental ray and the nervous ray.

3° **There are no IM images . in the second degree.**— Put this excavation of the white paper on contact with the **closed conrour** and **ascertain** that after this contact no part of the **excavation side** does not emit the specific radiations of the chosen metal nor any of the other **radiations** of the imago IM. There is therefore no i m a g o IM of the second **degree**, m C I fortiori, **do** higher degree. -

4° **The Little Carafe Game** : The essence of the little JCU of **society** which makes it possible to realize that **mental** impregnation can be produced on a liquid support and which has **generally** been very successful, **is the** following: A helper gives an operator, Z , c ô r é d o which has been placed a **decanter** full of water, a sheet of paper of which he keeps **the** double ot which contains a list of q r e è n q different metals. The oido, after **being** withdrawn **into** another room,

underline the name of .one of the metals on the list and think **about** it strongly, assuming it is deposited at the bottom of the carafe. For the operator, it is a question of discovering the metal chosen by his partner. This operation is easily done by elimination, after testing each of the metals on the list, by identifying the radiation that comes out of the decanter **using** the procedures indicated **on** page 159 of the book “Dowsing Method Physics”).

5° Radiance of a picture painted from nature : Such a painting constitutes a mental impregnation on the canvas (image 1 M) of the radiance of all the objects that the painter wanted to represent, after having thought about it for a long time and the to have captured **with** the help of the visual ray and the mental ray. On a part of the painting representing the ocean, we will find all the radiation of sea water, on the representation of a warship we will find the radiation of many metals, etc., etc...

6° Radiation of a copy of a painting : The painter who makes a copy has in view only the reproduction of the contours, tones and colors of the original. These are the elements that preoccupy him much more than the elements represented by the master's brush. The copy therefore radiates above all the colors of the original painting. If it emitted radiation from objects, it would be a quadratic LM **image** . Now, this image, as was said in 3°, does not exist. .

For us . to better account for it, let us carry out the following additional experiment:

The contour of the first degree impregnation of a metal having been traced on a sheet of paper, let us trace a second circle on the same sheet and try to mentally project the radiation of the first. As much as we will have succeeded easily in the first case, so we will fail in the second, as we can ascertain this with the aid of the pendulum.

It will therefore not be rash to affirm that an original painting is distinguished from a copy by the fact that, contrary to what happens in the first case, the copy does not emit the radiation of the objects represented. .

1_° Influence of a portrait from life : For the reasons set out in 5°. a portrait executed from nature will emit all the radiation of the person represented, **including** that of the illnesses with which she was affected at the time she posed. If the portrait emits neither the human wave, nor the nervous wave, nor the vital wave, nor the sympathetic wave, etc ..., we can be certain that we are in the presence of a copy . Do. experience on various works.

VARIETIES		2	3.	■	5	'
Abstract		ORIGIN	MECHANISM		ARMY, STABILITY, DURATION	POWER
Direct		1	1	1		
two species	D	Result of an accidental deviation , partial or total, of the inherent radiation of a radiant object, or of an emission source . •	Various	Very different, depending on the case: reflection, refraction, i tion, dispersion, or intervention of particular waves (phenomenon of corners, body suspended by a thread).	Variable, depending on the nature of the accident giving rise to them.	Same as body or source "from m diante whose radiation sart"3- form them.
	M-	Generated and maintained by the action on the radiant source of one or more external energies . the emission of particular electromagnetic waves , which use	Single: MS	• Magnetic images: M produced by the Ile- tio'n of light on the radiation of the magnetic charge specific to the generating body.	Bei indirect images can in relation to ngs, the general object tors;	The indirect images are someti of the same A polarity as the generating body or the original radiant source , sometimes Of opposite polarity. Most often, the polarity of the image corresponds to that of the exterior energy which gives rise to it; but this is not always correct because the law is quite complex.
	ERM	support for the specific radiation of the imago and which convey it. The different nature of the media determines different categories of images with very distinct physical properties.	Triple; ES — MS — RS	Electro-radio-magnetic images: ERM: produced by the action of light on the radiation of electrical charges and. own magnetic fields of the body, as well as on its own radioactive radiation .	Straight or slanted, Looks b l os or deformed, Identical, . amplified or reduced, Permanent or temporary , Fixed or mobile, according to the particular conditions of their formation and, also, according to whether the energy which gives birth to them is itself:	
	AR	We make an imago vanish by extinguishing only one Of its supports.	Single: RS ... ' Doubles:	Radioactive images: A : produced by the magnetic action of light on the radioactive radiation proper to the generating body.	--- YYYY ----	
	EM		Single: RS ... ' Doubles:	Electro-magnetic images: EM: produced by the action of an electric or magnetic field on the specific radiation of a body. - --	Permanent or temporary, - ■ ■Fixed or mobile.	Except for the ERM and ERA images, which radiate vertically or obliquely in the soft directions, the indirect images radiate exclusively towards the sky.
	IM		Duplicate: - ES — RS [Triple: ES — MS — '> j	Electro-radioactive images. : ERA: produced by the action of the terrestrial EM field on the electric charge and the own radioactivity of a body. ~ - Mental impregnation: MI: action of the terrestrial ■ EM field on the specific radiation of a body conveyed by the mental ray of an opera-		

GENERAL ON IMAGES

10

POLARITÉ

GROUPEMENT
DES IMAGES MULTIPLES

PROPRIÉTÉS PHYSIQUES
GÉNÉRALES

'OBSERVATIONS
pour un opérateur dont la main d'œuvre positive

is La same as this **lie ffu** ☐ Very variable depending on **the** case. The p h e n o m e n o d o s or of the ra- **source** ; co ; ns - and that of the suspension give birth whose **radiation ! ii** four images . _ _ _ symmetrical relative **2l los form.** **the** **object.**

' The indirect images are, ' Never isolated, the indirect images, are always of the same polarity
| Days gathered in groups of families.

quo lo generator body Or 1 Each family includes a number of individuals ! the **radiant** Like real bodies , give rise to the formation or ascending or descending **columns** i of families of images analogous to those that we . _ _ is on the vertical of each illuminated body.

j sometimes of **inverse** po la r i t y . , **generator** or Produce magnetic crosses and photogenic **triangles** with **fields** and **groups** of **families of** imagos , but **these** families are o **central** mother EM instead of **M. .**

Most often, the polarity • group includes a number of Limage families corresponding to col- ' , to U . mem o c h ; offer . •
lo of the external energy I , •
• ... d . _ The groups are almost always united on one
who gave birth to him

are **complex** , not always one **more group,** **gr o u p s .**

.nt i IS ^ a xact because the law is quite I Ai ns . , horizontally, los i magi of the Cross | magnetic o form
t

a **stand** of four groups. Those of the double
photogenic triangle, a population of six groups.

In each family, the central image, brighter than **the** others, is of
one dos quo **fro** J categories : M. ERM , **EM. ERA.** ' There are no vertical columns on **1** bone • .

■ The **other** images are all RA or disappear- _ _ _ _ _

are at sunset or **at the** extinction of a
artificial light. When the central image ost
i permanent , the grouping **is** reduced **in 1** ' ob-

nt **cur** ity to a stand of central imagos which

ros **t en t in 1** eur place.

Daytime images

Like real bodies , give rise to the formation or ascending or descending **columns** i of families of images analogous

to those that we . _ _ is on the vertical of each illuminated body.

Produce magnetic crosses and photogenic **triangles** with **fields** and **groups** of **families of** imagos , but **these** families are o **central** mother EM instead of **M. .**

Seem soundless • bles within the radius of an t e n ne . to
the visual ray and **to the** mental ray . _ _

Emit heat rays and heat rays _

chemica
ls.

Permanent images

' There are no vertical columns on **1** bone • .
i ma ge s per man o ntes .

They do not produce n . **magnetic** cross, n . **T r i** ang los pho I
t o g On i cs , but s • **m o** n . back f a mi ll os d imagos à m è I
ro con t rai o **EM** or i enté os North-South . **and** ' East-
Magnetic West.

ERA images are only found in vertical filas. at the
" magnetic cardinal points : of the generating object.

With a “t **6 m o n N** in the right hand, we do not turn on an image of a category other than an **EM**
image .

With a 'witness' in the left hand, the pendulum swung through all the images except the **EM**
imagos. 1 1 therefore oscillates on a real body and an EM image .

By the contact of two fingers of the left hand, of the same polarity
(for example, for the majority of the operators of the thumb and the
little finger) one is generally insensitive to the perception of any
indirect image whatsoever.

In sa.lissan **t** in the left hand a **disc of a subs- . any stance painted**
in black, one renders oneself insensitive **to** the perception of an indirect image
of any side of the country .

(**Process** indicated by **M. Di sc ry**) . _ _

Properties c O m m o n to all i n d i r e c t i m a g e s : _ _

All the **images** form the **ray of** union either with an object or with an
image of any other c . : _ _ _ that.

The six ca . eg o . i e s of i n d i r e c t images are captured by the .
fields of a magnet.

! we hold our breath during the gyration of a regulated **pendulum , this** gyration
which, on a real body, would persist. turns into. oscillations if the instrument rotates on an indirect
image. (P r o c e d e ' of M. Ch o u t eau).

Finally, the Schumfell Pendulum remains insensitive to any image on which it may **find** itself and only
manifests its action on the real body for which it is adjusted . This **is why** its use is
recommended . _ _ _ _ _ _ _ _ _

At verso: Tables 7 and 3.

2. Synoptic table of the particular Physical Properties of. PICTURES. DAYTIME

1	2	3	4	5	6	7	8
NAME BY CATEGORY - BY IMAGE	NAME of the TRAINING	MATERIALS	RADIATIONS MISCELLANEOUS ISSUED BY • THE IMAGE ----- ■! H ^{AA} .. •-.	PROC FROM EXTI	EDES NCHON.	SCREENS PHYSICAL stopping the ray -- Neg. —	ECR RADIEST making the operable 0 la „ A d__Ti pos. +
Pictures . magnetic;	Support single: MS	Action of _ light; on the magnetic charge i of the generative body.	CH. CH CL ! CL	And+ field Field _ M a g.+	Field El- Str Blade field. Mag.-:-	Field Mag.+ El+ field	Neg. — Mag Field— Field El— ! Resist to the rayo n ment of the Black Disc . _ _
(M) —	1.....		a—1	Neck Radius	J Colored cardboard , black. ly their black 1 1	Object on f e black color negative hand.	r or input object of the
7 *	! support:	electric loads ot	1 clean magnetic ' : body in memo time	• j ' ' to,	Field El— C ha mp 1 Mag _ ! Electrical insulation.	Electrical insulation. .	Resist the radiation of the Black Disc.
magnetic ' electro-radio images (ERM)	MS	only on sor, radiation	RS .	CH'CH _ _ CL' CL	Hair P radius Rayom do la Cou or E	Iron blade. Color card , black or blue e.	Point pointing towards A ground or Object of black or blue color held in the negative hand .
Z							
images rac-oacii* es (RA)	Bracket; Magnetic action single do: j the light on the ray- Clean radioactive element i of the generating body.			(1). Gti) . G "- , J ■ P M- , i ' , 0- , Y C H ' ' CH CL CL	• Tip. Radiation ■ Lead ; Where da the Color blue '	Lead blade. Cloth or Cardboard □ of blue color. Any c o u l o r ----- disc	Extinguished by the radiance of the black Disc. Do not form in blue or violet light . They only exist as satellites of an imagmero of another category.

? .cs.'-êûs
■-éra.'es

Daytime images of any category seem insensitive to the antenna ray, the visual ray and the mental ray.

3. 'Summary table' of Properties. Special Physics of PERMANENT IMAGES

NAME D ELA CATEGORY OF IMAGE NAME MODE (s) SUPPORTS : SUPPORTS TRAINING Imag os electro-magnetic E'EM 1 Double ; support i or Mag ES MS : Action of a P El field Ur lo rayonno ment specific to a body.	VARIOUS RADIATIONS Emitted by THE PICTURE pos. + J Neg. i ES + . ES—j MS + i MS—i JK	1 ROC D'EXTIN EDES TION pos. + AEI+ field Maq field. HAS “Poi nt. Radius of a wment Disc close	6 PHYSICAL SCREENS . stopping the specific radiation of the image El o ctric insulation . Iron blade .	7'g 1 _ DOWSING SCREENS making the operator insen- . noecom TIAMC sible to perception OBSERVATIONS Pole Mag+ CPole. Mag— i Pole El+ Pole El—, . Pointe diriqéo vars lo sol. Negative hand held record
Double j Action of the EM field, Pictures ; carrier . j terrestrial on load electro-radioactive i ES electrical or lo radrà- _ (ERA) specific activity of a RS cor Ps. “» ■ *	L M CJ. J* W. rJ- '1 . J -I, u v. I -, VS	Field 1EI+ I Field El— > ! . I Radiation _ of a nearby Disc t —•A— Cham p EI+ Field p El— i Field; Field M'+ e, "e. . Radiation of one. close disc Radiation of Blue OR Black Color	Lead blade. Electrical insulation. Lomo de.for. 1 Cloth or paper ! blue in color or black.	All images _ Load El+ 1 Load El— ges permanent. are destroyed at (Pocket stack). - times by radius _ n upstream of a DisA quo r ap r o ached and . - by that of the Di\$que hand held negotiva. black color. Pole Moq+ l Pole Mag— ! El+ pole Polo El— Point directed towards the ground . Disk held in the hand. negative: •
Tri p le: ■ i Action of the EM field 1 MS impregnation . terrestrial on the rayon- mental: , lies • specific of a Image ES 1 body conveyed by lt (IM) ! mind ray of an op A fo 1 rotator. ' .	1 ES+ 'ES — MS + MS — m, J : (•). ' t~ Spoke . Mental. Royon ' Nervous.	Cham p EI+ Field p El— i Field; Field M'+ e, "e. . Radiation of one. close disc Radiation of Blue OR Black Color	Electrical insulation. Lomo de.for. 1 Cloth or paper ! blue in color or black.	Point directed towards the ground . Disk held in the hand. negative: •

LESSON THIRTIETH

THE PITFALLS

DOCUMENTATION

. It's necessary. get **to** know the pitfalls that constantly tend **to** mislead the beginner. The most experienced graduates register, despite all their attention, failures, because they are sometimes due **to** unsuspected causes. The search for these causes of failure, in each particular case, will therefore be most instructive.

The operator should never say: “ **There** is this, **there** is that”, but-
.: “I think I feel this or that”.

Here are the main pitfalls **to** avoid:

The physical state of the radiesthesist : The nervous system playing a considerable role in the evolution of radiesthesic phenomena, the operator must beware of anything that could influence him • annoyingly: excessive fatigue, insomnia, meals too copious or too well watered, etc.... The operator must be in good spirits and preferably operate in the morning, or after a light meal.

In the beginning, devote only half an hour daily to the practical exercises of each lesson. Repeat the exercises of the previous day. Stop as soon as boredom or the ^{slightest} fatigue appears, the normal play of reflexes would be upset.

One can operate standing, the body straight, without stiffness, and the pendulum quite far from the chest. To avoid the fatigue of the first exercises, place the visible object to be studied on a fairly low support or table, making it easier for the forearm to

suspending the pendulum, a horizontal posture, the hanging hand. .

Avoid holding the free left hand in contact with the hips or another part of the body, which would have the effect of rotating the pendulum with wavelengths close to those which would be appropriate. Close the left hand to prevent its index finger from pointing towards the ground, which would sometimes prevent the pendulum from moving regularly.

It is also possible to operate seated, in front of a table low enough to allow the forearm not to rise above the horizontal, which would cause fatigue more quickly. Do not cross your legs. Have feet. flat on the ground.

Self-suggestion : What is often opposed to initial success is first of all the “lack of self-confidence”, that is to say the fear of not succeeding. The beginner thus puts himself in a state of absolute inferiority.

For some, on the other hand, what leads them to failure is a “spontaneous suggestion”, by too much insurance. These beginners are under the influence of autosuggestion, letting themselves be dominated by the thought that their pendulum will turn or oscillate, or that it will turn positive or negative. Their imagination is thus opposed to the free movements of their pendulum.

The beginner must therefore fight his imagination at all times. He will strive, on the contrary, to obtain from his mind a neutral state, a passive attitude, physically and mentally, as the physical method demands.

Indeed, he will find that with a little practice, it is easy to inspire the pendulum with the movement that one desires, that one wants. Our will can therefore act on it as it pleases. It is therefore necessary to “neutralize our will”. our brain, and to eliminate any preconceived idea. If necessary, think of anything other than your research, in order to let the reflexes occur freely.

The communication of the thought : If, while he works on your ground, you are at the sides of the dowser, saying to you incessantly interiorly : “ Ah ! if he could find water right there , how convenient it would be, how convenient it would be **for** me ! ”. you risk misleading him and making him proclaim an inaccurate result.

But the action envisaged is even more prejudicial if the operator works under the eyes of a hostile or simply skeptical observer who watches all his movements and comments on them. He has many chances, in this case, to cash a failure.

To conclude, we will say that if he wants to escape from the spotty mnuece of the communication of thought, the dowser must compel himself to work alone. If he cannot avoid a few people, let them move away from him at least four to five meters.

The influence of nerve rays : If we cover a coin with a transparent drinking glass , turned upside down, the vertical rays, to be electrically supported, will not pass through the insulation, and the pendulum set on the metal of this part, presented above the glass, will not react. But if we look at the room, by transparency, or even, if we are content to think of it, under the action of the visual nervous ray, or of the mental ray, the pendulum will begin to turn.

Place the same object on a table, and cover it with a piece of paper suitably placed and oriented. The vertical rays do not cross the electrical insulator which is the paper, we will only be able to trace the place of the points of emergence of the rays at 45° which escape from the coin. But if we look at the location of the position of the piece, or even just think about it, the visual nerve rays, or the men ray para. will carry through the paper the specific rays of its vertical waves and we will perceive them on the whole surface of the **dowsing** spot of the room, as if the paper were permeable to electricity.

These phenomena show how much. **if** we are not careful, we may be misled in experiments.

anything, and proclaim, for example, that cardboard, silk or rubber are crossed by the vertical layers **of** bodies. Being warned, we shall avoid the influence of the nervous rays, or of the mental ray. by ceasing to think about **the** experience we are having and, by closing our eyes, in order to let the reflexes occur freely.

The impregnation of the pendulum : We know that when a pendulum has received, for a certain time, the radiations of a body, it is impregnated and undergoes a kind of temporary self-adjustment. It turns permanently , even when it has ceased to be above this body.

If, therefore, we have to discriminate between different **objects on** the basis of the reaction of the pendulum alone (for example determining the sex of eggs), it is important **to** ensure that this reaction is indeed the fact of the object. on which the pendulum was presented, and not of the previous object.

For this, after each operation, we will give the pendulum one or more shocks until it stops rotating automatically, or we will give the pendulum, by its base, a slight shock on the experimental table . , then we will rub the right hand on the left hand in order to cancel the effect of the previous operations.

Fading : According to Abbé Mermet, the waves of radiotherapy sometimes present areas of fading, a phenomenon analogous to the fading of the TSF, and which is only temporary. Since prospecting in an area where success seemed very probable has not given results, it should not be abandoned without having made other attempts under slightly different conditions, or at another time. .

^ . Abbé Mermet further recommends directing your gaze and your left arm towards the sun, or in its direction if it is invisible, by extending your left fingers with an elongated and fine point: in many cases the phenomenon disappears . then after a short time.

The negatives : Some people, fortunately in fairly rare numbers, have the property, by their mere presence, of throwing all those they approach off balance and rendering them incapable of any radiesthetic perception. This is one more reason to operate alone...

The radiesthetic syncope : The absorption of a medicament acting on the nervous system is enough to temporarily prevent the radiesthetic reflex.

For aspirin, for example, the duration of syncope is, in most subjects, about three hours. It goes without saying that, until the restoration of the normal state, one should refrain from any operation.

Disturbing places : Move away, as much as possible, in the room where you operate. metallic masses (copper, aluminum, tin, nickel, silver, gold, etc., which are sometimes also found in drawers) that can produce rays of union with the substance studied. Also be wary, from this point of view, of floors above or below the one where you are operating. The ideal would be, for the first tests, the open air. Subsequently, all these precautions no longer have the same interest; a trained dowser can operate at all times and in all places, without any discomfort or fatigue.

But for the beginner, the electrical **current** , alternating (light and force), the body of the operator and the substance studied constitute three oscillating circuits, whose triple reciprocal action can produce disturbing effects. Consequently, in the beginning, work should be avoided in the evening, under electric light. Some authors condemn the wearing of rubber soles and silk socks.

With a little attention, observation and above all training , it will be easy to eliminate all these pitfalls.

It is possible that your beginnings are slower than you think, that you even find, at times, tiring **and** discouraging your exercises at the beginning. You will have to persevere, methodically, and train regularly. Quite natural skill is acquired in the long run for some; on the other hand, most obtain it very quickly.

ADVICE TO THE READER

The shortage of paper and its poor quality oblige us to limit ourselves, in this year 1943, to the **First Part** of the work that we had planned: GENERAL INSTRUCTION **and** DEVELOPMENT OF **THE SENSITIVITY OF THE BEGINNER**, in 30 Lessons and more than 150 Exercises .

We apologize to our readers and we •
momentarily ask them to refer to the work : “THE RADIESTHESIA **explained and practiced by the PHYSICAL METHOD**”, in particular to Chapters V and following, which already include many essential applications and developments on practical prospecting.

MENTAL METHOD

APPENDIX

ESSENTIAL CONCEPTS

of

RADIESTHESIA

following the

— **MENTAL METHOD** —

(CLASSICAL PSYCHO-PHYSIOLOGY AND METAPSYCHIA)

by

Pierre BEASSE
rechnicien-Council

Visa do controla of the Consuro Scientifique, General Secretariat
of Information : N° 8.089, of September 30 , 1943

Copyright by Pierro BÉASSE, Publisher, 1943

Reproduction, translation and adaptation
rights reserved for all countries

TABLE OF CONTENTS OF THE APPENDIX

on MENTAL Dowsing

	Pages
GENERAL	231

CHAPTER 1

The Instruments	238
-----------------------	-----

CHAPITRE II I

The "Mental Sense"	240
The mechanism $\wedge d \wedge j \wedge S \wedge es \wedge et \wedge al. \dots \dots \dots \wedge 2 \wedge 4 \wedge J$	..
Process of Mental	Inquiry 244

CHAPITRE III

The “mental desire ::” or “mental orientation:;>	245
“Mental selection:.. ..	246
The “mental convention”	247
The " mental questioning "	248

CHAPTER IV

Means of identification	250
General observations	251

CHAPTER V

The elements of perception.....	254
---------------------------------	-----

GENERAL

If it is true that the **Physical Method** (passive aHitude) requires neither special gift, nor mysterious or complicated initiation , that it

only proposes simple and rational procedures, which put it immediately within the reach of the great majority researchers, and as such, it should be recommended to all beginners, it is no less advisable, if they find it advantageous, to invite, by the . sui ie, these same researchers initiated to the physical method, to try the **Mérhode**. Mental (selective attitude or intuitive attitude), sometimes responding better to their special sensitivity or their particular abilities.

It is intended for those people who are psychically gifted, and because this exposition of the Mental Method. We have been very often asked that we have also been led to ascertain the internal subjective fact (physiological and psychic) which characterizes the mental method and which is just as indisputable as the external fact, physical or biological.

Our goal is to present in these few pages the essential principles by coordinating the knowledge acquired to date in this field. It will be, to a higher degree in practice, a complement to our physical method towards which our preferences go, however, in the initiation of beginners.

It **has** been said by authors who advocate **the** Mental Method, that the operators of the **Physical** Method do psychism without knowing it. It has also been said that exclusively mental operators, proceeding only

according to the Mental Method, also bring into play, without knowing it, light rays, azimuths, magnetic influence , etc., or adopt physical methods for calculating depth, flow, etc., and thus penetrate into the physical realm without their knowledge.

This is why, it can be wise to conclude that the radiesthesic operator acts more **or** less physically and mentally according to the capacities **of** his human compound: thus the two methods join for the capture •of the radiations. Doctor Andrée Besson, president of the Association des Médecins Radiesthesistes

de France, wrote : "To coordinate the triple physical, physiological and psychic education, such is the fundamental goal of the teaching of radiesthesists, **if** it rational and complete.

This amounts to confirming that Dowsing is " One ", whatever the way of practicing it.

It would therefore be a mistake to want to establish a “ dissociation ” **in the form of two** dowsings , one physical, the other mental. The need for the differences to be adopted arises only from the possibilities of the operators, from their degree of sensitivity, from the very different psychic gifts which devolve upon them from one subject to another. Because it is always necessary to - envisage the Dowsing in the human compound in oneself, and the study of the physical method must precede that of the mental method, this one being less with the range of 'out of the world.

*
**

Remember that there are three possible attitudes for an operator:

1. The passive attitude (that which is specially suited to physical processes) ;

selective attitude (in which the will intervenes in the choice of the waves to capture). mental process:

3° **The** intuitive attitude (which only suits people endowed with clairvoyance), a mental process.

We maintain a perfect unity of doctrine, by exposing in the following pages a Mental Method, based on the selection of waves, which is none other than what Mr. Christophe calls "mental orientation", and Brother Padey the "fluid of intention".

Physical operators abide by rules; by . On the other hand , the minds are most often concerned neither with images, nor with series, nor with wavelengths, nor with regulation, nor with é e e .romagnetism, etc... and yet, better than the physical ones, they penetrate psychically into immense domains to be explored, in spite of the complexity of the matters to be analysed. Abbé Mermet said: "It will be possible to prospect the human soul, as one prospecting for metals". And M" 'c de Mersseman, virtuoso of mental Radies thesis, would write to us in 1940 : "You have done well to republish your work on "Kadiesthesia Method Physics", because it is clear and practical for many people. In my opinion, the "Physical" Dowsing must be continued by the "Mental". This is not **within** everyone's reach; however, what complementary joys it gives! »

A little later, M" 'c de Mersseman wrote to us again:

"Our colleague, Pierre Béasse, asked me to explain my point of view on Mental Dowsing. Some

- believe that this is completely different from Physical Radiation thesis. I don't think so, since people accustomed to the mental method can easily carry out the experiments of their fellow physicists.

"Thus, I read in the work of Pierre Béasse on the Physical Method, pages on the work of MM. Engineers Voillaume and Correnson, on the adjustment of the pendulum, recommended by M . from France, on the waves, the harmonics, the interferences, etc... It is clearly explained. But in practice, a teleradiesthesia will leave the reasoning there, the search for cause and effect for

to use above all the mental method, as much as he can.

“Here's why: When it detects, the wave planes, initially remanent, follow each other rapidly. The operator must observe them with attention, orient his mind to have them in the direction of the research and change the length of the pendulum wire quickly in order to obtain gyrations in the fields of present influences. In the long run, sometimes, the detector seems to dominate the thought of the researcher, enlightens him by certain unexpected evolutions, which form, for example, the image of the things observed. This is why some rhabdologists question themselves by the pendulum. They can have good answers if they know how to control their imagination.

“We don't. recommend the mental method only to the most gifted, to pendulum virtuosos. There are many who ignore each other. Like the great violinists, they become one with their instrument, already following a short training. They quickly know him thoroughly and know what he can give. They vibrate together in the astral, which ends up giving them the secrets of Nature, its lights, its forms, its advice.

"On the surface, there seems to be a disagreement between the 'physical' and the 'psychic'. The physical wants to follow the possible experiences step by step, as in a laboratory, and not to go beyond what is physically controllable. The "psychic" can also be a scientist, but feeling in his subconscious and in his well-trained nervous system that there are other data, wants to observe further. So he searches into the unknown and soars to the stars to find out if they don't know the subject that interests him.

“I read on this subject in the “Revue d'Astrologie” of the distinguished mining engineer Dom Neroman, in constant contact with serious dowsers, that the stars know what man often seeks, the secret of his destiny. They reveal it to him with a logic, a sequence that marvels. Nature, when she shows us her beauties, does it with science !

“This is why many Confreres end up thinking, like me, that the studies of the two methods, physical and mental, come together in a complete scientific conjunction. In short, everything is science, and Dowsing is the bridge that connects the science of physicists or that of occultists, psychics, teledowers.

“Those, by the radiation of their fine sensations, penetrate into the infrared and then acquire new possibilities. They can thus compare their work to that of a wireless set, because they pick up very distant waves. From this follows the detection, on plan, of everything that exists on and in the earth's crust. Some “artists” even seize in space and time, diagrams relating to such or such deceased

character. By means of a "witness", they feel the radiations of a deceased who is dear to them pass through them.

"Scientific colleagues will perhaps think that I only have enthusiastic words for mental Dowsing, which is why I will end by telling them all the hopes that I put in them for the codification of the principles of the two methods in one science. Helped by psychics who perceive very far, the scientists will live the hope of the human race. Also, it is with great pleasure that I send a congratulations to these pioneers of a great and new science".

Thus expresses **Mrs.**^{ra} * of Mersseman, former professor of mental dowsing, very famous.

:t:
**

The processes of the mental method being often purely personal, the beginner will first realize the natural reaction of his pendulum, apart from any conscious auto-suggestion. He will write it down and check it many times, several days apart. constant regularity, before adopting it as definitive. The beginner will also realize that, in many cases, the movements of his pendulum will not respond to this natural reaction, but that he will have to adopt

a prior mental convention on the movements **to** be expected from its pendulum. Thus, his pendulum will respond to a conventional reaction.

As much as possible, it will be to his advantage to admit the codification of the most generally accepted conventions in the dowsing circles of the mental method, in order to avoid a new Tower of Babel. It will be, within this limit, the only acceptable conventional auto-suggestion, because of its positive advantages and for the progress of the dowsing.

We also recommend to the beginner to penetrate the difficulty by adopting two progressive levels of studies : 1° Exercises on visible objects, therefore certain ; 2° the Exercises on different known objects, but covered without the knowledge of the beginner. by an assistant; therefore invisible objects.

In these two cases, which will constitute the personal study of its own sensibility, the beginner will note, with the day. during the day, in a notebook, his smallest observations which he will compare with each other with perseverance. He will thus acquire a personal, indispensable experience that no teacher, however excellent, can inculcate in him. *

From random, his progress will become certain, unquestionable . Long work and severe discipline, constant personal discussions, will have been the basis of these prodigious results.



A definition which would seem to relate best to the mental Dowsing would be : “ the sensation by the nervous system and the subconscious of the radiations of beings and things ”.

Vis-a-vis this astonishing art, the independent spirit will not reject the facts. Quite the contrary, he will check them, analyze them, question them to unmask the prejudices of good or bad faith, and come to terms with the facts.

the human body be infinitely sensitive ? Recording sensations, could it not be a transmitter and receiver of electromagnetic waves, of radio-energy?

active, of vital fluid 7 To assign limits to this sublime mechanism that is man would be to lack scientific initiative. .

Everything radiates in inert or living nature. However, certain subjects have an exceptional disposition, a more or less accentuated "gift", for capturing radiation according to the mental method. And

as in music, we notice virtuosis there. Since many people are “born dowsers” who do not know themselves, why not facilitate their discovery ? Their "numan compound" is conducive to mental detection. All it takes is a little practice for them to reap the most valuable benefits in all areas of daily life.

It will be noted, however, that the human organism being one of the main elements in prospecting, its reactions will always be linked to the more or less favorable psychological state in which the operator will find himself at that moment.

The purpose of this short presentation is, above all, to specify the essential bases of the mental method, so that whoever can, can use it usefully.

We urge our readers, after a short initiation into the Physical Method, to gradually try out their abilities in the selective and even intuitive Mental Method: they will perhaps discover natural possibilities — what is called the "gift — . which will save them precious time in the practice of this amazing art. Very many are the gifted dowsers who are unaware of each other. .

INSTRUMENTS

Unlike most animals, and especially insects, the nerve centers of man have . need, in the majority of the cases, to feel and mentally select the external waves, of the intervention of a "support". some would say, of an “amplifier”, others would say; the wand and the pendulum will be ideal. We will therefore use the same instruments as in the physical method.

The wand : We refer the reader **to** what has been • written on its construction and holding, page **34** of this book.

The pendulum : We will not go back further on * the presentation made • about it, page **39**. We will recommend . however: — either a black neutral spherical pendulum, of the PS type, **mentally tuned** to the wave or radiation to be received, the process of which is described on page **81**.

— Or, preferably, a magnetic **pendulum** . such as Schumfell Developer, **without 'adjustment**.

Indeed, in the mental method, there is no question, **in principle** , as is essential in the **physical method, of precisely tuning** the pendulum to the wavelength of the radiation or of the body. **to** detect. For many operators, the length of the suspension wire remains approximately 10 centimeters. Definitely, for all bodies and radiation. For some, **the** length of the wire is about half a centimeter, that is to say one of the shortest line lengths. For others, however, the maximum gyration is more frank if they

seek, without precision moreover, the best length of wire for a given body or radiation.

As we can see, everyone will decide according to their individual conveniences, remembering that Abbé Ferran spoke simply of "favourabilities". length of wire, while Abbé Mermet had come to precise adjustment, which he considered "very important to do a good job".

In mental method, the magnetic North-South orientation of the Schumfell pendulum ring (see page **121**) . is not necessary in stretching exercises on visible objects. This orientation will however be an excellent precaution, later, during practical prospecting, in the fight against "images": it will thus avoid many errors, for operators who feel more or less sensitive to images, one of which complete description has been given on pages **183** and following.

*

THE “MENTAL SENSE ”

We have always thought, with many authors, that dowsing, seen as a whole, belonged to classical psycho-physiology, on the one hand, because of the obligatory mental action in research, as it belongs to physics , on the other hand, since his concern is, among other things, the perception of the radiation of matter. But we will not exclude the examination of its attachments to metapsychics, when it implements the interesting phenomena of “ mental convention and questioning ”.

In the course of dowsing by the physical method, which precedes this appendix, we are above all attached to perceiving this radiation of matter which is the proper and limited domain of this method. In the following pages, our attention will focus more on the part of the mental action of the dowser, whose domain extends not only to the mental perception of invisible bodies, but also to research of an intellectual, immaterial order.

And each, according to his special dispositions, or the training of his sensibility, will ultimately choose the method which best corresponds to his aptitudes, or will use both, according to the circumstances.



We have pointed out that most animals, and especially insects, possessed an instinct enabling them to select and smell. We said that in man

civilized, this instinct, enlightened by intelligence, was, in our day, very wrongly, neglected, for the sole benefit of intelligence and, as a result, had atrophied; that it would however suffice to awaken and develop it, which is easy with a little practice. •

But what is, basically, this human instinct ? **We** are still reduced to imprecise hypotheses, although many authors agree with us that this mental **sense** of the radiés thesiste is entirely dependent on his **subconscious** . .

Metapsychists **believe** that this subconscious, also called unconscious, is a "second person" in us, having an activity alongside consciousness and will. We gladly agree with this opinion. As there is in us, the " subconscious self", or " unconscious self", there is also in us the " conscious self", intelligence and reasoning. It is moreover through our "conscious self", that is to say through our intelligent thought, that we solicit the activity of our " subconscious self".

This subconscious, or inner sense, or sixth mental sense, as we want to call it, assists us in what cannot be seen, recognized, felt by our other five outer senses.

This subconscious is a real guide in the immaterial order, as are, in the material order, the other five outer senses. It guides us, while being a kind of liaison between the different faculties of our thought, creative or ideas or movement.

We are going. account, therefore, of all the use that can be made of this "second person", of this subconscious which is in us, which most people are unaware of, or if they have a vague knowledge of it, abandon. absolutely - useless...

The truth is that to use the subconscious requires a concentrated and sustained effort of thought, unpleasant at the beginning of initiation, and one more easily meets people willing to make a physical effort, walking, singing, etc., than to make the more delicate effort of attention and volume for the solution of a problem

abstract intellectual, in the exercise of an art, which because of the state of empiricism in which it has long been, does not exceed. hardly, **to** torr, the framework of amateurism, **apart** from a few thousand trained and experienced professionals.

The truth is also that one cannot silence an appeal **to one's** subconscious — and this is absolutely recognized by all practitioners — except for praiseworthy and honest reasons. Otherwise, the subconscious, reproving .all that goes **against** conscience, altruism, good, or fails to respond, or lets the provocateur get bogged down in 'failure and error' .

On the other hand, when we seriously consider the prodigious material and intellectual wealth that can be drawn from the subconscious, even in all the most ordinary circumstances of life, however little we want to solicit it, we remain the thought that such a treasure had been thus neglected, abandoned.

**

How can we explain that we have been so long away from so great a source of precious advantages ? This is because, on the one hand, dowsing, the last trace of the secrets of the highest antiquity , vegetated until our days in a certain empiricism; it is also that science considering that abnormal (metapsychic) powers do not . are not scientifically explainable, avoid introducing them into his field. Faced with their obviousness, she does not dispute these powers of the metapsychic (.erm created by her, moreover), but she considers them as scientifically non-existent. We do not have **to** discuss such an attitude here, but in such conditions of reluctance, sometimes of marked hostility, the dowsing could not progress.

In summary, if the spiritualists give the explanations that we know, the materialists are content to remain in scientific skepticism. And yet, throughout the world, cultivated minds, great intelligences continue to speak of "subconscious, of consciousness, of soul, of inner personality beyond the borders of the material body,

etc .. » . »•

Be that **as it** may, not disputing, as for us, the existence of the spiritual, involving the existence of the soul, we support, with many other authors, the theory of the mental method in dowsing, on its essential foundation: the subconscious.

The mechanism of mental sense

All our training will therefore contribute to developing in us the activity of our subconscious, our unconscious self , our second person.

For this purpose, we will always remember that in the solicitation of our subconscious, we must absolutely dismiss the action of the will, of intelligence, of memory or of reason, all conscious faculties, that the subconscious is however willing to help.

We will see further that the will will intervene however, but at most to support our attention and our desire, in the radiesthetic research.

In any case, our subconscious will respond to the call of our desire, as soon as we have met the required conditions, to assist us in our daily, material or intellectual difficulties, and this only within the limits of the exercise of good.

By material difficulties, we mean the multitude of discoveries which interest us, of what is hidden, invisible, and which we could not, without the intervention of the subconscious, procure for ourselves, in order to satisfy our natural or protect us; search for lost objects, water, metals or precious products, material analyses, etc... The response of the subconscious will result, in this first case, in a neuro-muscular reaction transmitted by the hands to the amplifier what is the pendulum or the rod.

By intellectual difficulties, we mean the vast field of questions of all kinds that arise in our daily life, **including** views on the future: research on plans, photos, writing, witnesses, concerning vitality, profession. qualities and defects. the state of health, etc... The response of the subconscious translates

in. this second case by the spontaneous emergence, in our mind, of ideas or thoughts relating with precision and perfection to the question posed. Here again this thought is creative of the manifestations of the pendulum and the rod. It goes without saying that this last kind of mental dowsing, which would be more akin to metapsychiatry, requires a more intense spiritual development than in the researches of a material order in the domain of classical psychophysiology .

Mental inquiry process

Success will be all the more perfect if we have succeeded in bringing together the following conditions as well as possible:

- 1° Physical, physiological and psychic state summed up in an attitude of calm, of absolute self-control, of material and mental isolation, and finally of self-confidence;
- 2° Careful implementation of the mental principles which are the subject of the following chapter.

CHAPITRE II 1

"MENTAL DESIRE " OR "MENTAL ORIENTATION"

It is a purely psychological fact, depending on the phenomena of the human soul, of the **subconscious**, and not of the organic senses.

This means of detecting radiation is called, according to the authors, " mental desire ", or " mental orientation ", or even " fluid of intention ".

He asks, in the appeal addressed to the subconscious, only the intervention of two faculties: attention and will.

Attention:

1° By **concentrating all his thought** solely on the object or the problem of the research; creating a void around oneself, to the point of seeing the object in imagination and being insensitive to **any** external distraction:

2° By **desiring to become receptive** to the activity of the subconscious ;

3° By **avoiding any emotion**, any modification of the normal rhythm of breathing.

The will:

1. By reinforcing the **desire or the formal intention to be receptive** to the visible or invisible radiations of such and such an object, if it exists, or to be enlightened on such and such a question, if the solution to it it exists;

2° By intervening in the **selection** of the radiations **to be captured**, prohibiting the reception of any other radiation (selective attitude):

3° By becoming neutral as to the results to come.

Thus the "mental desire", under the control of the attention and the will, causes the mental detection of the radiations of the object of research.

This desire still needs to be expressed in some form, simple, short and clear. This mental desire must also be expressed in the subconscious in the particularly supple form of an invitation, of a prayer, thus excluding all demand, all will, a fortiori all violence. And, as a prayer is more "semie", more perfect, when it is said aloud, than in a low voice, or internally, the oral form heard clearly by the operator himself will be the most indicated.

This is how all practitioners of the mental method have adopted the now classic formula: "I want to make myself sensitive to such radiation, from such body, or object." of such handwriting, such person, etc.

Engage in some exercises.

Let us note in passing that this same mental desire, strongly supported on such and such an object of research, is enough, for many experienced operators of the mental method - by a sort of appropriation or mental agreement - to no longer make it necessary to setting their pendulum on the object in question. Indeed, this facility is naturally offered to the mental dowser, as soon as he acquires a certain operating habit.

. MENTAL SELECTION •

The subconscious is the basis of mental selection. Just as it is possible for our ear to discern more particularly such and such a musical instrument in an orchestra, just as it is easy for our eyes to adapt visually to successively more distant planes, up to to perceive such insignificant detail in a vast and distant landscape, in the same way it is possible for our subconscious to select among the innumerable wave emissions which assail us from all sides and at all times, the only wave or radiation that we are looking for and which interests us.

In other words, when thought has made its choice on a radiation, among the multitude of radiations which cross us and surround us on all sides, this one will be retained by our subconscious only from the moment when our thought takes care of it with attention and will. in the intense desire to be receptive to this wave.

Whereas in a physical method, the selection of waves and radiations is made by a precise adjustment of the suspension wire of the pendulum and by the addition to the pendulum of a "witness" (filter of radiations), the pendulum no

longer reacting that in the presence of a body identical to this witness, in mental method the selection is made by a mental adaptation of our subconscious to the wave or radiation sought.

Practice it.-

MENTAL CONVENTION

Quite difficult to analyze, the “ mental convention :. seems to come from metapsychics, that is to say from the science of the phenomena of the human soul, which goes beyond ordinary psychology. This is what makes it (like 'mental inquiry ') so different from 'mental desire '. We then seem to be in the very special domain of "psychesthesia " (intuitive attitude), also called, perhaps improperly, "psychic dowsing ", because the pendulum which is used in the mental convention seems to be only as an accessory, in the absence of any other device used by occultists or people gifted with clairvoyance.

The “mental convention ” establishes a general rule, accepted by oneself alone or by a group of users. It results, not from the nature of things, but from a tacit, conventional, prior and precise agreement with our subconscious to understand and interpret the movements of the pendulum . The latter then adopts exactly **the** movements which suit the circumstances, movements studied in advance.

Indeed, if the operator bases the movements of his pendulum on prior mental conventions, the results are all the more comprehensible and certain, as these conventions will have been more precise and as his attention will have been more sustained.

Let us give as an example of mental convention the arbitrary adoption of any unit of evaluation to measure the flow rate of a source, by what is commonly called “mental calculation ”. It consists of a convention of measurement, prior, with the subconscious, which responds by the pendulum. Thus, having discovered a source, you carry yourself on the vertical of the current, where the pendulum turns at its maximum. Having agreed that each gyration will represent a litre, or a decalitre, or hectolitre, etc..., you venically raise your pendulum in gyration; it then begins to oscillate, then to turn. You count how many new gyrations are carried out in this way until the stop, determined by your subconscious, to let you know the number of units to retain, that is to say the measurement of the flow.

The most embarrassing cases can thus be solved by mental convention.

For some operators, it is absolutely appropriate (as we have already

seen in the case of the expression of mental desire) that the mental convention be expressed aloud, otherwise, for them, the pendulum does not react appropriately . , or not at all. This means that in such cases the inner concentration of thought is insufficient.

Perform a few such exercises and repeat them.

MENTAL QUESTIONING

As much as the "mental convention", the " mental questioning" seems to belong to metapsychics, going beyond ordinary psychology. Being in the realm of "psychesthesia ", it allows us to successfully ask the subconscious questions ad infinitum, in the most different domains, within the framework of good and honesty.

That is to say that the operator, thus superiorly gifted, will have to surround himself, in controlling the results, with the greatest prudence and a great deal of modesty.

These operations, which we consider clear-sighted, can only be undertaken successfully after a certain training in material research (with a selective attitude).

In the opinion of many authors, research on plan, phorography, writing, etc., can be considered among the immaterial research leading directly to mental questioning.

Let us first point out that the same conditions of attention, and of will, required vis-à-vis the concentration of thought, the desire to become receptive, mental selection, and the mental convention , must precede the mental questioning.

The question should be formulated calmly, simply, with clarity. The more intelligently it is asked, the quicker and more precise the survey will be. The operator will have the greatest knowledge in the feeling that the reality, the truth can be thus known.

The usual sentence is this: "I want to be honest with you on such and such a question", followed by: "Is it true that..."; or "I want to make myself sensitive to water radiation", followed by "Is there water here?" » ; or "I wish to make myself sensitive to the radiation of such a misplaced object ", followed by "Is this object here?" », « Is it. in certe d nec r we ? ". and so ...

r-it;let's take the case of an unknown fruit before us: Is it good or bad for your human health? I concentrate my thoughts on this result (attention and volume, selection) I thunder aloud my desire to make myself receptive

to its desires (desire), excluding all other exhausting radiations (selection) . or I will advance that the response of my subconscious will be positive (yes, if my pendulum turns clockwise; or that the response will be negative (no; if the gyration takes place in the opposite direction, that this answer will be, finally, doubtful, or unaffected, in the case of swinging (oscillations) of my pendulum In practice, the succession of these mental operations is very rapid.

The application of these principles is therefore easy in innumerable cases. He presents himself to the court at any time; of a day. This is an opportunity for many trials and exercises.

MEANS OF IDENTIFICATION

We studied, in our “ Course of Physical Method”, the **physical processes** of identification of the bodies which are the **adjustment**, the series or **particular number o b i e c t s** , the **fundamental ray** . _ _ _ It will suffice, on the other hand, in the mental method, to establish, prior to prospecting, a **mental representation** of the image of the body with the sensation of perception that one has fixed by a convention of personal mental ion.

Most mental dowsers consider the only three controls that constitute the **mental witness, the fundamental ray undergone, to be sufficient. if**, the subjective series .

The association of ideas which they have thus applied mentally, once and for all and particularly **to** each body or radiation, gives them, **in** training, such a mastery of their sensibility, which—taking into account the the more or less favorable psychological state in which they find themselves at the time of their activity assures them of remarkable security of operation. ■

Faced with such convenience which is offered to them by their very special "gift", we do not hesitate **to** recommend to new beginners of the mental method to adopt, for the ease of relations between the adepts of the method physics and themselves, the objective results obtained by physical processes in terms of polarity of bodies or of fundamental radius, or of particular number of series. They will not thus have **to** fix themselves a mental representation at random of their inspiration, but will adopt precise data physically established and appearing in the Table of identification of the simple Bodies, page 96.

GENERAL OBSERVATIONS

In principle, the **Mental Method** requires the following conditions: *

1° Strive to let the pendulum move by itself; consequently, the **hand** and the **fingers** will be compelled to a maximum physical **passivity** .

2° To bind oneself to an absolute **neutrality of mind as to the results** which will ensue; therefore self-suggestion **regarding** these results should be avoided.

3° On the other hand, to restrict oneself to an **activity of mind** pushed in the **desire** that one has to see the subconscious answer **such** a question of the present, or of the past, or even of the future, by the pendulum or the wand, or of the **desire** to perceive the radiations emitted by such a well-defined body in thought, if this one exists, and **to** the exclusion of any other body. Then, wait until the pendulum or the rod are brought into contact, by the subconscious, with this body.

To have the firm opinion, the very assurance, to finally believe, with more resolution than the reasons persuade, that the dowsing "fact" will then inevitably occur. if the natural conditions exist, there must be the state of mind. Thus stubborn, thought will go its way with fearlessness, despising all obstacles. They cannot be disappointed by the always exact manifestations of the subconscious by the pendulum.

The various manifestations of the pendulum (oscillations, gyrations , series, etc.) or of the rod (straightening, immobility, lowering) will then be revealed **according to the code** that we will have previously and personally established for ourselves; or **according to collectively accepted rules** to form a general method, ensuring a uniform interpretation desirable for all dowsers, whatever method they practice.

However, there **are** three pitfalls to avoid:

1° To want to obtain or to **expect** such a result, to such an extent that the subconscious commands or pushes in some way to make the pendulum turn or oscillate obeys, so to speak, our will. This would be to fall into the grave error of an unfortunate auto-suggestion.

On the contrary, it is advisable to remain impartial, but attentive, and ask questions, for example: "Is there water here? Is it drinkable? », etc.

Auto-suggestion still consists in having the **impression**, even **spontaneously believing** ^{that} an invisible body exists under our pendulum and in these illusory conditions, estimating that the pendulum or the rod must logically react on this imaginary body. In a way, this amounts to "commanding" by apparently logical deduction, to one's pendulum, and the latter is executed. Indeed, because of such a mental influence on the pendulum, it is to be noted that the action of the brain inadvertently produces the muscular reflex which fatally involves the movement of the pendulum.

Be that as it may, the harmful action of conscious, or unconscious, or disordered auto-suggestion, being thus revealed, it is easy for the beginner to avoid it.

. When in doubt, it will be very wise and convenient to control the reactions of the rod using the pendulum, and vice versa.

2° Exaggeratedly fearing auto-suggestion : Thus, imagining that the slightest movement of the hand, communicated to the pendulum, is the result of a nervous reflex incompatible with the dowsing reflex and, for this, stiffening, wrongly, arm, hand and fingers.

3° Misinterpreting the observed manifestations, for lack of a previously well-defined convention. This prior agreement should not be established at random. It depends either on your own previous observations on the behavior of the pendulum with regard to you, or still on the collected observations of professional dowsers.

Of the choice between the “Schumfell” pendulum and the wand

Thanks to its particular sensitivity in a large number of cases. the pendulum - “Schumfell” has the advantage. in mental method. to be able to be handled quite easily, without precise **adjustment** , **nor orientation**, and to be more precise than the rod whose use often requires more experience. The Schumfell is also slower in its manifestations than the study PS black neutral spherical pendulum, which makes it easier to interpret the results.

Furthermore. By orienting magnetic North-South (by means of the compass) the great ring of Schumfell, prospecting operations are considerably simplified by eliminating multiple “images” of any kind. who watch at all times for the most knowledgeable dowser. and which are so many “mirages” in relation to reality. A full description of the Schumfell is given on page 46. Professional - radiologists use it routinely, considering it a first-rate control instrument.

However, on the ground. any pendulum is more easily thwarted by walking, wind and rain, than the rod. This is also more convenient for quickly “clearing” the ground to be ^{explored} . by immediately identifying the most interesting points to work on afterwards with the pendulum. This one is also best suited indoors and for card work, photo. plan, etc.

It still happens that certain dowsers, excellent at the pendulum, are not sensitive to the rod or vice versa , fault, most often. to have practiced sufficiently with one or the other instrument.

THE ELEMENTS OF PERCEPTION

As in the physical method, we can use the following means in the mental method to reinforce our action and make it more effective:

The visual ray : This nervous ray **leaves the eyes** to move towards the object sought and provokes the reaction of the pendulum. It corresponds to the denomination of "radius capital", in some authors.

The digital ray : Also called "antenna ray", this other nervous ray **comes out of the index finger** or arm, as an antenna, to move towards the object sought.

The mental ray : Studied on page 176, it is the result of our thought on our subconscious which triggers the reaction of the pendulum as soon as this mental ray encounters the object sought. This ray is the most important in the mental method, and also the most convenient. Indeed, it instantly connects the subconscious of the operator to the body sought, at any time, and from any place where the operator is.

1 o Being **on the ground**, it is enough to pivot on the spot, pendulum suspended with the fingers, arm and index in antenna, and to carry out circular turns around. self, towers moving away - successively by 10, 50 or 100 meters, etc..., according to the needs, to prospect strips of land more and more distant, until one encounters the reaction of the pendulum on a distant point, aimed with the eyes and the index finger, and as soon as the mental ray has encountered the object sought.

When one has thus acquired a distant point, in any direction, it is necessary to clarify this point by cross-checking two or three other similar investigations. We then go to two or three other places of departure, and it is at the point of intersection of these other "aims" with the first, that we obtain a precise point.

It will then be appropriate to go to this intersected point to perceive the vertical rays, either of the object sought, or the limit of such a deposit, or the shore or banks of such a stream of water sought. The best is, often, to bend down, or even to squat, the base of the pendulum about 5 centimeters from the ground, the index finger of the free hand in antenna, touching the ground. This is how the best results are obtained.

The use of the visual, digital, and mental rays combined constitutes, in itself, one of the best methods of perception.

2° Working **on** a plan, or on a photograph, or a geographical map, or an anatomical map, our impressions will be exactly the same as if we were on the ground. We will proceed in the same way. It will suffice to suspend the pendulum by the right hand, base **of the** pendulum at the height and away from the map, oriented magnetic North-South, and to walk for example a very pointed pencil and held very vertically by the free left hand, on this case, to trigger the reaction of the pendulum, as soon as our mental ray and the vertical point of the pencil will be on the point where the object sought is. This work of 'ele^dowsing will require, it goes without saying, much more will and attention, desire and selection than the work on the ground.

There **is** always an interest in prospecting only on geographical maps of the largest possible scale, giving as much detail as possible; thus the radiesthetist becomes better aware of the atmosphere of the place. A sketch, even in pencil, but very precise, is sufficient.

As can be seen, there is hardly any difference in process between work on the ground up to a circle of visible horizon, and work on a geographical map which amounts to prospecting a circle of invisible horizon, located sometimes thousands of kilometers from the operator. When working in the field, the operator realizes "visu" the detail of the terrain, whereas on the map he "represents" by the precise details of the latter the true configuration of the terrain, which amounts to the same.

In terms of depth, either on the ground or on the map, the operator imagines that a kind of decameter slides from the surface of the ground 'in the interior of the earth to the surface or the bottom of the source located, or to the buried object, and thus measures the depth. As for the flow, it suffices for him to count how many strokes (or gyrations according to the previously adopted mental convention) the pendulum will make, each stroke or gyration "representing "

for him either one meter or ten meters, according to his prior convention.

With a little practice, you can then, by “questioning” yes or no the subconscious, to obtain information of all kinds. These facts are continually verified in very gifted mental dowsers.

. The photogenic ray : Studied on page 128, it is emitted by any light source (sun, artificial light, moon). He heads straight for the wanted body. When the pendulum meets this ray, it reacts.

The use of this photogenic ray, of the mental ray and of the fundamental ray still constitutes for other dowsers a sufficient whole to assure the control of prospecting.

Lines or zones of influence : These are vertical rays, or with fixed inclinations, or at 45° or 135° . Studied on page 62, these rays are as many emissions (vertical, horizontal or at 45°), starting from the body sought, on which the pendulum reacts successively.

In the case of the search for a source, the rays at 45° come from each of the two banks of this current. The meeting with the ground on each side of the current of these two rays at 45° constitutes what dowsers commonly call the two great parallels. Between these rays inclined at 45° and the vertical rays emitted by the current, there are six other lines coming from rays with fixed inclinations, called parallels, more or less perceptible. The distance on the ground, between the vertical radius of an edge and the radius at 45° of this edge, gives the measure of the depth. (See page 169).

The subjective series : Not everyone is sensitive to series. Contrary to the objective series, studied on page 105, the subjective series figure is the result either of a very personal observation, or of a pure prior personal agreement.

After a normal gyration on the vertical rays of a body, it suffices to slightly raise the pendulum with a flick of the wrist, which begins to oscillate. From this moment, note first one or two or three ellipses, then a first giraiion, then a second, a third and so on until a new or more ellipses. It is the total (or figure) of these gyrations which constitutes the subjective series. This figure is personal, if you have not previously agreed with yourself on a conventional and constant series figure, for a given body.

If you do not adopt a table of conventions such as there is one on page 96, you will have to establish one yourself. You will then search for the subjective series of many usual fields, and note the result. This work completed, and checked several times, it will happen that you will now be able to use the subjective series as a means of checking for the identification of buried bodies.

Indeed, if you then wish to be receptive to the series of such body that you believe to be buried, the number of gyrations that you will obtain will effectively be the one that you will have obtained for this body during the establishment of your table of series; or the one, known to you, that you will have adopted from the identification table on page 96.

Indeed, your subconscious has received, during the first experiments on each body or radiation, a kind of definitive education, and you will have instructed it once and for all to make your pendulum react in the same way (polarity, series, radius fundamental) in the presence of each of these bodies or radiations.

The subjective fundamental ray : Studied on page 1 13, **it** was discovered by Abbé Mermet. It is emitted by any body or radiation in a fixed direction, which is proper to it. Upon meeting it, the pendulum oscillates in this fixed direction, normally at the end of a series. .

The mental witness : It is obviously very convenient. when looking for a body. to concentrate one's desire to be receptive to the radiations of this body on a specimen of this body which one has before one's eyes and which one can, as a "search factor", place between the thumb and the index, attached to the rod of his pendulum.

But it is even more convenient, especially if one does not have an identical body at hand, to "represent" it strongly in the mind and in a precise way. Such is the 'mental witness' .

The images : Few are the dowzers who are not confronted with the multiple images which move around a body. These images sometimes direct research towards one or more false positions of the real body. Sometimes the image still appears in the form of an afterglow, a sort of immaterial trace of the body which has stayed in this place previously. Finally, the mass of a body can impregnate all the ground which surrounds it with its own waves; the pendulum then reacts far from the body on all the surroundings, sometimes with a greater intensity than on the body itself.

It is therefore necessary to eliminate these various images, the causes of so many failures and errors. The systematic use of the Schumfell pendulum renders the most valuable services in this respect.

But some images can become very useful. This is the case of the image-impregnation for the search, for example, of the body of a drowned person, of which we will use the body linen used by him, to capture the personal waves.

nal. This will also be the case for the study of a writing which has received the image-impregnation, by the hand of the writer, of the latter's personal waves, waves on which we can question the subconscious.

The impregnation, studied on page 208, whether on paper, on clothing, in the ground, on liquid, or in the air, is indeed an image. It can be compared to a vibration given by a bow to a tuning fork, and maintained. This phenomenon is completely different from that of afterglow with which it is sometimes confused.

The remanence, studied on page 207, has a deceptive influence on the black neutral spherical pendulum. It takes a long practice to arrive, with this pendulum, to distinguish the radiation of a real body, from that of one of the images to which this radiation has given rise. The remanence consists in the artificial impression left in the medium where a body has stayed, to the point of producing on the black spherical neutral pendulum, the same reactions as the real body itself.

The Schumfell electro-magnetic pendulum, patented by SGD, was precisely designed for the elimination of images of all categories, including remanence and impregnation. Its use relieves the operator of the concern arising from this delicate question, which, out of his mind, one comes up against at all times.

*..
:i: *

Such are the processes of the mental method.

We have endeavored to summarize, as clearly as possible, the particular processes of this method, so as not to embarrass the beginner, while instructing him sufficiently so that he then progresses, on his own, quickly.

We will be especially happy if we succeed, after having developed the sensitivity of many readers, thanks to the progressive exercises of flexibility of the Physical Method, to lead them to the practice, more comfortable and superior, of the Mental Method, which will allow them would most often have been discouraging and impractical if they had embarked on it without this judicious preparation.

Let us note, indeed, that innumerable are the admirers of the dowsing; but they remained mere amateur ' pendulants ', because their initial momentum was shattered by inevitable setbacks. Legion will be, on the other hand, from now on, the true dowsers; those will have committed themselves to a methodical physical preparation, the results of which will

already be considerable. But they will then be able to benefit, with faith and success, from the ^Aprodigious possibilities of Thought.

Register of Trades: Nice 51 20
Register of Commwrcc : Nice 50.652
Printed in France.

B - ----- - ----- -û1

Printing
of • L'Eclaireur d-.1 Nico •
21, Ruo Mt'yorbccr, 21

- ----- -

Deposit Il-gal 2B<*-&.-1 No. 38 ..

1
1
1
1