

# 01204211 Discrete Mathematics

## Lecture 6: Mathematical Induction 1

Jittat Fakcharoenphol

September 1, 2015

# Mathematical Induction

- ▶ In this lecture, we will focus on how to prove properties on natural numbers.

# Mathematical Induction

- ▶ In this lecture, we will focus on how to prove properties on natural numbers.
- ▶ For example, we may want to prove that for any integer  $n \geq 1$ ,

$$\sum_{i=1}^n i = i(i+1)/2,$$

# Mathematical Induction

- ▶ In this lecture, we will focus on how to prove properties on natural numbers.
- ▶ For example, we may want to prove that for any integer  $n \geq 1$ ,

$$\sum_{i=1}^n i = i(i+1)/2,$$

or for any integer  $n \geq 1$ ,

$$\sum_{i=1}^n i^2 = \frac{n}{6}(n+1)(2n+1),$$

# Mathematical Induction

- ▶ In this lecture, we will focus on how to prove properties on natural numbers.
- ▶ For example, we may want to prove that for any integer  $n \geq 1$ ,

$$\sum_{i=1}^n i = i(i+1)/2,$$

or for any integer  $n \geq 1$ ,

$$\sum_{i=1}^n i^2 = \frac{n}{6}(n+1)(2n+1),$$

or “We can pay any integer amount  $x \geq 4$  baht with 2-baht coins and 5-baht coins.”

## A review of the summation notation (by examples)

$$\blacktriangleright \sum_{i=1}^{10} i =$$

## A review of the summation notation (by examples)

►  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)



# A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) =$

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$   
(reads “sum from  $i = 7$  to 9 of  $i^2 + i$ ” or “sum of  $i^2 + i$  from  $i = 7$  to 9”)

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$   
(reads “sum from  $i = 7$  to 9 of  $i^2 + i$ ” or “sum of  $i^2 + i$  from  $i = 7$  to 9”)
- ▶ The range of the index may be sets. For example, let  $A = \{1, 2, 4, 15\}$ , we have that  $\sum_{i \in A} i^2 =$

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$   
(reads “sum from  $i = 7$  to 9 of  $i^2 + i$ ” or “sum of  $i^2 + i$  from  $i = 7$  to 9”)
- ▶ The range of the index may be sets. For example, let  $A = \{1, 2, 4, 15\}$ , we have that  $\sum_{i \in A} i^2 = 1^2 + 2^2 + 4^2 + 15^2.$

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$   
(reads “sum from  $i = 7$  to 9 of  $i^2 + i$ ” or “sum of  $i^2 + i$  from  $i = 7$  to 9”)
- ▶ The range of the index may be sets. For example, let  $A = \{1, 2, 4, 15\}$ , we have that  $\sum_{i \in A} i^2 = 1^2 + 2^2 + 4^2 + 15^2.$
- ▶ What is  $\sum_{i=5}^2 i$ ?

## A review of the summation notation (by examples)

- ▶  $\sum_{i=1}^{10} i = 1 + 2 + \cdots + 10.$   
(reads “sum from  $i = 1$  to 10 of  $i$ ” or “sum of  $i$  from  $i = 1$  to 10”)
- ▶  $\sum_{i=7}^9 (i^2 + i) = (7^2 + 7) + (8^2 + 8) + (9^2 + 9).$   
(reads “sum from  $i = 7$  to 9 of  $i^2 + i$ ” or “sum of  $i^2 + i$  from  $i = 7$  to 9”)
- ▶ The range of the index may be sets. For example, let  $A = \{1, 2, 4, 15\}$ , we have that  $\sum_{i \in A} i^2 = 1^2 + 2^2 + 4^2 + 15^2.$
- ▶ What is  $\sum_{i=5}^2 i$ ? Note that in this case, the range is empty. This sum is called an **empty sum**. By convention, we define it to be zero.

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ :



# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1,

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ ,

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ :

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ ,

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ ,

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ , OK
- ▶ Try  $n = 3$ :

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ , OK
- ▶ Try  $n = 3$ : LHS:  $1 + 2 + 3 = 6$ ,

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ , OK
- ▶ Try  $n = 3$ : LHS:  $1 + 2 + 3 = 6$ , RHS:  $3(3+1)/2 = 6$ ,



# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ , OK
- ▶ Try  $n = 3$ : LHS:  $1 + 2 + 3 = 6$ , RHS:  $3(3+1)/2 = 6$ , OK
- ▶ Try ...

# Informal arguments

- ▶ Let's try to check that  $\sum_{i=1}^n i = i(i+1)/2$ , for any integer  $n \geq 1$ , by experimentation.
- ▶ Try  $n = 1$ : LHS: 1, RHS:  $1(1+1)/2 = 1$ , OK
- ▶ Try  $n = 2$ : LHS:  $1 + 2 = 3$ , RHS:  $2(2+1)/2 = 3$ , OK
- ▶ Try  $n = 3$ : LHS:  $1 + 2 + 3 = 6$ , RHS:  $3(3+1)/2 = 6$ , OK
- ▶ Try ...
- ▶ With this approach, we can't actually prove this statement.