01204211 Discrete Mathematics Lecture 13: Binomial Coefficients

Jittat Fakcharoenphol

September 20, 2015

The binomial coefficients

There is a reason why the term $\binom{n}{k}$ is called the binomial coefficients. In this lecture, we will discuss

- the Pascal's triangle,
- the binomial theorem, and
- advanced counting with binomial coefficients.

The equation

Last time we proved that, for n, k > 0,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

While we can prove this equation algebraically using definitions of binomial coefficients, proving the fact by describing the process of choosing k-subsets reveals interesting insights. This equation also hints us how to compute the value of $\binom{n}{k}$ using values of $\binom{n}{\cdot}$'s. So, let's try to do it.

The table

We shall use the fact that $\binom{n}{0}=1$ and $\binom{n}{k}=\binom{n-1}{k-1}+\binom{n-1}{k}$ to fill in the following table.

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

You can note that the table is left-right symmetric. This is true because of the fact that $\binom{n}{k} = \binom{n}{n-k}$.

The Triangle

If we move the numbers in the table slightly to the right, the table becomes the Pascal's triangle.

```
6
     10
                10
15
          20
                     15
```

The table and the binomial coefficients have many other interesting properties.