# 01204211 Discrete Mathematics Lecture 13: Binomial Coefficients

Jittat Fakcharoenphol

September 21, 2015

#### The binomial coefficients<sup>1</sup>

There is a reason why the term  $\binom{n}{k}$  is called the binomial coefficients. In this lecture, we will discuss

- the Pascal's triangle,
- the binomial theorem, and
- advanced counting with binomial coefficients.

¹This lecture mostly follows Chapter 3 of [LPV]. ←□ → ←② → ←② → ←② → → ② → ○② ←

## The equation

Last time we proved that, for n, k > 0,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

## The equation

Last time we proved that, for n, k > 0,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

While we can prove this equation algebraically using definitions of binomial coefficients, proving the fact by describing the process of choosing k-subsets reveals interesting insights. This equation also hints us how to compute the value of  $\binom{n}{k}$  using values of  $\binom{n}{\cdot}$ 's.

### The equation

Last time we proved that, for n, k > 0,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

While we can prove this equation algebraically using definitions of binomial coefficients, proving the fact by describing the process of choosing k-subsets reveals interesting insights. This equation also hints us how to compute the value of  $\binom{n}{k}$  using values of  $\binom{n}{\cdot}$ 's. So, let's try to do it.

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1						

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				

$\overline{n}$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1						

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1						

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1						

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1						

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

We shall use the fact that  $\binom{n}{0}=1$  and  $\binom{n}{k}=\binom{n-1}{k-1}+\binom{n-1}{k}$  to fill in the following table.

n	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

You can note that the table is left-right symmetric. This is true because of the fact that  $\binom{n}{k} = \binom{n}{n-k}$ .

## The Triangle

If we move the numbers in the table slightly to the right, the table becomes the Pascal's triangle.

# The Triangle

If we move the numbers in the table slightly to the right, the table becomes the Pascal's triangle.

```
10
                10
15
          20
                     15
```

The table and the binomial coefficients have many other interesting properties.

- $(x+y)^1 = x+y$
- $(x+y)^2 =$

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$
- $(x+y)^3 =$

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$
- $(x+y)^3 = x^3 + 3 \cdot x^2 y + 3 \cdot xy^2 + y^3$

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$
- $(x+y)^3 = x^3 + 3 \cdot x^2y + 3 \cdot xy^2 + y^3$
- $(x+y)^4 =$

Let's start by looking at polynomial of the form  $(x+y)^n$ . Let's start with small values of n:

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$
- $(x+y)^3 = x^3 + 3 \cdot x^2y + 3 \cdot xy^2 + y^3$
- $(x+y)^4 = x^4 + 4 \cdot x^3y + 6 \cdot x^2y^2 + 4 \cdot xy^3 + y^4.$

Let's focus on the coefficient of each term. You may notice that terms  $x^n$  and  $y^n$  always have 1 as their coefficients. Why is that?

Let's start by looking at polynomial of the form  $(x+y)^n$ . Let's start with small values of n:

- $(x+y)^1 = x+y$
- $(x+y)^2 = x^2 + 2 \cdot xy + y^2$
- $(x+y)^3 = x^3 + 3 \cdot x^2y + 3 \cdot xy^2 + y^3$
- $(x+y)^4 = x^4 + 4 \cdot x^3y + 6 \cdot x^2y^2 + 4 \cdot xy^3 + y^4.$

Let's focus on the coefficient of each term. You may notice that terms  $x^n$  and  $y^n$  always have 1 as their coefficients. Why is that? Let's look further at the coefficients of terms  $x^{n-1}y$ . Do you see any pattern in their coefficients? Can you explain why?

Let's take a look at  $(x+y)^4$  again. It is

$$(x+y)(x+y)(x+y)(x+y).$$

▶ How do we get  $x^4$  in the expansion?

Let's take a look at  $(x+y)^4$  again. It is

$$(x+y)(x+y)(x+y)(x+y).$$

- ▶ How do we get  $x^4$  in the expansion? For every factory, you have to pick x.
- ▶ How do we get  $xy^3$  in the expansion?

Let's take a look at  $(x+y)^4$  again. It is

$$(x+y)(x+y)(x+y)(x+y).$$

- ▶ How do we get  $x^4$  in the expansion? For every factory, you have to pick x.
- ▶ How do we get  $xy^3$  in the expansion? Out of the 4 factors, you have to pick y in one of the factor (or you have to pick x in 3 of the factors).

Let's take a look at  $(x+y)^4$  again. It is

$$(x+y)(x+y)(x+y)(x+y).$$

- ▶ How do we get  $x^4$  in the expansion? For every factory, you have to pick x.
- ▶ How do we get  $xy^3$  in the expansion? Out of the 4 factors, you have to pick y in one of the factor (or you have to pick x in 3 of the factors). Thus there are  $\binom{4}{3} = \binom{4}{1}$  ways to do so.

#### The binomial theorem

Theorem: If you expand  $(x+y)^n$ , the coefficient of the term  $x^ky^{n-k}$  is  $\binom{n}{k}$ .

That is,

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} =$$

$$\binom{n}{n} x^n + \binom{n}{n-1} x^{n-1} y^1 + \binom{n}{n-2} x^{n-2} y^2 + \dots + \binom{n}{1} x y^{n-1} + \binom{n}{0} y^n.$$

## Additional applications of the binomial theorem

The binomial theorem can be used to prove various identities regarding the binomial coefficients. For example, if we let x=1 and y=1, we get that

$$(1+1)^n = 2^n = \binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n-1} + \binom{n}{n}.$$

## Additional applications of the binomial theorem

The binomial theorem can be used to prove various identities regarding the binomial coefficients. For example, if we let x=1 and y=1, we get that

$$(1+1)^n = 2^n = \binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n-1} + \binom{n}{n}.$$

Quick check. Can you prove that

$$\binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots = 0.$$

Note that this statements says that the number of odd subsets equals the number of even subsets.

# More on counting

We shall see more techniques for counting when we consider the following problems.

- How many anagrams does the word "KASETSARTUNIVERSITY" have? (They do not have to be real English words.)
- ▶ How can you give out *n* different presents to *k* students when student *i* has to get *n<sub>i</sub>* pieces of presents?
- ▶ How many ways can you distribute n baht coins to k children?

► An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of ADD are ADD, DAD, and DDA.

- ▶ An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of *ADD* are *ADD*, *DAD*, and *DDA*.
- ► How many anagrams does "ABC" have?

- ▶ An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of *ADD* are *ADD*, *DAD*, and *DDA*.
- ► How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.

- ▶ An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of *ADD* are *ADD*, *DAD*, and *DDA*.
- ▶ How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.
- ▶ How many anagrams does "ABCC" have?

- ► An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of ADD are ADD, DAD, and DDA.
- ▶ How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.
- ▶ How many anagrams does "ABCC" have?
  - ► This time we have to be careful because the answer of 4! is too large as it over counts many anagrams.

- ► An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of ADD are ADD, DAD, and DDA.
- ▶ How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.
- ▶ How many anagrams does "ABCC" have?
  - ► This time we have to be careful because the answer of 4! is too large as it over counts many anagrams.
  - ▶ How many times does "CABC" get counted in 4!?

- ▶ An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of *ADD* are *ADD*, *DAD*, and *DDA*.
- ▶ How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.
- ▶ How many anagrams does "ABCC" have?
  - ▶ This time we have to be careful because the answer of 4! is too large as it over counts many anagrams.
  - ▶ How many times does "CABC" get counted in 4!?
  - If we treat two C's differently as  $C_1$  and  $C_2$ , we can see that CABC is counted twice as  $C_1ABC_2$  and  $C_2ABC_1$ . This is true for any anagram of ABCC.

- ▶ An anagram of a particular word is a word that uses the same set of alphabets. For example, the anagrams of *ADD* are *ADD*, *DAD*, and *DDA*.
- ► How many anagrams does "ABC" have?
  - ▶ 3!, because every permutation of A B or C is a different anagram.
- ▶ How many anagrams does "ABCC" have?
  - ▶ This time we have to be careful because the answer of 4! is too large as it over counts many anagrams.
  - ▶ How many times does "CABC" get counted in 4!?
  - ▶ If we treat two C's differently as  $C_1$  and  $C_2$ , we can see that CABC is counted twice as  $C_1ABC_2$  and  $C_2ABC_1$ . This is true for any anagram of ABCC.
  - Since each anagram is counted in 4! twice, the number of anagrams is  $4!/2 = 4 \cdot 3 = 12$ .

## General anagrams

Let's try to use the same approach to count the anagram of HELLOWORLD. (It has 3 L's, 2 O's, H, E, W, R, and D.)

### General anagrams

Let's try to use the same approach to count the anagram of HELLOWORLD. (It has 3 L's, 2 O's, H, E, W, R, and D.)

The number of permutation of alphabets in HELLOWORLD, treating each character differently is 10!. However, each anagram is counted for 3!2! times because of the 3 copies of L and the 2 copies of O. Therefore, the number of anagrams is

$$\frac{10!}{3!2!}$$

### Distributing presents

I have 9 different presents. I want to give them to 3 students: A, B, and C. I want to give each student 3 presents. In how many ways can I do it?

## Distributing presents

I have 9 different presents. I want to give them to 3 students: A, B, and C. I want to give each student 3 presents. In how many ways can I do it?

This is essentially an anagram problem. You can think of one particular way of present distribution as anagram of AAABBCCC. Thus, the number is

$$\frac{9!}{3!3!3!}$$