AFRICAST

TIDY TIME SERIES & FORECASTING IN R

Exponential smoothing

Session 7 of 10



SECTION 1 OF 5

Exponential smoothing







1 **Exponential smoothing** 2 Trend methods 3 Seasonal methods 4 ETS taxonomy 5 Non-Gaussian forecast distributions



From simple methods to Exponential Smoothing

- Naive method: Use only the last observation.
- Average method: Use all observations.
- Want something in between naive and average methods.
- Most recent data should have more weight.
- This is exactly the concept behind exponential smoothing.

Pegels' classification

Trend	Seasonality		
	None	Additive	Multiplicative
None			
Additive			
Additive Damped			
Multiplicative			
Multiplicative Damped			

Historical perspective

- Developed in the 1950s and 1960s as methods (algorithms) to produce point forecasts.
- Combine a “level”, “trend” (slope) and “seasonal” component to describe a time series.
- The rate of change of the components are controlled by “smoothing parameters”: α, β and γ respectively.
- Need to choose best values for the smoothing parameters (and initial states).
- Equivalent ETS state space models developed in the 1990s and 2000s.

A model for levels, trends, and seasonalities

We want a model that captures the level (ℓ_t), trend (b_t) and seasonality (s_t).

How do we combine these elements?

Additively?

$$y_t = \ell_{t-1} + b_{t-1} + s_{t-m} + \varepsilon_t$$

Multiplicatively?

$$y_t = \ell_{t-1} b_{t-1} s_{t-m} (1 + \varepsilon_t)$$


Perhaps a mix of both?

$$y_t = (\ell_{t-1} + b_{t-1}) s_{t-m} + \varepsilon_t$$

How do the level, trend and seasonal components evolve over time?

ETS models

General notation E T S : **E**xpon**e**n**T**ial **S**moothing



Error: Additive (`"A"`) or multiplicative (`"M"`)

Trend: None (`"N"`), additive (`"A"`), multiplicative (`"M"`), or damped (`"Ad"` or `"Md"`).

Seasonality: None (`"N"`), additive (`"A"`) or multiplicative (`"M"`)

ETS(A,N,N): SES with additive errors

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T \\
\text{Measurement equation} \quad & y_t = \ell_{t-1} + \varepsilon_t \\
\text{State equation} \quad & \ell_t = \ell_{t-1} + \alpha\varepsilon_t
\end{aligned}
$$

where $\varepsilon_t \sim \text{NID}(0, \sigma^2)$.

- "innovations" or "single source of error" because equations have the same error process, ε_t.
- Measurement equation: relationship between observations and states.
- Transition/state equation(s): evolution of state(s) over time.

ETS(M,N,N): SES with multiplicative errors

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T \\
\text{Measurement equation} \quad & y_t = \ell_{t-1}(1+\varepsilon_t) \\
\text{State equation} \quad & \ell_t = \ell_{t-1}(1+\alpha\varepsilon_t)
\end{aligned}
$$

where $\varepsilon_t \sim \text{NID}(0, \sigma^2)$.

- Models with additive and multiplicative errors with the same parameters generate the same point forecasts but different prediction intervals.

SECTION 2 OF 5

Trend methods

1 Exponential smoothing
2 **Trend methods**
3 Seasonal methods
4 ETS taxonomy
5 Non-Gaussian forecast distributions



Holt's linear trend

Additive errors: ETS(A,A,N)

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T + hb_T \\
\text{Measurement equation} \quad & y_t = \ell_{t-1} + b_{t-1} + \varepsilon_t \\
\text{State equations} \quad & \ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t \\
& b_t = b_{t-1} + \beta\varepsilon_t
\end{aligned}
$$

Multiplicative errors: ETS(M,A,N)

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T + hb_T \\
\text{Measurement equation} \quad & y_t = (\ell_{t-1} + b_{t-1})(1 + \varepsilon_t) \\
\text{State equations} \quad & \ell_t = (\ell_{t-1} + b_{t-1})(1 + \alpha\varepsilon_t) \\
& b_t = b_{t-1} + \beta(\ell_{t-1} + b_{t-1})\varepsilon_t
\end{aligned}
$$

Example: Australian population

```
aus_economy <- global_economy |>
  filter(Country == "Australia") |>
  mutate(Pop = Population / 1e6)
fit <- aus_economy |>
  model(AAN = ETS(Pop ~ error("A") + trend("A") + season("N")))
report(fit)
```

```
Series: Pop
Model: ETS(A,A,N)
  Smoothing parameters:
    alpha = 1
    beta  = 0.327

  Initial states:
 l[0]  b[0]
 10.1 0.222

  sigma^2:  0.0041

  AIC  AICc   BIC
-77.0 -75.8 -66.7
```


Example: Australian population

```
components(fit)
```

```
# A dable: 59 x 7 [1Y]
# Key:     Country, .model [1]
# :        Pop = lag(level, 1) + lag(slope, 1) + remainder
   Country   .model  Year   Pop level slope remainder
   <fct>     <chr>  <dbl> <dbl> <dbl> <dbl>     <dbl>
 1 Australia AAN     1959  NA    10.1 0.222 NA
 2 Australia AAN     1960  10.3  10.3 0.222 -0.000145
 3 Australia AAN     1961  10.5  10.5 0.217 -0.0159
 4 Australia AAN     1962  10.7  10.7 0.231  0.0418
 5 Australia AAN     1963  11.0  11.0 0.223 -0.0229
 6 Australia AAN     1964  11.2  11.2 0.221 -0.00641
 7 Australia AAN     1965  11.4  11.4 0.221 -0.000314
 8 Australia AAN     1966  11.7  11.7 0.235  0.0418
 9 Australia AAN     1967  11.8  11.8 0.206 -0.0869
10 Australia AAN     1968  12.0  12.0 0.208  0.00350
# i 49 more rows
```


Example: Australian population

```
components(fit) |> autoplot()
```



Example: Australian population

```
fit |>
  forecast(h = 20) |>
  autoplot(aus_economy) +
  labs(y = "Population", x = "Year")
```



ETS(A,Ad,N): Damped trend method


Additive errors

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T + (\phi + \phi^2 + \cdots + \phi^h) b_T \\
\text{Measurement equation} \quad & y_t = (\ell_{t-1} + \phi b_{t-1}) + \varepsilon_t \\
\text{State equations} \quad & \ell_t = (\ell_{t-1} + \phi b_{t-1}) + \alpha \varepsilon_t \\
& b_t = \phi b_{t-1} + \beta \varepsilon_t
\end{aligned}
$$

- Damping parameter $0 < \phi \leq 1$.
- If $\phi = 1$, identical to Holt's linear trend.
- As $h \to \infty$, $\hat{y}_{T+h|T} \to \ell_T + \phi b_T / (1 - \phi)$.
- Short-run forecasts trended, long-run forecasts constant.

Example: Australian population

```
aus_economy |>
  model(damped = ETS(Pop ~ error("A") + trend("Ad") + season("N"))) |>
  forecast(h = 20) |>
  autoplot(aus_economy)
```



Example: Nigerian population

```
global_economy |>
  filter(Country == "Nigeria") |>
  mutate(Pop = Population / 1e6) |>
  model(
    ANN = ETS(Pop ~ error("A") + trend("N") + season("N")),
    AAN = ETS(Pop ~ error("A") + trend("A") + season("N")),
    AAdN = ETS(Pop ~ error("A") + trend("Ad") + season("N"))
  ) |>
  glance() |>
  select(.model, AICc)
```

```
# A tibble: 3 × 2
  .model  AICc
  <chr>  <dbl>
1 ANN     360.
2 AAN     -37.7
3 AAdN     26.9
```

Which ETS model is best for Nigeria's population?

ETS(A,N,N) **0** 0%

ETS(A,A,N) **0** 0%

ETS(A,Ad,N) **0** 0%

poll.nectric.com.au/s/africast-ets

0 responses

● CLOSED 0 connected

Example: National populations

```
fit <- global_economy |>
  mutate(Pop = Population / 1e6) |>
  model(ets = ETS(Pop))
fit
```

```
# A mable: 263 x 2
# Key:     Country [263]
   Country                     ets
   <fct>                   <model>
 1 Afghanistan          <ETS(A,A,N)>
 2 Albania              <ETS(M,A,N)>
 3 Algeria              <ETS(M,A,N)>
 4 American Samoa       <ETS(M,A,N)>
 5 Andorra              <ETS(M,A,N)>
 6 Angola               <ETS(M,A,N)>
 7 Antigua and Barbuda  <ETS(M,A,N)>
 8 Arab World           <ETS(M,A,N)>
 9 Argentina            <ETS(A,A,N)>
10 Armenia              <ETS(M,A,N)>
# i 253 more rows
```

Example: National populations

```
fit |>
  forecast(h = 5)
```

```
# A fable: 1,315 x 5 [1Y]
# Key:     Country, .model [263]
   Country     .model  Year
   <fct>       <chr>  <dbl>
 1 Afghanistan ets     2018
 2 Afghanistan ets     2019
 3 Afghanistan ets     2020
 4 Afghanistan ets     2021
 5 Afghanistan ets     2022
 6 Albania     ets     2018
 7 Albania     ets     2019
 8 Albania     ets     2020
 9 Albania     ets     2021
10 Albania     ets     2022
# i 1,305 more rows
# i 2 more variables: Pop <dist>, .mean <dbl>
```

SECTION 3 OF 5

Seasonal methods

1 Exponential smoothing
2 Trend methods
3 **Seasonal methods**
4 ETS taxonomy
5 Non-Gaussian forecast distributions



ETS(A,A,A): Holt-Winters additive method

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = \ell_T + hb_T + s_{T+h-m(k+1)} \\
\text{Measurement equation} \quad & y_t = \ell_{t-1} + b_{t-1} + s_{t-m} + \varepsilon_t \\
\text{State equations} \quad & \ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t \\
& b_t = b_{t-1} + \beta\varepsilon_t \\
& s_t = s_{t-m} + \gamma\varepsilon_t
\end{aligned}
$$

- $k =$ integer part of $(h-1)/m$.
- $\sum_i s_i \approx 0$.
- Parameters: $0 \leq \alpha \leq 1$, $0 \leq \beta \leq \alpha$, $0 \leq \gamma \leq 1-\alpha$ and $m =$ period of seasonality (e.g. $m = 4$ for quarterly data).

ETS(M,A,M): Holt-Winters multiplicative method

$$
\begin{aligned}
\text{Forecast equation} \quad & \hat{y}_{T+h|T} = (\ell_T + hb_T)s_{T+h-m(k+1)} \\
\text{Measurement equation} \quad & y_t = (\ell_{t-1} + b_{t-1})s_{t-m}(1+\varepsilon_t) \\
\text{State equations} \quad & \ell_t = (\ell_{t-1} + b_{t-1})(1+\alpha\varepsilon_t) \\
& b_t = b_{t-1} + \beta(\ell_{t-1} + b_{t-1})\varepsilon_t \\
& s_t = s_{t-m}(1+\gamma\varepsilon_t)
\end{aligned}
$$


- k is integer part of $(h-1)/m$.
- $\sum_i s_i \approx m$.
- Parameters: $0 \leq \alpha \leq 1$, $0 \leq \beta \leq \alpha$, $0 \leq \gamma \leq 1-\alpha$ and $m =$ period of seasonality (e.g. $m = 4$ for quarterly data).

Example: Australian holiday tourism

```
holidays <- tourism |>
  filter(Purpose == "Holiday")
fit <- holidays |> model(ets = ETS(Trips))
fit
```

```
# A mable: 76 x 4
# Key:     Region, State, Purpose [76]
   Region          State Purpose          ets
   <chr>           <chr> <chr>        <model>
 1 Adelaide        SA    Holiday <ETS(A,N,A)>
 2 Adelaide Hills  SA    Holiday <ETS(A,A,N)>
 3 Alice Springs   NT    Holiday <ETS(M,N,A)>
 4 Ballarat        VIC   Holiday <ETS(M,N,A)>
 5 Barkly          NT    Holiday <ETS(A,N,A)>
 6 Barossa         SA    Holiday <ETS(A,N,N)>
 7 Bendigo Loddon  VIC   Holiday <ETS(M,N,N)>
 8 Blue Mountains  NSW   Holiday <ETS(M,N,M)>
 9 Brisbane        QLD   Holiday <ETS(A,A,N)>
10 Bundaberg       QLD   Holiday <ETS(A,N,A)>
# i 66 more rows
```

Example: Australian holiday tourism



Which ETS model suits Snowy Mountains holiday trips?

ETS(?,A,A)	0	0%
ETS(?,N,A)	0	0%
ETS(?,N,M)	0	0%
ETS(?,N,N)	0	0%

poll.nectric.com.au/s/africast-ets

0 responses

● CLOSED 0 connected

Example: Australian holiday tourism

```
fit |>
  filter(Region == "Snowy Mountains") |>
  report()
```

```
Series: Trips
Model: ETS(M,N,A)
  Smoothing parameters:
    alpha = 0.157
    gamma = 1e-04

  Initial states:
 l[0] s[0] s[-1] s[-2] s[-3]
  142  -61   131 -42.2 -27.7

  sigma^2:  0.0388

 AIC AICc  BIC
 852  854  869
```


Example: Australian holiday tourism

```
fit |>
  filter(Region == "Snowy Mountains") |>
  components()
```

```
# A dable: 84 x 9 [1Q]
# Key:     Region, State, Purpose, .model [1]
# :        Trips = (lag(level, 1) + lag(season, 4)) * (1 + remainder)
   Region           State Purpose .model Quarter Trips level season remainder
   <chr>            <chr> <chr>   <chr>    <qtr> <dbl> <dbl>  <dbl>     <dbl>
 1 Snowy Mountains NSW   Holiday ets    1997 Q1   NA     NA   -27.7    NA
 2 Snowy Mountains NSW   Holiday ets    1997 Q2   NA     NA   -42.2    NA
 3 Snowy Mountains NSW   Holiday ets    1997 Q3   NA     NA   131.     NA
 4 Snowy Mountains NSW   Holiday ets    1997 Q4   NA    142.  -61.0    NA
 5 Snowy Mountains NSW   Holiday ets    1998 Q1  101.   140.  -27.7    -0.113
 6 Snowy Mountains NSW   Holiday ets    1998 Q2  112.   142.  -42.2     0.154
 7 Snowy Mountains NSW   Holiday ets    1998 Q3  310.   148.  131.      0.137
 8 Snowy Mountains NSW   Holiday ets    1998 Q4   89.8  148.  -61.0     0.0335
 9 Snowy Mountains NSW   Holiday ets    1999 Q1  112.   147.  -27.7    -0.0687
10 Snowy Mountains NSW   Holiday ets    1999 Q2  103.   147.  -42.2    -0.0199
# i 74 more rows
```

Example: Australian holiday tourism

```
fit |>
  filter(Region == "Snowy Mountains") |>
  components() |>
  autoplot()
```




Example: Australian holiday tourism

```
fit |> forecast()
```

```
# A fable: 608 x 7 [1Q]
# Key:     Region, State, Purpose, .model [76]
   Region          State Purpose .model Quarter
   <chr>           <chr> <chr>   <chr>    <qtr>
 1 Adelaide        SA    Holiday ets    2018 Q1
 2 Adelaide        SA    Holiday ets    2018 Q2
 3 Adelaide        SA    Holiday ets    2018 Q3
 4 Adelaide        SA    Holiday ets    2018 Q4
 5 Adelaide        SA    Holiday ets    2019 Q1
 6 Adelaide        SA    Holiday ets    2019 Q2
 7 Adelaide        SA    Holiday ets    2019 Q3
 8 Adelaide        SA    Holiday ets    2019 Q4
 9 Adelaide Hills SA    Holiday ets    2018 Q1
10 Adelaide Hills SA    Holiday ets    2018 Q2
# i 598 more rows
# i 2 more variables: Trips <dist>, .mean <dbl>
```

Example: Australian holiday tourism

```
fit |>
  forecast() |>
  filter(Region == "Snowy Mountains") |>
  autoplot(holidays) +
  labs(x = "Quarter", y = "Overnight trips (thousands)")
```



SECTION 4 OF 5

ETS taxonomy

1 Exponential smoothing
2 Trend methods
3 Seasonal methods
4 **ETS taxonomy**
5 Non-Gaussian forecast distributions



Exponential smoothing models

Additive Error		N	A	M
Trend Component		**(None)**	**(Additive)**	**(Multiplicative)**
N	(None)	A,N,N	A,N,A	~~A,N,M~~
A	(Additive)	A,A,N	A,A,A	~~A,A,M~~
A_d	(Additive damped)	A,A_d,N	A,A_d,A	~~A,A_d,M~~

Multiplicative Error		N	A	M
Trend Component		**(None)**	**(Additive)**	**(Multiplicative)**
N	(None)	M,N,N	M,N,A	M,N,M
A	(Additive)	M,A,N	M,A,A	M,A,M
A_d	(Additive damped)	M,A_d,N	M,A_d,A	M,A_d,M

Estimating ETS models

- Smoothing parameters α, β, γ and ϕ, and the initial states ℓ_0, b_0, $s_0, s_{-1}, \ldots, s_{-m+1}$ are estimated by maximising the "likelihood" = the probability of the data arising from the specified model.
- For models with additive errors equivalent to minimising SSE.
- For models with multiplicative errors, **not** equivalent to minimising SSE.

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2\log(L) + 2k$$

where L is the likelihood and k is the number of parameters and initial states estimated in the model.

Corrected AIC

$$\text{AIC}_\text{c} = \text{AIC} + \frac{2k(k+1)}{T-k-1}$$

which is the AIC corrected (for small sample bias).

Bayesian Information Criterion

$$\text{BIC} = \text{AIC} + k(\log(T) - 2).$$

AIC and cross-validation

Minimising the AIC assuming Gaussian residuals is asymptotically equivalent to minimising one-step time series cross validation MSE.

Automatic forecasting

From Hyndman et al. (IJF, 2002):

1. Apply each model that is appropriate to the data. Optimise parameters and initial values using MLE.
2. Select best method using AICc.
3. Produce forecasts using best method.
4. Obtain forecast intervals using underlying state space model.

- Method performed very well in M3 competition.
- Used as a benchmark in the M4 competition.

SECTION 5 OF 5


Non-Gaussian forecast distributions

1 Exponential smoothing
2 Trend methods
3 Seasonal methods
4 ETS taxonomy
5 **Non-Gaussian forecast distributions**




Non-Gaussian forecast distributions

```
vic_cafe <- aus_retail |>
  filter(State == "Victoria",
         Industry == "Cafes, restaurants and catering services") |>
  select(Month, Turnover)
vic_cafe |>
  autoplot(Turnover) +
  labs(title = "Monthly turnover of Victorian cafes", y = "Turnover ($AUD million)")
```



Forecasting with transformations

```
vic_cafe |> autoplot(box_cox(Turnover, lambda = 0.2))
```




Forecasting with transformations

```
vic_cafe |>
  model(
    raw = ETS(Turnover),
    log = ETS(log(Turnover))
  ) |>
  glance() |>
  select(.model, AICc)
```

```
# A tibble: 2 × 2
  .model   AICc
  <chr>   <dbl>
1 raw     4613.
2 log       82.8
```

Which model is better for Victorian cafe turnover?



Forecasting with transformations

```
fit <- vic_cafe |>
  model(ets = ETS(box_cox(Turnover, 0.2)))
fit
```

```
# A mable: 1 x 1
           ets
       <model>
1 <ETS(A,A,A)>
```


```
(fc <- fit |> forecast(h = "3 years"))
```

```
# A fable: 36 x 4 [1M]
# Key:     .model [1]
   .model    Month        Turnover .mean
   <chr>     <mth>          <dist> <dbl>
 1 ets    2019 Jan  t(N(13, 0.02))  608.
 2 ets    2019 Feb t(N(13, 0.028))  563.
 3 ets    2019 Mar t(N(13, 0.036))  629.
 4 ets    2019 Apr t(N(13, 0.044))  615.
 5 ets    2019 May t(N(13, 0.052))  613.
 6 ets    2019 Jun t(N(13, 0.061))  593.
 7 ets    2019 Jul t(N(13, 0.069))  624.
 8 ets    2019 Aug t(N(13, 0.077))  640.
 9 ets    2019 Sep t(N(13, 0.085))  630.
10 ets    2019 Oct t(N(13, 0.093))  642.
# i 26 more rows
```

- `t(N)` denotes a transformed normal distribution.
- The distribution is back-transformed and the mean (`.mean`) is bias-adjusted automatically.

Forecasting with transformations

```
fc |> autoplot(vic_cafe)
```




Bootstrapped forecast distributions

```
sim <- fit |> generate(h = "3 years", times = 5, bootstrap = TRUE)
sim
```

```
# A tsibble: 180 x 4 [1M]
# Key:       .model, .rep [5]
   .model    Month .rep   .sim
   <chr>     <mth> <chr> <dbl>
 1 ets    2019 Jan 1      584.
 2 ets    2019 Feb 1      560.
 3 ets    2019 Mar 1      635.
 4 ets    2019 Apr 1      601.
 5 ets    2019 May 1      586.
 6 ets    2019 Jun 1      567.
 7 ets    2019 Jul 1      589.
 8 ets    2019 Aug 1      640.
 9 ets    2019 Sep 1      597.
10 ets    2019 Oct 1      651.
# i 170 more rows
```

Bootstrapped forecast distributions

```
vic_cafe |>
  filter(year(Month) >= 2008) |>
  ggplot(aes(x = Month)) +
  geom_line(aes(y = Turnover)) +
  geom_line(aes(y = .sim, colour = as.factor(.rep)), data = sim) +
  labs(title = "Monthly turnover of Victorian cafes") +
  guides(colour = "none")
```



Bootstrapped forecast distributions

```
fc <- fit |> forecast(h = "3 years", bootstrap = TRUE)
fc
```

```
# A fable: 36 x 4 [1M]
# Key:     .model [1]
   .model    Month     Turnover .mean
   <chr>     <mth>       <dist> <dbl>
 1 ets    2019 Jan sample[5000]  608.
 2 ets    2019 Feb sample[5000]  562.
 3 ets    2019 Mar sample[5000]  629.
 4 ets    2019 Apr sample[5000]  615.
 5 ets    2019 May sample[5000]  612.
 6 ets    2019 Jun sample[5000]  593.
 7 ets    2019 Jul sample[5000]  624.
 8 ets    2019 Aug sample[5000]  640.
 9 ets    2019 Sep sample[5000]  630.
10 ets    2019 Oct sample[5000]  642.
# i 26 more rows
```

Bootstrapped forecast distributions

```
fc |> autoplot(vic_cafe) +
  labs(title = "Monthly turnover of Victorian cafes")
```

