AFRICAST

TIDY TIME SERIES & FORECASTING IN R

Basic modelling and forecasting



Session 4 of 10

Outline

- Statistical forecasting
- What can we forecast?
- Benchmark methods
- Specify and estimate
- Produce forecasts
- Fitted values and residuals


SECTION 1 OF 6

Statistical forecasting

1 **Statistical forecasting** 2 What can we forecast? 3 Benchmark methods 4 Specify and estimate 5 Produce forecasts 6 Fitted values and residuals



Forecasting workflow




Statistical forecasting steps

- Prepare data.
- Visualise data.
- Choosing and fitting models (specify and train models).
- Produce forecast.
- Evaluate quality.


What is a forecast?

A forecast is an estimate of the probability distribution of a variable to be observed in the future.



What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What is a forecast?


A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model


Prediction interval

A forecast is an estimate of the probability distribution of a variable to be observed in the future.




Simulated futures from an ETS model

Visualising forecast distributions



Forecast distribution



Statistical forecasting

- Thing to be forecast: y_{T+h}.
- What we know: $y_1, \dots, y_T$.
- Forecast distribution: $y_{T+h|t} = y_{T+h} \mid \{y_1, y_2, \dots, y_T\}$.
- Point forecast: $\hat{y}_{T+h|T} = \mathrm{E}[y_{T+h} \mid y_1, \dots, y_T]$.
- Forecast variance: $\mathrm{Var}[y_t \mid y_1, \dots, y_T]$
- Prediction interval is a range of values of y_{T+h} with high probability.

SECTION 2 OF 6


What can we forecast?

1 Statistical forecasting
2 **What can we forecast?**
3 Benchmark methods
4 Specify and estimate
5 Produce forecasts
6 Fitted values and residuals



What can we forecast?

A forecast is an estimate of the probability distribution of a variable to be observed in the future.



Simulated futures from an ETS model

What can we forecast?



What can we forecast?




What can we forecast?



What can we forecast?



What can we forecast?



What can we forecast?



What can we forecast?



Which is easiest to forecast?

daily electricity demand in 3 days time	0	0%
timing of next Halley's comet appearance	0	0%
time of sunrise this day next year	0	0%
Google stock price tomorrow	0	0%
Google stock price in 6 months time	0	0%
maximum temperature tomorrow	0	0%
exchange rate of $US/AUS next week	0	0%
total sales of drugs in Australian pharmacies next month	0	0%



poll.nectric.com.au/s/africast-modelling

0 responses

CLOSED 0 connected

Which is hardest to forecast?

daily electricity demand in 3 days time	0	0%
timing of next Halley's comet appearance	0	0%
time of sunrise this day next year	0	0%
Google stock price tomorrow	0	0%
Google stock price in 6 months time	0	0%
maximum temperature tomorrow	0	0%
exchange rate of $US/AUS next week	0	0%
total sales of drugs in Australian pharmacies next month	0	0%



poll.nectric.com.au/s/africast-modelling

0 responses

CLOSED 0 connected

- how do we measure "easiest"?

Factors affecting forecastability

Something is easier to forecast if:

- we have a good understanding of the factors that contribute to it
- there is lots of data available;
- the forecasts cannot affect the thing we are trying to forecast.
- there is relatively low natural/unexplainable random variation.
- the future is somewhat similar to the past


SECTION 3 OF 6

Benchmark methods

1 Statistical forecasting
2 What can we forecast?
3 **Benchmark methods**
4 Specify and estimate
5 Produce forecasts
6 Fitted values and residuals




Some simple forecasting methods



How would you forecast these series?

Some simple forecasting methods



How would you forecast these series?


Some simple forecasting methods



How would you forecast these series?

`MEAN(y)`: Average method


- Forecast of all future values is equal to mean of historical data $\{y_1, \ldots, y_T\}$.
- Forecasts: $\hat{y}_{T+h|T} = \bar{y} = (y_1 + \cdots + y_T)/T$



Some simple forecasting methods


`NAIVE(y)`: Naïve method

- Forecasts equal to last observed value.
- Forecasts: $\hat{y}_{T+h|T} = y_T$.
- Consequence of efficient market hypothesis.



`SNAIVE(y ~ lag(m))`: Seasonal naïve method

- Forecasts equal to last value from same season.
- Forecasts: $\hat{y}_{T+h|T} = y_{T+h-m(k+1)}$, where $m =$ seasonal period and k is the integer part of $(h-1)/m$.




`RW(y ~ drift())`: Drift method

- Forecasts equal to last value plus average change.
- Forecasts:

$$\begin{aligned}\hat{y}_{T+h|T} &= y_T + \frac{h}{T-1}\sum_{t=2}^{T}(y_t - y_{t-1}) \\ &= y_T + \frac{h}{T-1}(y_T - y_1).\end{aligned}$$

- Equivalent to extrapolating a line drawn between first and last observations.

Drift method



SECTION 4 OF 6

Specify and estimate

1 Statistical forecasting
2 What can we forecast?
3 Benchmark methods
4 **Specify and estimate**
5 Produce forecasts
6 Fitted values and residuals



Model specification

- Model specification in fable supports a formula based interface
- A model formula in R is expressed using `response ~ terms`
  - the formula's left side describes the response
  - the right describes terms used to model the response.
- `Attention`: MODEL name is in capital letters, e.g. `SNAIVE`

Model estimation

The `model()` function trains models on data.

- It returns a `mable` object.

```
# Fit the models
my_mable <- my_data |>
  model(
    choose_name1 = MODEL1(response ~ term1+...),
    choose_name2 = MODEL2(response ~ term1+...),
    choose_name3 = MODEL3(response ~ term1+...),
    choose_name4 = MODEL4(response ~ term1+...)
  )
```

Model fitting: example

The `model()` function trains models on data.

```
beer_fit <- aus_production |>
  model(
    `Seasonal_naïve` = SNAIVE(Beer),
    `Naïve` = NAIVE(Beer),
    Drift = RW(Beer ~ drift()),
    Mean = MEAN(Beer)
  )
```

```
# A mable: 1 x 4
  Seasonal_naïve   Naïve          Drift    Mean
          <model> <model>        <model> <model>
1       <SNAIVE> <NAIVE> <RW w/ drift>  <MEAN>
```

A `mable` is a model table, each cell corresponds to a fitted model.

Extract information from `mable`

```
beer_fit |> select(`Seasonal_naïve`) |> report()
beer_fit |> tidy()
beer_fit |> glance()
```

- The `report()` function gives a formatted model-specific display.
- The `tidy()` function is used to extract the coefficients from the models.
- We can extract information about some specific model using the `filter()` and `select()` functions.

Check model performance

Once a model has been fitted, it is important to check how well it has performed on the data. We come back to this later.

SECTION 5 OF 6

Produce forecasts

1 Statistical forecasting
2 What can we forecast?
3 Benchmark methods
4 Specify and estimate
5 **Produce forecasts**
6 Fitted values and residuals



Producing forecasts

- The `forecast()` function is used to produce forecasts from estimated models.
- **h** can be specified with:
  - a number (the number of future observations)
  - natural language (the length of time to predict)
  - provide a dataset of future time periods

Producing forecasts


```
beer_fc <- beer_fit |>
  forecast(h = "5 years")
```

```
# A fable: 80 x 4 [1Q]
# Key:     .model [4]
  .model Quarter         Beer .mean
  <chr>    <qtr>       <dist> <dbl>
1 Seaso… 2010 Q3 N(419, 373)   419
2 Seaso… 2010 Q4 N(488, 373)   488
3 Seaso… 2011 Q1 N(414, 373)   414
4 Seaso… 2011 Q2 N(374, 373)   374
# i 76 more rows
```

A `fable` is a forecast table with point forecasts and distributions.

Visualising forecasts

```
beer_fc |>
  autoplot(aus_production, level = NULL) +
  labs(title = "Forecasts for quarterly beer production",
       x = "Quarter", y = "Megalitres") +
  guides(colour = guide_legend(title = "Forecast"))
```



Forecast distributions

- A forecast $\hat{y}_{T+h|T}$ is (usually) the mean of the conditional distribution $y_{T+h} \mid y_1, \dots, y_T$.
- Most time series models produce normally distributed forecasts.
- The forecast distribution describes the probability of observing any future value.

Forecast distributions - normal distribution

Assuming residuals are normal, uncorrelated, sd = $\hat{\sigma}$:

Mean:	$\hat{y}_{T+h\|T} \sim N(\bar{y}, (1+1/T)\hat{\sigma}^2)$
Naïve:	$\hat{y}_{T+h\|T} \sim N(y_T, h\hat{\sigma}^2)$
Seasonal naïve:	$\hat{y}_{T+h\|T} \sim N(y_{T+h-m(k+1)}, (k+1)\hat{\sigma}^2)$
Drift:	$\hat{y}_{T+h\|T} \sim N(y_T + \frac{h}{T-1}(y_T - y_1), h\frac{T+h}{T}\hat{\sigma}^2)$

where k is the integer part of $(h-1)/m$.

Note that when $h=1$ and T is large, these all give the same approximate forecast variance: $\hat{\sigma}^2$.

Forecast distributions from bootstrapping

When a normal distribution for the residuals is an unreasonable assumption, one alternative is to use bootstrapping, which only assumes that the residuals are uncorrelated with constant variance.


- A one-step forecast error is defined as $e_t = y_t - \hat{y}_{t|t-1}$, $y_t = \hat{y}_{t|t-1} + e_t$.
- So we can simulate the next observation of a time series using $y_{T+1} = \hat{y}_{T+1|T} + e_{T+1}$
- Adding the new simulated observation to our data set, we can repeat the process to obtain $y_{T+2} = \hat{y}_{T+2|T+1} + e_{T+2}$

Generate many possible future using `generate()`

```
beer_2000 <- aus_production |> filter(year(Quarter) >= 2000) |> select(Beer)
fit <- beer_2000 |>
  model(NAIVE(Beer))
sim <- fit |> generate(h = 12, times = 5, bootstrap = TRUE)
sim
```

```
# A tsibble: 60 x 4 [1Q]
# Key:       .model, .rep [5]
   .model       Quarter .rep   .sim
   <chr>          <qtr> <chr> <dbl>
 1 NAIVE(Beer) 2010 Q3 1      320.
 2 NAIVE(Beer) 2010 Q4 1      362.
 3 NAIVE(Beer) 2011 Q1 1      290.
 4 NAIVE(Beer) 2011 Q2 1      272.
 5 NAIVE(Beer) 2011 Q3 1      199.
 6 NAIVE(Beer) 2011 Q4 1      221.
 7 NAIVE(Beer) 2012 Q1 1      178.
 8 NAIVE(Beer) 2012 Q2 1      220.
 9 NAIVE(Beer) 2012 Q3 1      172.
10 NAIVE(Beer) 2012 Q4 1      100.
# i 50 more rows
```

Generate 5 different futures



Use `forecast()` function to generate probabilistic forecasts

```
fc <- fit |> forecast(h = 12, bootstrap = TRUE)
fc
```

```
# A fable: 12 x 4 [1Q]
# Key:     .model [1]
   .model       Quarter          Beer .mean
   <chr>          <qtr>       <dist> <dbl>
 1 NAIVE(Beer) 2010 Q3 sample[5000]  374.
 2 NAIVE(Beer) 2010 Q4 sample[5000]  374.
 3 NAIVE(Beer) 2011 Q1 sample[5000]  374.
 4 NAIVE(Beer) 2011 Q2 sample[5000]  373.
 5 NAIVE(Beer) 2011 Q3 sample[5000]  374.
 6 NAIVE(Beer) 2011 Q4 sample[5000]  374.
 7 NAIVE(Beer) 2012 Q1 sample[5000]  374.
 8 NAIVE(Beer) 2012 Q2 sample[5000]  373.
 9 NAIVE(Beer) 2012 Q3 sample[5000]  374.
10 NAIVE(Beer) 2012 Q4 sample[5000]  373.
11 NAIVE(Beer) 2013 Q1 sample[5000]  373.
12 NAIVE(Beer) 2013 Q2 sample[5000]  371.
```

Prediction intervals

- Forecast intervals can be extracted using the `hilo()` function.

```
fit <- aus_production |> select(Beer) |> model(NAIVE(Beer))
forecast(fit) |> hilo(level = 95) |> unpack_hilo("95%")
```

```
# A tsibble: 8 x 6 [1Q]
# Key:       .model [1]
  .model Quarter          Beer .mean
  <chr>    <qtr>         <dist> <dbl>
1 NAIVE… 2010 Q3  N(374, 4580)   374
2 NAIVE… 2010 Q4  N(374, 9159)   374
3 NAIVE… 2011 Q1 N(374, 13739)   374
4 NAIVE… 2011 Q2 N(374, 18319)   374
5 NAIVE… 2011 Q3 N(374, 22898)   374
6 NAIVE… 2011 Q4 N(374, 27478)   374
7 NAIVE… 2012 Q1 N(374, 32057)   374
8 NAIVE… 2012 Q2 N(374, 36637)   374
# i 2 more variables: `95%_lower` <dbl>, `95%_upper` <dbl>
```

Prediction intervals

```
beer_fc |>
  hilo(level = c(50, 75))
```

```
# A tsibble: 80 x 6 [1Q]
# Key:       .model [4]
   .model       Quarter          Beer
   <chr>          <qtr>        <dist>
 1 Seasonal_n… 2010 Q3  N(419, 373)
 2 Seasonal_n… 2010 Q4  N(488, 373)
 3 Seasonal_n… 2011 Q1  N(414, 373)
 4 Seasonal_n… 2011 Q2  N(374, 373)
 5 Seasonal_n… 2011 Q3  N(419, 747)
 6 Seasonal_n… 2011 Q4  N(488, 747)
 7 Seasonal_n… 2012 Q1  N(414, 747)
 8 Seasonal_n… 2012 Q2  N(374, 747)
 9 Seasonal_n… 2012 Q3 N(419, 1120)
10 Seasonal_n… 2012 Q4 N(488, 1120)
# i 70 more rows
# i 3 more variables: .mean <dbl>, `50%` <hilo>, `75%` <hilo>
```

Prediction intervals

```
beer_fc |>
  hilo(level = c(50, 75)) |>
  mutate(lower = `50%`$lower, upper = `50%`$upper)
```

```
# A tsibble: 80 x 8 [1Q]
# Key:       .model [4]
   .model       Quarter          Beer
   <chr>          <qtr>        <dist>
 1 Seasonal_n… 2010 Q3  N(419, 373)
 2 Seasonal_n… 2010 Q4  N(488, 373)
 3 Seasonal_n… 2011 Q1  N(414, 373)
 4 Seasonal_n… 2011 Q2  N(374, 373)
 5 Seasonal_n… 2011 Q3  N(419, 747)
 6 Seasonal_n… 2011 Q4  N(488, 747)
 7 Seasonal_n… 2012 Q1  N(414, 747)
 8 Seasonal_n… 2012 Q2  N(374, 747)
 9 Seasonal_n… 2012 Q3 N(419, 1120)
10 Seasonal_n… 2012 Q4 N(488, 1120)
# i 70 more rows
# i 5 more variables: .mean <dbl>, `50%` <hilo>, `75%` <hilo>, lower <dbl>, upper <dbl>
```

SECTION 6 OF 6

Fitted values and residuals

1 Statistical forecasting
2 What can we forecast?
3 Benchmark methods
4 Specify and estimate
5 Produce forecasts
6 **Fitted values and residuals**



Fitted values

- $\hat{y}_{t|t-1}$ is the forecast of y_t based on observations $y_1, \ldots, y_{t-1}$.
- We call these “fitted values”.
- Sometimes drop the subscript: $\hat{y}_t \equiv \hat{y}_{t|t-1}$.
- Often not true forecasts since parameters are estimated on all data.

For example:

- $\hat{y}_t = \bar{y}$ for average method.
- $\hat{y}_t = y_{t-1} + (y_T - y_1)/(T-1)$ for drift method.

Forecasting residuals

Residuals in forecasting: difference between observed value and its fitted value:
$e_t = y_t - \hat{y}_{t|t-1}$.


Beer production - augment

```
fit <- aus_production |> select(Beer) |> model(SNAIVE(Beer))
augment(fit)
```

```
# A tsibble: 218 x 6 [1Q]
# Key:       .model [1]
   .model        Quarter  Beer .fitted .resid .innov
   <chr>           <qtr> <dbl>   <dbl>  <dbl>  <dbl>
 1 SNAIVE(Beer) 1956 Q1   284      NA     NA     NA
 2 SNAIVE(Beer) 1956 Q2   213      NA     NA     NA
 3 SNAIVE(Beer) 1956 Q3   227      NA     NA     NA
 4 SNAIVE(Beer) 1956 Q4   308      NA     NA     NA
 5 SNAIVE(Beer) 1957 Q1   262     284    -22    -22
 6 SNAIVE(Beer) 1957 Q2   228     213     15     15
 7 SNAIVE(Beer) 1957 Q3   236     227      9      9
 8 SNAIVE(Beer) 1957 Q4   320     308     12     12
 9 SNAIVE(Beer) 1958 Q1   272     262     10     10
10 SNAIVE(Beer) 1958 Q2   233     228      5      5
# i 208 more rows
```

Beer production - fitted values

```
augment(fit) |>
  ggplot(aes(x = Quarter)) +
  geom_line(aes(y = Beer, colour = "Data")) +
  geom_line(aes(y = .fitted, colour = "Fitted"))
```



Beer production - residuals

```
augment(fit) |>
  autoplot(.resid) +
  labs(x = "Quarter", y = "", title = "Residuals from snaïve method")
```

