

## Worksheet 6: Giant Planets and Small Bodies

### Full solutions

Covers Lectures 11 and 12: giant-planet interiors, the Roche limit and rings, radioactive dating, Kirkwood gaps, cometary activity

Planetary Systems (WBAS002-05) • Kapteyn Astronomical Institute, University of Groningen

This worksheet is ungraded practice, with the same shape and level as the final exam. Plan up to 2 hours for a serious pass. Work each problem through on paper before you open the solutions, and show your algebra: the exam asks for the steps, not only the number. Some parts state an intermediate result ("show that..."); use it to check your work, and to continue if a step resists. Carry at least four significant figures through the intermediate steps (the worked solutions print five) and round only at the end. All parts are analytical; no computer is needed.

#### Constants and data

$$G = 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}, \quad \ln 2 = 0.6931$$

Jupiter:  $M = 1.898 \times 10^{27} \text{ kg}$ ,  $R = 6.991 \times 10^7 \text{ m}$ , semi-major axis  $a_J = 5.203 \text{ AU}$ , orbital period  $T_J = 11.86 \text{ yr}$ . Saturn:  $M = 5.683 \times 10^{26} \text{ kg}$ , volumetric mean radius  $R_p = 58,232 \text{ km}$ , bulk density  $\rho_p = 687 \text{ kg m}^{-3}$

Metallic-hydrogen transition near 100 GPa. Ring and moon densities: water ice  $\rho_s = 1000 \text{ kg m}^{-3}$ . Saturn A-ring outer edge observed at  $\sim 137,000 \text{ km}$

$^{26}\text{Al}$ : half-life  $t_{1/2} = 0.717 \text{ Myr}$ , Solar-System initial ratio  $(^{26}\text{Al}/^{27}\text{Al})_0 = 5.2 \times 10^{-5}$ .

The uniform-density central pressure of a self-gravitating sphere in hydrostatic equilibrium is  $P_c = 3GM^2/(8\pi R^4)$ ; the fluid Roche limit is  $d_R = 2.46 R_p (\rho_p/\rho_s)^{1/3}$  and the rigid-body value is  $d_R = 1.26 R_p (\rho_p/\rho_s)^{1/3}$ . Data as in the Lecture 11 and 12 notes.

### Problem 1 Giant-planet interiors and the metallic-hydrogen transition

Jupiter and Saturn are fluid hydrogen-helium planets. Their deep interiors reach pressures high enough to squeeze hydrogen into a metallic, electrically conducting state. A crude uniform-density model already shows where and why this happens.

(a) Compute Jupiter's bulk density from its mass and radius.

#### Solution.

$$\rho = \frac{M}{\frac{4}{3}\pi R^3} = \frac{1.898 \times 10^{27}}{\frac{4}{3}\pi (6.991 \times 10^7)^3} = \frac{1.898 \times 10^{27}}{\frac{4}{3}\pi (3.4168 \times 10^{23})},$$

$$\text{so with } \frac{4}{3}\pi (3.4168 \times 10^{23}) = 1.4312 \times 10^{24} \text{ m}^3,$$

$$\rho = \frac{1.898 \times 10^{27}}{1.4312 \times 10^{24}} = 1326 \text{ kg m}^{-3}.$$

**$\rho \approx 1326 \text{ kg m}^{-3}$ , close to the bulk density of the Sun and far below rock, so most**

**of the planet is hydrogen and helium.**

**(b)** Using the uniform-density result  $P_c = 3GM^2/(8\pi R^4)$ , estimate Jupiter's central pressure. Express it in GPa and compare it with the  $\sim 100$  GPa metallic-hydrogen transition.

**Solution.**

$$P_c = \frac{3GM^2}{8\pi R^4} = \frac{3(6.674 \times 10^{-11})(1.898 \times 10^{27})^2}{8\pi(6.991 \times 10^7)^4}.$$

The numerator is  $3 \times 6.674 \times 10^{-11} \times 3.6024 \times 10^{54} = 7.2121 \times 10^{44}$ , and the denominator is  $8\pi \times 2.3887 \times 10^{31} = 6.0034 \times 10^{32}$ , so

$$P_c = \frac{7.2121 \times 10^{44}}{6.0034 \times 10^{32}} = 1.201 \times 10^{12} \text{ Pa} = 1201 \text{ GPa}.$$

This is about 12 times the 100 GPa needed to metallize hydrogen.

**Even the uniform-density estimate,  $\sim 1200$  GPa, exceeds the metallization pressure by an order of magnitude, so most of Jupiter's hydrogen is metallic. The real centrally-condensed value is a few thousand GPa, higher still.**

**(c)** Repeat the estimate for Saturn. Compare the two central pressures against the metallization pressure and state what this implies for the relative size of each planet's metallic-hydrogen region.

**Solution.**

$$P_c = \frac{3(6.674 \times 10^{-11})(5.683 \times 10^{26})^2}{8\pi(5.8232 \times 10^7)^4}.$$

The numerator is  $3 \times 6.674 \times 10^{-11} \times 3.2296 \times 10^{53} = 6.4669 \times 10^{43}$ , and the denominator is  $8\pi \times 1.1499 \times 10^{31} = 2.8900 \times 10^{32}$ , so

$$P_c = \frac{6.4669 \times 10^{43}}{2.8900 \times 10^{32}} = 2.238 \times 10^{11} \text{ Pa} = 224 \text{ GPa}.$$

Saturn's central pressure is about 2.2 times the metallization value, against Jupiter's factor of 12.

**Jupiter sits far above the transition throughout most of its interior, so it has a large metallic-hydrogen region. Saturn only just crosses it near the centre, so its metallic region is a much smaller fraction of the planet.**

**(d)** Saturn radiates more energy than it absorbs from the Sun, more than slow contraction alone can supply, while Jupiter's excess luminosity is closer to what contraction gives. Explain in two or three sentences how the smaller metallic-hydrogen region you found in part (c) is linked to the extra energy source (helium rain) that Saturn needs.

**Solution.** Helium becomes immiscible in metallic hydrogen at the pressures and temperatures of the deep interior, so helium droplets condense and sink, releasing gravitational energy. This process operates in the metallic-hydrogen layer. Saturn is cooler and its metallic region, though smaller, spans the pressure-temperature range where helium separates, so helium rain runs strongly and supplies the extra luminosity. Jupiter is hotter and its interior has only recently entered the helium-separation range, so its excess heat is still dominated by contraction.

**The size and thermal state of the metallic-hydrogen region set whether helium rain is active; Saturn needs it, Jupiter barely does.**

## Problem 2 The Roche limit and Saturn's rings

The inner edge of the region where a moon is tidally torn apart, rather than held together by its own gravity, is the Roche limit. It sets where rings can exist and where moons can survive. Saturn is the clearest case.

**(a)** Compute the rigid-body Roche limit for an icy body at Saturn,  $d_R = 1.26 R_p (\rho_p / \rho_s)^{1/3}$ . Express it in units of  $R_p$ .

**Solution.** With  $(\rho_p / \rho_s)^{1/3} = (687 / 1000)^{1/3} = 0.882$ ,

$$d_R = 1.26 \times 58,232 \text{ km} \times 0.882 = 64,710 \text{ km} = 1.111 R_p.$$

**The rigid-body limit is  $\approx 64,710 \text{ km}$ , only  $1.11 R_p$ , just above Saturn's surface: a rigid body could in principle survive almost down to the cloud tops.**

**(b)** Compute the fluid Roche limit,  $d_R = 2.46 R_p (\rho_p / \rho_s)^{1/3}$ . Compare it with the observed outer edge of the A ring at  $\sim 137,000 \text{ km}$ .

**Solution.**

$$d_R = 2.46 \times 58,232 \text{ km} \times 0.882 = 126,350 \text{ km} = 2.170 R_p.$$

The A-ring outer edge lies at  $137,000 \text{ km}$ , so the fluid estimate matches to

$$\frac{137,000 - 126,350}{137,000} = 7.8\%.$$

**The fluid Roche limit,  $\approx 126,350 \text{ km}$ , matches the A-ring outer edge to within about 8%: the main rings end near where a self-gravitating fluid body would be disrupted.**

**(c)** Small moons such as Pan and Daphnis orbit inside the A ring, well inside the fluid Roche limit of part (b) but outside the rigid-body limit of part (a). Explain in two or three sentences how both facts can be true, and what the two limits bracket.

**Solution.** The fluid limit assumes a strengthless body that deforms and disrupts easily; the rigid limit assumes a body held together as a solid sphere. A real icy moonlet has finite material strength, so its true disruption distance lies between the two. Pan and Daphnis sit between  $64,710 \text{ km}$  and  $126,350 \text{ km}$ , so their own gravity plus modest material strength holds them together even though a strengthless fluid body there would be torn apart.

**The rigid and fluid limits bracket the real disruption distance; bodies with finite strength survive between them, which is why solid moonlets exist inside the fluid Roche limit.**

## Problem 3 Dating the Solar System with radioactive decay

Meteorites carry two kinds of radioactive clock. Short-lived isotopes such as  $^{26}\text{Al}$  are now extinct but timed the first few million years; long-lived isotopes such as uranium give absolute ages of billions of years. You will use both.

(a) For radioactive decay  $N = N_0 e^{-\lambda t}$  with  $\lambda = \ln 2 / t_{1/2}$ , compute the decay constant of  $^{26}\text{Al}$  and the fraction of the initial  $^{26}\text{Al}$  left after 2 Myr.

**Solution.**

$$\lambda = \frac{\ln 2}{t_{1/2}} = \frac{0.6931}{0.717} = 0.9667 \text{ Myr}^{-1}.$$

After 2 Myr,

$$\frac{N}{N_0} = e^{-\lambda t} = e^{-0.9667 \times 2} = e^{-1.9334} = 0.1447.$$

**|  $\lambda \approx 0.9667 \text{ Myr}^{-1}$ ; about 14.5% of the  $^{26}\text{Al}$  survives after 2 Myr, so most of it decays within the first few million years.**

(b) Radiogenic heating scales with the amount of live  $^{26}\text{Al}$  present at accretion. A planetesimal that accretes 2 Myr after the first solids (CAIs) starts with the Solar-System initial  $^{26}\text{Al}/^{27}\text{Al}$  multiplied by the surviving fraction you found in part (a); compute this ratio. State the factor by which its heating rate is lower than a body that formed at CAI time, and what this implies for melting.

**Solution.**

$$\left(\frac{^{26}\text{Al}}{^{27}\text{Al}}\right) = 5.2 \times 10^{-5} \times 0.1447 = 7.52 \times 10^{-6}.$$

The heating rate is proportional to the live  $^{26}\text{Al}$ , so it is lower by the factor  $1/0.1447 = 6.9$ .

**| A body forming 2 Myr late has  $\approx 7.52 \times 10^{-6}$  live  $^{26}\text{Al}/^{27}\text{Al}$ , a factor 6.9 less heating. Early planetesimals melt and differentiate; those forming more than a couple of million years late stay cold and undifferentiated.**

(c) Explain in two or three sentences why the short-lived  $^{26}\text{Al}$  clock gives high time resolution but only relative ages, while the long-lived U-Pb clock gives absolute ages but coarser resolution over the first few million years.

**Solution.**  $^{26}\text{Al}$  decays in a few million years, so small time differences change its abundance a lot, giving fine relative timing; but it is now extinct, so it cannot date against a present-day measurement and needs an anchor to an absolute clock. U-Pb decays over billions of years, so it survives to be measured today and gives an absolute age; but over the first few million years the accumulated radiogenic lead changes very little, so its resolution there is coarse.

**| Short-lived isotopes resolve fine intervals but only relative to an anchor; long-lived isotopes give the absolute age but blur the earliest few million years.**

## Problem 4 Kirkwood gaps and mean-motion resonances

The asteroid belt has gaps at specific distances where an asteroid's orbital period is a simple fraction of Jupiter's. These mean-motion resonances pump orbital eccentricities

and clear the gaps. They are the Kirkwood gaps.

**(a)** At an interior  $j : k$  resonance the asteroid completes  $j$  orbits for every  $k$  of Jupiter's, so its period is  $T = (k/j)T_J$ . Using Kepler's third law  $a^3 = T^2$  (with  $a$  in AU and  $T$  in yr, both scaled to  $a_J$ ), show that the resonance semi-major axis is  $a = a_J (k/j)^{2/3}$ , and compute the 3 : 1 location.

**Solution.** From  $a^3 \propto T^2$ , the ratio of two orbits gives  $a/a_J = (T/T_J)^{2/3} = (k/j)^{2/3}$ . For the 3 : 1 resonance,  $k/j = 1/3$ , so

$$a = 5.203 \text{ AU} \times (1/3)^{2/3} = 5.203 \times 0.4807 = 2.501 \text{ AU}.$$

**|  $a \approx 2.50 \text{ AU}$ , matching the observed inner Kirkwood gap.**

**(b)** Compute the locations of the 5 : 2 and 2 : 1 resonances.

**Solution.**

$$a_{5:2} = 5.203 \times (2/5)^{2/3} = 5.203 \times 0.5429 = 2.825 \text{ AU},$$

$$a_{2:1} = 5.203 \times (1/2)^{2/3} = 5.203 \times 0.6300 = 3.278 \text{ AU}.$$

**| The 5 : 2 gap is at  $\approx 2.82 \text{ AU}$  and the 2 : 1 gap at  $\approx 3.28 \text{ AU}$ , the outer edge of the main belt.**

**(c)** An asteroid exactly at a Kirkwood gap and one just outside it look identical in a snapshot of positions. Explain in two or three sentences what physically empties the gap, and why the gaps are visible in a plot of semi-major axis but not in an instantaneous map of the belt.

**Solution.** At the resonance Jupiter kicks the asteroid at the same orbital phase every cycle, and these kicks add coherently, so the eccentricity grows over many orbits. A high eccentricity eventually makes the orbit cross Mars or Jupiter, and a close encounter removes the asteroid. The clearing acts on the orbit's semi-major axis, not its instantaneous position, so it shows only when bodies are binned by  $a$ ; at any instant a resonant asteroid can be anywhere along its orbit and looks ordinary.

**| Coherent resonant kicks pump eccentricity until the orbit becomes planet-crossing and the body is removed; the depletion appears only in semi-major-axis space.**