

Worksheet 2: Thermal Evolution and Differentiation

Full solutions

Covers Lectures 3 and 4: heat sources, convection, core formation, dynamos

Planetary Systems (WBAS002-05) • Kapteyn Astronomical Institute, University of Groningen

This worksheet is ungraded practice, with the same shape and level as the final exam. Plan up to 3 hours for a serious pass. Work each problem through on paper before you open the solutions, and show your algebra: the exam asks for the steps, not only the number. Some parts state an intermediate result ("show that..."); use it to check your work, and to continue if a step resists. Carry at least four significant figures through the intermediate steps (the worked solutions print five) and round only at the end. Problem 3 has a short computational part; everything else is analytical.

Constants and data

1 yr = 3.156×10^7 s 1 d = 86 400 s $M_{\oplus} = 5.972 \times 10^{24}$ kg $R_{\oplus} = 6371$ km $R_{\text{Moon}} = 1737$ km

Thermal diffusivity of rock $\kappa = 10^{-6} \text{ m}^2 \text{ s}^{-1}$ Age of the solar system 4.57 Gyr Earth's core mass fraction $x = 0.32$ silicate heat capacity $c_p = 1250 \text{ J kg}^{-1} \text{ K}^{-1}$

Isotope	Half-life (Gyr)	Heat production (W per kg isotope)	BSE concentration
^{238}U	4.47	9.5×10^{-5}	20 ppb
^{235}U	0.704	5.7×10^{-4}	0.14 ppb
^{232}Th	14.0	2.6×10^{-5}	80 ppb
^{40}K	1.25	2.9×10^{-5}	28 ppb

Isotope table as in the Lecture 3 notes (BSE = bulk silicate Earth; 1 ppb = 10^{-9} by mass). Hf-W constants as in the Lecture 4 notes: $Q_W = 10^4$, $(^{182}\text{Hf}/^{180}\text{Hf})_0 = 1.02 \times 10^{-4}$, $\lambda = 7.79 \times 10^{-2} \text{ Myr}^{-1}$ ($t_{1/2} = 8.9 \text{ Myr}$).

Problem 1 The radiogenic clock

Radioactive decay powers planetary interiors on two very different timescales. Here you reconstruct both from the isotope table, then place them in the planetary context.

(a) The isotope table refers to the bulk silicate Earth (BSE): the mantle plus crust, without the core. Compute the BSE mass from $M_{\oplus}$ and the core mass fraction x (show that $M_{\text{BSE}} \approx 4.06 \times 10^{24} \text{ kg}$), then compute the present-day specific heating rate of BSE rock by summing the four isotopes (show that it is $4.872 \times 10^{-12} \text{ W kg}^{-1}$). Using the round value $M_{\text{BSE}} = 4.0 \times 10^{24} \text{ kg}$, give Earth's present radiogenic power in TW and compare it with the value quoted in the notes.

Solution. The BSE mass is the non-core fraction of Earth's mass:

$$M_{\text{BSE}} = (1 - 0.32) \times 5.972 \times 10^{24} = 4.0610 \times 10^{24} \text{ kg}.$$

Each isotope contributes (concentration) $\times$ (heat production per kg of isotope):

$$^{238}\text{U} : 20 \times 10^{-9} \times 9.5 \times 10^{-5} = 1.9000 \times 10^{-12} \text{ W kg}^{-1},$$

$$^{235}\text{U} : 0.14 \times 10^{-9} \times 5.7 \times 10^{-4} = 7.9800 \times 10^{-14}, \quad ^{232}\text{Th} : 80 \times 10^{-9} \times 2.6 \times 10^{-5} = 2.0800 \times 10^{-12},$$

$$^{40}\text{K} : 28 \times 10^{-9} \times 2.9 \times 10^{-5} = 8.1200 \times 10^{-13}.$$

The sum is

$$h_{\text{today}} = (1.9000 + 0.0798 + 2.0800 + 0.8120) \times 10^{-12} = 4.8718 \times 10^{-12} \text{ W kg}^{-1},$$

which matches the printed checkpoint. With the round BSE mass:

$$P_{\text{today}} = 4.872 \times 10^{-12} \times 4.0 \times 10^{24} = 1.9488 \times 10^{13} \text{ W} \approx 19.5 \text{ TW}.$$

| $P_{\text{today}} \approx 19.5 \text{ TW}$, consistent with the 19–20 TW quoted in the Lecture 3 notes.

(b) Each isotope was more abundant in the past by a factor $2^{t/t_{1/2}}$ with $t = 4.5 \text{ Gyr}$. Compute this factor for each isotope, then the specific heating rate at Earth's formation (show that it is $2.297 \times 10^{-11} \text{ W kg}^{-1}$), the corresponding power, and the ratio of the heating rate at formation to today.

Solution. The abundance factors over $t = 4.5 \text{ Gyr}$:

$$^{238}\text{U} : 2^{4.5/4.47} = 2^{1.0067} = 2.0093, \quad ^{235}\text{U} : 2^{4.5/0.704} = 2^{6.3920} = 83.98,$$

$$^{232}\text{Th} : 2^{4.5/14.0} = 2^{0.32143} = 1.2497, \quad ^{40}\text{K} : 2^{4.5/1.25} = 2^{3.6000} = 12.126.$$

Scaling each present-day contribution from part (a) by its factor:

$$1.9000 \times 2.0093 = 3.8177, \quad 0.0798 \times 83.98 = 6.7016,$$

$$2.0800 \times 1.2497 = 2.5994, \quad 0.8120 \times 12.126 = 9.8463,$$

all in units of $10^{-12} \text{ W kg}^{-1}$. The sum is

$$h_{\text{form}} = 2.2965 \times 10^{-11} \text{ W kg}^{-1},$$

matching the checkpoint. The power and the ratio:

$$P_{\text{form}} = 2.297 \times 10^{-11} \times 4.0 \times 10^{24} = 9.188 \times 10^{13} \text{ W} \approx 92 \text{ TW}, \quad \frac{h_{\text{form}}}{h_{\text{today}}} = \frac{2.297 \times 10^{-11}}{4.872 \times 10^{-12}} = 4.71.$$

| $P_{\text{form}} \approx 92 \text{ TW}$ and $h_{\text{form}}/h_{\text{today}} \approx 4.7$: exactly the factor and the formation-era power printed in the Lecture 3 notes and figure.

(c) The radiogenic-heating figure in the Lecture 3 notes shows the short-lived isotope ^{26}Al ($t_{1/2} = 0.72 \text{ Myr}$) crossing below the summed long-lived curve. Read off the approximate crossover time from the figure. How many ^{26}Al half-lives is that (show that it is ≈ 14), and by what factor has ^{26}Al decayed by then?

Solution. The dashed ^{26}Al curve crosses the solid long-lived total at roughly 10 Myr

after CAI formation. That is

$$\frac{10 \text{ Myr}}{0.72 \text{ Myr}} = 13.9 \approx 14 \text{ half-lives,}$$

so the isotope has decayed by

$$2^{13.9} = 1.5 \times 10^4.$$

I Crossover near 10 Myr, about 14 half-lives, a decline by a factor $\sim 1.5 \times 10^4$: short-lived heating is a property of the first few Myr only.

(d) In 2 to 3 sentences: why do U, Th, and K reside in the mantle and crust rather than in the core, and what does this imply for *where* radiogenic heat is released inside Earth?

Solution. U, Th, and K are lithophile in the Goldschmidt classification: during metal-silicate separation they stay in the silicate phase rather than following the iron into the core. As incompatible elements they are further concentrated into the last-crystallising liquids, and so into the crust. Radiogenic heat is therefore released in the silicate shell, roughly half of it in the thin crust, and essentially none inside the core, whose heat budget is set by cooling and crystallisation instead.

I Lithophile behaviour keeps the heat producers in the silicate shell; the core receives essentially no radiogenic heating.

Problem 2 Conduction fails, convection takes over

Whether a body can cool by conduction alone is a question of size. Here you find the size limit, then show that Earth's mantle convects instead, and how vigorously.

(a) Compute the conductive cooling timescale $\tau = L^2/\kappa$ for the Moon, using $L = R_{\text{Moon}}$ (show that $\tau \approx 9.6 \times 10^{10}$ yr), and compare with the age of the solar system. Then invert the timescale: what is the largest body size that *can* cool conductively within 4.57 Gyr?

Solution.

$$L^2 = (1.737 \times 10^6)^2 = 3.0172 \times 10^{12} \text{ m}^2, \quad \tau = \frac{3.0172 \times 10^{12}}{10^{-6}} = 3.0172 \times 10^{18} \text{ s.}$$

In years:

$$\tau = \frac{3.0172 \times 10^{18}}{3.156 \times 10^7} = 9.5602 \times 10^{10} \text{ yr,}$$

matching the checkpoint: about 21 times the age of the solar system, so even the Moon cannot have cooled through its full radius by conduction alone. Inverting, with the age in seconds $t = 4.57 \times 10^9 \times 3.156 \times 10^7 = 1.4423 \times 10^{17}$ s:

$$L = \sqrt{\kappa t} = \sqrt{10^{-6} \times 1.4423 \times 10^{17}} = \sqrt{1.4423 \times 10^{11}} = 3.798 \times 10^5 \text{ m} \approx 380 \text{ km.}$$

I $\tau \approx 9.6 \times 10^{10}$ yr $\gg$ 4.57 Gyr; only bodies below ≈ 380 km cool conductively, larger ones must convect, or stay hot.

(b) Compute the Rayleigh number of Earth's mantle,

$$Ra = \frac{\alpha \rho g \Delta T d^3}{\kappa \eta},$$

with $\alpha = 2 \times 10^{-5} \text{ K}^{-1}$, $\rho = 4000 \text{ kg m}^{-3}$, $g = 10 \text{ m s}^{-2}$, $\Delta T = 2500 \text{ K}$, mantle depth $d = 3000 \text{ km}$, and dynamic viscosity $\eta = 10^{21} \text{ Pa s}$ (show that $Ra \approx 5.4 \times 10^7$). Compare with the critical value $Ra_c \approx 1708$.

Solution. Numerator first:

$$\alpha \rho g \Delta T = 2 \times 10^{-5} \times 4000 \times 10 \times 2500 = 2000 \text{ kg m}^{-3} \text{ s}^{-2}, \quad d^3 = (3 \times 10^6)^3 = 2.7 \times 10^{19} \text{ m}^3,$$

so the numerator is $2000 \times 2.7 \times 10^{19} = 5.4 \times 10^{22}$. The denominator:

$$\kappa \eta = 10^{-6} \times 10^{21} = 10^{15}.$$

Therefore

$$Ra = \frac{5.4 \times 10^{22}}{10^{15}} = 5.4 \times 10^7,$$

matching the checkpoint, and

$$\frac{Ra}{Ra_c} = \frac{5.4 \times 10^7}{1708} = 3.162 \times 10^4.$$

| $Ra \approx 5.4 \times 10^7$, **over 3×10^4 times critical: the mantle convects vigorously despite its enormous viscosity.**

(c) The Nusselt number, normalised to vanish at onset, is $Nu = (Ra/Ra_c)^{1/3}$. Compute Nu for the mantle and state what it means for the heat flux.

Solution. From the printed ratio of part (b):


$$Nu = (3.162 \times 10^4)^{1/3} = 31.62 \approx 32.$$

The Nusselt number is the ratio of the actual (convective) heat flux to the flux that conduction alone would carry across the same layer and temperature drop: convection moves heat about 32 times faster than conduction would.

| $Nu \approx 32$, **inside the 18–39 range quoted in the notes for $Ra = 10^7$ – 10^8 .**

(d) Sketch the horizontally averaged temperature profile through Earth's mantle, from the cold surface to the core-mantle boundary at 2890 km depth. Label the two thermal boundary layers and the nearly adiabatic interior, with approximate thicknesses and the anchor temperatures from the notes. In one sentence: why is almost the entire temperature drop confined to the two thin boundary layers?

Solution. The profile has three segments: a steep conductive gradient across the cold thermal boundary layer (the lithosphere, ~100 km); a shallow, nearly adiabatic gradient (~0.5 K km⁻¹) through the stirred interior, reaching ~2500 K at the base of the mantle; and a second steep gradient across the hot boundary layer (the D'' layer, ~200 km), rising to $T_{\text{CMB}} \sim 4000 \text{ K}$.



The interior is stirred by convection, which keeps it close to the shallow adiabat: a large gradient cannot survive there, so the full contrast between surface and core is squeezed into the two thin conductive layers at the top and bottom.

Steep gradients exist only where heat moves by conduction: in the two thin boundary layers. The convecting interior follows the adiabat.

(e) Judge this claim in 2 to 3 sentences: “Doubling the mantle’s temperature contrast ΔT would double the convective heat flux.”

Solution. False. The convective flux is $q \propto \text{Nu} \times \Delta T$ with $\text{Nu} \propto \text{Ra}^{1/3} \propto \Delta T^{1/3}$, so $q \propto \Delta T^{4/3}$: doubling ΔT raises the flux by $2^{4/3} = 2.52$, more than double. A hotter mantle does not just carry more heat per parcel, it also convects more vigorously.

False: $q \propto \Delta T^{4/3}$, so the flux rises by a factor 2.52.

Problem 3 A minimal thermal evolution model

The heat budget of the Lecture 3 notes (radiogenic heating in, convective heat loss out) can be turned into a one-line model of Earth’s thermal history. This problem assembles the model on paper and then integrates it in Python.

(a) Problem 2(e) already used the scaling $q \propto \Delta T^{4/3}$; derive it now in one line from $\text{Nu} \propto \text{Ra}^{1/3}$ and $\text{Ra} \propto \Delta T$ (all other parameters fixed). Writing T for the mantle temperature contrast, energy conservation for the silicate shell then reads

$$C \frac{dT}{dt} = H(t) - Q(T), \quad Q(T) = Q_0 \left(\frac{T}{T_0} \right)^{4/3}, \quad H(t) = H_f e^{-t/\tau_r},$$

with $C = M_{\text{BSE}} c_p$ the heat capacity of the silicate shell. State in one sentence each what the two right-hand terms represent.

Solution. The conductive flux across the layer is $q_{\text{cond}} \propto \Delta T$, and convection multiplies it by Nu:

$$Q \propto \text{Nu} \times \Delta T \propto \Delta T^{1/3} \times \Delta T = \Delta T^{4/3}.$$

$H(t)$ is the radiogenic heat input, a single exponential standing in for the summed

decay of the four long-lived isotopes; $Q(T)$ is the convective heat loss through the surface, rising steeply with mantle temperature. The balance of the two drives dT/dt .

$Q \propto \Delta T^{4/3}$; heating in, parametrised convective cooling out.

(b) Take today's values $H = 20$ TW, $Q_0 = 47$ TW at $T_0 = 2500$ K, and $C = 4.0 \times 10^{24} \times 1250 = 5.0 \times 10^{27}$ J K⁻¹. Show that the model's present-day cooling rate is $dT/dt \approx -170$ K per Gyr. Name one reason this simple estimate overstates the true mantle cooling rate.

Solution.

$$\frac{dT}{dt} = \frac{(20 - 47) \times 10^{12}}{5.0 \times 10^{27}} = -5.400 \times 10^{-15} \text{ K s}^{-1}.$$

Per Gyr (3.156×10^{16} s):

$$\frac{dT}{dt} = -5.400 \times 10^{-15} \times 3.156 \times 10^{16} = -170.4 \text{ K Gyr}^{-1},$$

matching the checkpoint. The estimate charges the full 47 TW to the silicate shell, but roughly 10 TW of it is heat leaving the core (which is resupplied by core cooling and crystallisation, not by the mantle's own reservoir), so the mantle itself cools more slowly; estimates from petrology give closer to 50–100 K per Gyr.


$dT/dt \approx -170 \text{ K Gyr}^{-1}$, an overestimate because part of the surface heat flow is paid by the core, not the mantle.

(c) The template `thermal_evolution.py` (next to this sheet on the course site) integrates the model from $T(0) = 3000$ K over 4.5 Gyr with $H_f = 94$ TW (today's round 20 TW multiplied by the factor 4.7; the more precise chain of Problem 1(b) gives 92 TW) and $\tau_r = 2.91$ Gyr. Complete the two marked lines (the expressions for $H(t)$ and $Q(T)$), run the script, and report: the maximum temperature and when it occurs, the final temperature $T(4.5 \text{ Gyr})$, and the final Urey ratio H/Q . Your run should give $T(4.5 \text{ Gyr}) \approx 2670$ K. Describe the shape of the curve in one or two sentences.

Solution. The completed lines are

```
H = H_F * np.exp(-t / TAU_R)
Q = Q_0 * (T / T_0) ** (4.0 / 3.0)
```

The run gives a maximum of $T \approx 3110$ K at $t \approx 1.2$ Gyr, a final temperature $T(4.5 \text{ Gyr}) \approx 2670$ K, and a final Urey ratio $H/Q = 20.0/51.4 \approx 0.39$. The curve first *rises*: at 3000 K the convective loss (≈ 60 TW) is smaller than the initial heating (94 TW), so the interior warms until Q catches up with H ; from ~ 1.2 Gyr onward the planet tracks the decaying heat source downward, cooling ever after. The solution figure shows $T(t)$, $H(t)$, and $Q(t)$:



Peak ≈ 3110 K at ≈ 1.2 Gyr; $T(4.5) \approx 2670$ K; Urey ratio ≈ 0.39 . Early self-heating, then long secular cooling.

(d) Set $H = 0$ in the model and solve it analytically: show that at late times the temperature falls off as $T \propto t^{-3}$. What does this limit say about how quickly a small, radiogenically dead body fades?

Solution. With $H = 0$,

$$C \frac{dT}{dt} = -\frac{Q_0}{T_0^{4/3}} T^{4/3}.$$

Separate variables:

$$T^{-4/3} dT = -\frac{Q_0}{C T_0^{4/3}} dt \quad \Rightarrow \quad -3 T^{-1/3} = -\frac{Q_0}{C T_0^{4/3}} t + \text{const},$$

so $T^{-1/3}$ grows linearly with t , and at late times

$$T \propto t^{-3}.$$

A body without radiogenic heating cools fast at first (the $T^{4/3}$ loss is strong while it is hot) but fades on a long power-law tail rather than exponentially: cooling slows as the driving contrast collapses.

$T \propto t^{-3}$: fast early cooling, then a slow power-law fade.

Problem 4 Building a core and dating it

Core formation is fast, chemically selective, and datable. Here you follow one iron droplet down, predict what it leaves behind, and read the isotopic clock it starts.

(a) An iron droplet of radius $r = 5$ mm sinks through a magma ocean with density contrast $\Delta\rho = 4000 \text{ kg m}^{-3}$, gravity $g = 5 \text{ m s}^{-2}$, and melt viscosity $\mu = 0.1 \text{ Pa s}$. Compute its Stokes settling velocity (show that $v \approx 1.11 \text{ m s}^{-1}$) and the time to cross a magma ocean 1500 km deep, in days. State also why this crossing time is a lower limit.

Solution.

$$v = \frac{2}{9} \frac{\Delta\rho g r^2}{\mu} = \frac{2}{9} \times \frac{4000 \times 5 \times (5 \times 10^{-3})^2}{0.1} = \frac{2}{9} \times \frac{2 \times 10^4 \times 2.5 \times 10^{-5}}{0.1} = \frac{2}{9} \times 5.0 = 1.1111 \text{ m s}^{-1},$$

matching the checkpoint. The crossing time:

$$t = \frac{1.5 \times 10^6}{1.111} = 1.3501 \times 10^6 \text{ s} = \frac{1.3501 \times 10^6}{86400} = 15.6 \text{ d.}$$

| $v \approx 1.11 \text{ m s}^{-1}$; the droplet crosses 1500 km in about 16 days. The time is a lower limit: at 5 mm the flow around the droplet is already turbulent (the notes place the laminar limit near 1 mm), so the true settling is slower and the crossing takes weeks rather than days.

(b) Let x be the core mass fraction and $D = C^{\text{metal}}/C^{\text{silicate}}$ the metal-silicate partition coefficient of a trace element. Assuming the whole planet equilibrated once, show from mass conservation that the mantle retains a fraction

$$\frac{C^{\text{silicate}}}{C^{\text{bulk}}} = \frac{1}{1 + x(D - 1)}$$

of the bulk concentration. Evaluate it for iridium with $D = 10^4$ and $x = 0.32$ (show that the fraction is $\approx 3.1 \times 10^{-4}$).

Solution. Per unit planet mass, the element is conserved:

$$C^{\text{bulk}} = x C^{\text{metal}} + (1 - x) C^{\text{silicate}} = C^{\text{silicate}} [xD + (1 - x)] = C^{\text{silicate}} [1 + x(D - 1)],$$

using $C^{\text{metal}} = D C^{\text{silicate}}$. Inverting gives the stated fraction. For iridium:

$$x(D - 1) = 0.32 \times 9999 = 3200, \quad \frac{C^{\text{silicate}}}{C^{\text{bulk}}} = \frac{1}{3201} = 3.124 \times 10^{-4}.$$

| The mantle should retain only $\approx 3.1 \times 10^{-4}$ of its bulk iridium; with the larger D values in the notes' range (10^4 – 10^6) the predicted remainder is smaller still.

(c) The observed iridium depletion of Earth's mantle is a factor ~ 1000 , i.e. the mantle retains $\sim 10^{-3}$ of chondritic. Compare with your prediction from part (b) and explain, in 2 to 3 sentences, what the discrepancy implies. What single observation about the element ratios clinches the interpretation?

Solution. The mantle holds at least 3 times more iridium than one-stage equilibration predicts (and orders of magnitude more for the higher D values in the notes' range). The standard resolution is the late veneer: after core formation ended, Earth accreted a final $\sim 0.5\%$ of its mass in chondritic material whose siderophile elements stayed

in the mantle, because no sinking metal remained to extract them. The clinching observation is that the highly siderophile elements sit in *chondritic proportions* to one another: core extraction would have fractionated them (their D values differ), whereas a late chondritic addition preserves the ratios.

The excess records a late veneer of ~ 0.5% Earth masses of chondritic material after core formation; the unfractionated (chondritic) element ratios are the fingerprint.

(d) A planetesimal's mantle shows a tungsten excess $\epsilon^{182}\text{W} = +20$ with an Hf/W enrichment factor $(\text{Hf}/\text{W})/(\text{Hf}/\text{W})_{\text{CHUR}} = 26$, so the bracketed factor in the notes' two-stage equation is 25. Using the Hf-W constants from the data box, show that $e^{-\lambda t_c} = 0.784$ and find the two-stage core-formation age t_c . Compare with the accretion-timeline figure in the Lecture 4 notes.

Solution. The two-stage equation rearranges to

$$e^{-\lambda t_c} = \frac{\epsilon^{182}\text{W}}{Q_{\text{W}} \times 25 \times ({}^{182}\text{Hf}/{}^{180}\text{Hf})_0} = \frac{20}{10^4 \times 25 \times 1.02 \times 10^{-4}} = \frac{20}{25.5} = 0.784,$$

the printed checkpoint. Solving for t_c :

$$t_c = -\frac{\ln 0.784}{\lambda} = \frac{0.24335}{7.79 \times 10^{-2}} = 3.1238 \approx 3.1 \text{ Myr}.$$

This sits squarely in the planetesimal band of the accretion-timeline figure, which shows planetesimal cores differentiating within the first few Myr, well before the planets finish growing.

$t_c \approx 3.1$ Myr: this body formed its core in the planetesimal era.

(e) In 2 to 3 sentences: why can the Hf-W chronometer not date any core-forming event later than roughly 45 Myr after CAIs, and what tool takes over for such late events?

Solution. Beyond ~5 half-lives (≈ 45 Myr) the parent ${}^{182}\text{Hf}$ is effectively extinct: whatever ${}^{182}\text{W}$ exists is already made and distributed, so a later core extraction changes nothing in the mantle's $\epsilon^{182}\text{W}$ and the clock reads nothing. Late events (such as the nucleation of Earth's inner core, billions of years after formation) are instead constrained by thermal-evolution modelling, as in the notes.

After ~45 Myr the parent isotope is extinct and the chronometer is blind; thermal-evolution modelling takes over.

Problem 5 Dynamos and shields

A convecting metallic core can maintain a magnetic field, and that field carves a cavity in the solar wind. Here you test both, for two very different bodies.

(a) Start from pressure balance at the dayside magnetopause, $\frac{1}{2}\rho_{\text{sw}}v_{\text{sw}}^2 = B_{\text{mp}}^2/(2\mu_0)$, with a dipole field $B(r) = B_0(R_p/r)^3$. Show that the standoff distance obeys

$$r_{\text{mp}} = R_p \left(\frac{B_0^2}{\mu_0 \rho_{\text{sw}} v_{\text{sw}}^2} \right)^{1/6}.$$

During a coronal mass ejection the solar-wind density rises tenfold at fixed speed. Starting from Earth's typical $r_{\text{mp}} = 10 R_{\oplus}$, show that the magnetopause moves in to $\approx 6.8 R_{\oplus}$.

Solution. Insert the dipole law at $r = r_{\text{mp}}$ into the pressure balance:

$$\frac{1}{2} \rho_{\text{sw}} v_{\text{sw}}^2 = \frac{B_0^2 (R_p / r_{\text{mp}})^6}{2\mu_0} \Rightarrow \left(\frac{r_{\text{mp}}}{R_p} \right)^6 = \frac{B_0^2}{\mu_0 \rho_{\text{sw}} v_{\text{sw}}^2},$$

which is the stated result after taking the sixth root. The standoff therefore scales as $\rho_{\text{sw}}^{-1/6}$ at fixed B_0 and v_{sw} :

$$r'_{\text{mp}} = 10 R_{\oplus} \times 10^{-1/6} = \frac{10}{1.4678} R_{\oplus} = 6.813 R_{\oplus} \approx 6.8 R_{\oplus}.$$

| $r_{\text{mp}} \propto (B_0^2 / \rho_{\text{sw}} v_{\text{sw}}^2)^{1/6}$; a tenfold density surge compresses Earth's magnetopause from 10 to $\approx 6.8 R_{\oplus}$. The sixth root is why the magnetosphere is so stable against solar-wind buffeting.

(b) Ganymede's dynamo operates in a liquid iron shell of thickness $L \approx 300$ km. Taking the convective speed to be the same as in Earth's core, $U = 5 \times 10^{-4} \text{ m s}^{-1}$, and magnetic diffusivity $\eta = 1 \text{ m}^2 \text{ s}^{-1}$, compute the magnetic Reynolds number $\text{Rm} = UL/\eta$ (show that $\text{Rm} = 150$) and judge dynamo feasibility against the critical range $\text{Rm}_c \sim 10\text{--}100$.

Solution.

$$\text{Rm} = \frac{5 \times 10^{-4} \times 3.0 \times 10^5}{1} = 150,$$

matching the checkpoint. This clears the upper end of the critical range: even a shell only 300 km thick can sustain a dynamo at Earth-like convective speeds, consistent with the observed $\sim 0.7 \mu\text{T}$ surface field. The margin is modest, though: a few times slower convection would put Ganymede below threshold, which is why its dynamo is considered surprising and its driving (compositional buoyancy, possibly tidally modulated) matters.

| $\text{Rm} = 150 \geq \text{Rm}_c$: feasible, with little margin.

(c) If Ganymede's convection stopped, its field would decay ohmically on the timescale $\tau_{\text{ohm}} = L^2/\eta \approx 3 \times 10^3 \text{ yr}$ (verify this in one line if you wish). Explain in 2 to 3 sentences what this timescale, set against the age of the solar system, proves about the field the Galileo spacecraft measured in 1996, and why a "fossil field" interpretation fails.

Solution. The one-line check: $\tau_{\text{ohm}} = (3.0 \times 10^5)^2 / 1 = 9.0 \times 10^{10} \text{ s} = 9.0 \times 10^{10} / 3.156 \times 10^7 = 2852 \text{ yr}$. Against 4.57 Gyr this is an instant: any field created in Ganymede's past would have decayed away millions of times over. A fossil field therefore cannot survive; the field Galileo measured must be regenerated continuously, so the 1996 detection is direct proof of a dynamo operating now.

| $\tau_{\text{ohm}} \approx 3 \times 10^3 \text{ yr} \ll 4.57 \text{ Gyr}$: no fossil field survives, so the measured field proves a currently running dynamo.

(d) Judge this claim in 2 to 3 sentences: "Venus has no magnetic field because it rotates too slowly."

Solution. Not supported. Numerical dynamo simulations show that slow rotation alone does not prevent dynamo action, and the scaling laws for field strength are essentially independent of rotation rate. The likely culprits are the absence of a crystallising inner core (no compositional buoyancy) and Venus's stagnant lid, which insulates the core and throttles the convective power available to a dynamo; rotation is at most a secondary factor.

False as stated: the field is starved of convective power, not of rotation.