

A vibrant green aurora australis (southern lights) is visible against a dark blue night sky. The aurora appears as a bright, glowing band of light that curves across the upper right portion of the frame. The background is a deep, dark blue, suggesting a clear night sky.

Lecture 4: Chemical Differentiation & Magnetospheres

Planetary Systems

Tim Lichtenberg

Kapteyn Astronomical Institute
University of Groningen

2026

Learning objectives

By the end of this lecture, you will be able to:


- Explain metal–silicate separation and core formation
- Use siderophile element depletion as evidence for differentiation
- Describe the Hf–W chronometer for dating core formation
- Derive the magnetic Reynolds number and assess dynamo feasibility
- Compare magnetic fields across the solar system

Recap: Lecture 3


- Four heat sources: accretional, differentiation, radioactive, tidal
- Conduction too slow for planets: $\tau \sim L^2/\kappa$
- Rayleigh number: $Ra \sim 10^7\text{--}10^8$ for Earth $\Rightarrow$ vigorous convection
- Plate tectonics (mobile lid) vs. stagnant lid
- Tidal heating: Io, Enceladus, Europa

Today: The *chemical* consequences of melting, and how differentiation enables magnetic fields.

1. Accretion, melting, & core formation



Metal-silicate separation




Credit: Wikimedia, CC BY-SA 4.0

- Accretional heating + $^{26}\text{Al} \rightarrow$ **magma ocean**
- Iron ($\rho \sim 7000 \text{ kg m}^{-3}$) is immiscible with silicate ($\rho \sim 3000 \text{ kg m}^{-3}$)
- Iron droplets sink through silicate melt
- **Stokes settling velocity:**


$$v_{\text{Stokes}} = \frac{2}{9} \frac{\Delta \rho g r^2}{\mu}$$

- For cm-sized droplets: $v \sim 4 \text{ m s}^{-1}$
- Traverses magma ocean in **days**



Stokes settling of iron droplets in a silicate magma ocean. Course figure.

Reading the settling-velocity plot



- The line is $v \propto r^2$ on log-log axes (slope 2).
- Markers: 1 mm at 0.044 m s⁻¹, 1 cm at 4.4 m s⁻¹; left of the dashed line ($Re \approx 1$), Stokes' law holds exactly.
- To its right the flow is turbulent and the true speed at 1 cm is nearer 0.5 m s⁻¹, still crossing a 1000 km magma ocean within weeks.

Iron rain empties a magma ocean of its metal within weeks: on geological clocks, core formation is instantaneous.

Why the magma ocean stage is essential

Metal–silicate separation needs a **liquid** mantle.

- **Liquid mantle:** iron droplets sink at m s^{-1} , a core forms within weeks
- **Solid mantle:** rock flows so slowly ($\mu \sim 10^{21} \text{ Pa s}$, 22 orders of magnitude higher than silicate melt) that even a km-sized iron body would need longer than the age of the universe to sink to the centre

Core formation happens while the planet is molten, or not at all; chondritic asteroids that never melted never formed cores.

The magma ocean state

After accretion, the outer few thousand kilometres of a terrestrial planet are molten.


- Depth: several hundred to >2000 km (whole-mantle for large impacts)
- Surface temperature: 2000–4000 K depending on atmosphere
- Viscosity: 10^{-1} – 10^0 Pa s (like motor oil, for molten silicate)
- Convection is **turbulent** (Rayleigh number $\gtrsim 10^{25}$)
- Cooling is set by surface radiation and greenhouse blanketing

Lifetime of a magma ocean is 10^3 – 10^8 yr: long enough for core formation, short compared to planetary evolution.

The Moon-forming giant impact

- Mars-sized body (Theia) collided with proto-Earth ~4.5 Ga
- Re-melted most of Earth's mantle, vaporised part of it
- Created the final deep magma ocean
- Established present core-mantle structure
- Debris disk → Moon (confirmed by lunar $\epsilon^{182}\text{W}$ matching Earth's mantle)

Core formation is not a single event but a series of episodes, culminating in the giant impact.



SPH simulation of the Moon-forming giant impact: (a) Mars-sized Theia approaches the proto-Earth, (b) contact and tidal disruption, (c) part of the debris falls back while outer clumps survive, (d) a satellite on a wide orbit. Kegerreis et al. (2022).

► Play animation

► Source video


Giant impacts that remove mantles

Not every giant impact ends in net accretion: an oblique, fast collision can strip more silicate than it delivers.

- **Mercury:** core mass fraction 65–70% (Earth ~32%, Mars ~24%); a hit-and-run impact (an oblique collision whose remnant re-accretes rather than merging) stripped ~50–70% of the pre-impact mantle
- **Mars:** the hemispheric dichotomy (~6 km elevation, ~30 km crustal-thickness contrast) is the relaxed scar of a Borealis-scale impact; crust moved sideways, not lost
- **Earth–Moon:** the opposite endpoint, net accretion of the impactor


Mercury and Earth are the two endpoints of the same process: giant impacts set the final core-to-mantle ratio.

Source: Benz et al. (2007); Chau et al. (2018); Andrews-Hanna et al. (2008); Marinova et al. (2008)




SPH (particle-based hydrodynamics) simulation of a mantle-stripping giant impact on proto-Mercury. Chau et al. (2018).

Reading the Mercury impact frames



- A proto-Mercury of 2.25 present Mercury masses (red core, yellow mantle) is hit by an impactor of half its mass (turquoise core, blue mantle) at impact parameter $b = 0.5$ and $v = 30 \text{ km s}^{-1}$; the white bar marks $1 R_{\oplus}$.
- Silicate mantle is preferentially ejected while the metal-rich remnant recollapses.
- After 53 hours the largest fragment holds 0.95 of Mercury's present mass with iron mass fraction $Z_{\text{Fe}} = 0.67$, matching the observed core mass fraction.

One oblique collision at several escape velocities converts an ordinary protoplanet into iron-rich Mercury.



Mars's hemispheric dichotomy in MOLA global topography: smooth, low northern plains (blue) against cratered southern highlands (orange to red); the deep blue basin on the right is Hellas. Credit: NASA/JPL.

The late veneer

Highly siderophile elements (Ir, Pt, Os, Au) are **overabundant** in Earth's mantle compared to the deep-core equilibration prediction.

- Mantle abundances $\sim 200\times$ higher than expected for full metal extraction
- Relative ratios are **chondritic** (matching primitive meteorites), not fractionated
- Interpretation: an extra $\sim 0.5\%$ of Earth's mass accreted **after** core formation closed
- This “late veneer” delivered its siderophile elements to the mantle without equilibrating with the core

Source: Walker (2009); Day et al. (2016)

A smaller late veneer is also inferred for Mars and the Moon from their platinum-group element abundances.

Goldschmidt classification

1 1.008 H hydrogen	2 6.94 Li lithium	3 9.012 Be beryllium	4 12.01 B boron	5 10.81 C carbon	6 14.01 N nitrogen	7 16.00 O oxygen	8 19.00 F fluorine	9 20.18 Ne neon	10 4.003 He helium
11 22.99 Na sodium	12 24.31 Mg magnesium	13 26.98 Al aluminum	14 28.09 Si silicon	15 30.97 P phosphorus	16 32.07 S sulfur	17 35.45 Cl chlorine	18 39.95 Ar argon	19 39.09 K potassium	20 40.08 Ca calcium
21 39.09 K potassium	22 40.08 Ca calcium	23 44.96 Sc scandium	24 47.88 Ti titanium	25 50.94 V vanadium	26 52.00 Cr chromium	27 54.94 Mn manganese	28 55.85 Fe iron	29 58.93 Co cobalt	30 58.93 Ni nickel
31 69.72 Ga gallium	32 72.64 Ge germanium	33 74.92 As arsenic	34 78.96 Se selenium	35 79.90 Br bromine	36 83.80 Kr krypton	37 85.47 Rb rubidium	38 87.62 Sr strontium	39 88.91 Y yttrium	40 90.92 Zr zirconium
41 132.91 Cs cesium	42 132.91 Ba barium	43 137.33 La lanthanum	44 138.91 Ce cerium	45 140.12 Pr praseodymium	46 140.91 Nd neodymium	47 143.92 Pm promethium	48 150.36 Sm samarium	49 151.96 Eu europium	50 157.25 Gd gadolinium
51 132.91 Cs cesium	52 132.91 Ba barium	53 137.33 La lanthanum	54 138.91 Ce cerium	55 140.12 Pr praseodymium	56 140.91 Nd neodymium	57 143.92 Pm promethium	58 150.36 Sm samarium	59 151.96 Eu europium	60 157.25 Gd gadolinium
61 132.91 Cs cesium	62 132.91 Ba barium	63 137.33 La lanthanum	64 138.91 Ce cerium	65 140.12 Pr praseodymium	66 140.91 Nd neodymium	67 143.92 Pm promethium	68 150.36 Sm samarium	69 151.96 Eu europium	70 157.25 Gd gadolinium
71 132.91 Cs cesium	72 132.91 Ba barium	73 137.33 La lanthanum	74 138.91 Ce cerium	75 140.12 Pr praseodymium	76 140.91 Nd neodymium	77 143.92 Pm promethium	78 150.36 Sm samarium	79 151.96 Eu europium	80 157.25 Gd gadolinium
81 132.91 Cs cesium	82 132.91 Ba barium	83 137.33 La lanthanum	84 138.91 Ce cerium	85 140.12 Pr praseodymium	86 140.91 Nd neodymium	87 143.92 Pm promethium	88 150.36 Sm samarium	89 151.96 Eu europium	90 157.25 Gd gadolinium
91 132.91 Cs cesium	92 132.91 Ba barium	93 137.33 La lanthanum	94 138.91 Ce cerium	95 140.12 Pr praseodymium	96 140.91 Nd neodymium	97 143.92 Pm promethium	98 150.36 Sm samarium	99 151.96 Eu europium	100 157.25 Gd gadolinium
101 132.91 Cs cesium	102 132.91 Ba barium	103 137.33 La lanthanum	104 138.91 Ce cerium	105 140.12 Pr praseodymium	106 140.91 Nd neodymium	107 143.92 Pm promethium	108 150.36 Sm samarium	109 151.96 Eu europium	110 157.25 Gd gadolinium
111 132.91 Cs cesium	112 132.91 Ba barium	113 137.33 La lanthanum	114 138.91 Ce cerium	115 140.12 Pr praseodymium	116 140.91 Nd neodymium	117 143.92 Pm promethium	118 150.36 Sm samarium	119 151.96 Eu europium	120 157.25 Gd gadolinium
121 132.91 Cs cesium	122 132.91 Ba barium	123 137.33 La lanthanum	124 138.91 Ce cerium	125 140.12 Pr praseodymium	126 140.91 Nd neodymium	127 143.92 Pm promethium	128 150.36 Sm samarium	129 151.96 Eu europium	130 157.25 Gd gadolinium
131 132.91 Cs cesium	132 132.91 Ba barium	133 137.33 La lanthanum	134 138.91 Ce cerium	135 140.12 Pr praseodymium	136 140.91 Nd neodymium	137 143.92 Pm promethium	138 150.36 Sm samarium	139 151.96 Eu europium	140 157.25 Gd gadolinium
141 132.91 Cs cesium	142 132.91 Ba barium	143 137.33 La lanthanum	144 138.91 Ce cerium	145 140.12 Pr praseodymium	146 140.91 Nd neodymium	147 143.92 Pm promethium	148 150.36 Sm samarium	149 151.96 Eu europium	150 157.25 Gd gadolinium
151 132.91 Cs cesium	152 132.91 Ba barium	153 137.33 La lanthanum	154 138.91 Ce cerium	155 140.12 Pr praseodymium	156 140.91 Nd neodymium	157 143.92 Pm promethium	158 150.36 Sm samarium	159 151.96 Eu europium	160 157.25 Gd gadolinium
161 132.91 Cs cesium	162 132.91 Ba barium	163 137.33 La lanthanum	164 138.91 Ce cerium	165 140.12 Pr praseodymium	166 140.91 Nd neodymium	167 143.92 Pm promethium	168 150.36 Sm samarium	169 151.96 Eu europium	170 157.25 Gd gadolinium
171 132.91 Cs cesium	172 132.91 Ba barium	173 137.33 La lanthanum	174 138.91 Ce cerium	175 140.12 Pr praseodymium	176 140.91 Nd neodymium	177 143.92 Pm promethium	178 150.36 Sm samarium	179 151.96 Eu europium	180 157.25 Gd gadolinium
181 132.91 Cs cesium	182 132.91 Ba barium	183 137.33 La lanthanum	184 138.91 Ce cerium	185 140.12 Pr praseodymium	186 140.91 Nd neodymium	187 143.92 Pm promethium	188 150.36 Sm samarium	189 151.96 Eu europium	190 157.25 Gd gadolinium
191 132.91 Cs cesium	192 132.91 Ba barium	193 137.33 La lanthanum	194 138.91 Ce cerium	195 140.12 Pr praseodymium	196 140.91 Nd neodymium	197 143.92 Pm promethium	198 150.36 Sm samarium	199 151.96 Eu europium	200 157.25 Gd gadolinium
201 132.91 Cs cesium	202 132.91 Ba barium	203 137.33 La lanthanum	204 138.91 Ce cerium	205 140.12 Pr praseodymium	206 140.91 Nd neodymium	207 143.92 Pm promethium	208 150.36 Sm samarium	209 151.96 Eu europium	210 157.25 Gd gadolinium
211 132.91 Cs cesium	212 132.91 Ba barium	213 137.33 La lanthanum	214 138.91 Ce cerium	215 140.12 Pr praseodymium	216 140.91 Nd neodymium	217 143.92 Pm promethium	218 150.36 Sm samarium	219 151.96 Eu europium	220 157.25 Gd gadolinium
221 132.91 Cs cesium	222 132.91 Ba barium	223 137.33 La lanthanum	224 138.91 Ce cerium	225 140.12 Pr praseodymium	226 140.91 Nd neodymium	227 143.92 Pm promethium	228 150.36 Sm samarium	229 151.96 Eu europium	230 157.25 Gd gadolinium
231 132.91 Cs cesium	232 132.91 Ba barium	233 137.33 La lanthanum	234 138.91 Ce cerium	235 140.12 Pr praseodymium	236 140.91 Nd neodymium	237 143.92 Pm promethium	238 150.36 Sm samarium	239 151.96 Eu europium	240 157.25 Gd gadolinium
241 132.91 Cs cesium	242 132.91 Ba barium	243 137.33 La lanthanum	244 138.91 Ce cerium	245 140.12 Pr praseodymium	246 140.91 Nd neodymium	247 143.92 Pm promethium	248 150.36 Sm samarium	249 151.96 Eu europium	250 157.25 Gd gadolinium
251 132.91 Cs cesium	252 132.91 Ba barium	253 137.33 La lanthanum	254 138.91 Ce cerium	255 140.12 Pr praseodymium	256 140.91 Nd neodymium	257 143.92 Pm promethium	258 150.36 Sm samarium	259 151.96 Eu europium	260 157.25 Gd gadolinium
261 132.91 Cs cesium	262 132.91 Ba barium	263 137.33 La lanthanum	264 138.91 Ce cerium	265 140.12 Pr praseodymium	266 140.91 Nd neodymium	267 143.92 Pm promethium	268 150.36 Sm samarium	269 151.96 Eu europium	270 157.25 Gd gadolinium
271 132.91 Cs cesium	272 132.91 Ba barium	273 137.33 La lanthanum	274 138.91 Ce cerium	275 140.12 Pr praseodymium	276 140.91 Nd neodymium	277 143.92 Pm promethium	278 150.36 Sm samarium	279 151.96 Eu europium	280 157.25 Gd gadolinium
281 132.91 Cs cesium	282 132.91 Ba barium	283 137.33 La lanthanum	284 138.91 Ce cerium	285 140.12 Pr praseodymium	286 140.91 Nd neodymium	287 143.92 Pm promethium	288 150.36 Sm samarium	289 151.96 Eu europium	290 157.25 Gd gadolinium
291 132.91 Cs cesium	292 132.91 Ba barium	293 137.33 La lanthanum	294 138.91 Ce cerium	295 140.12 Pr praseodymium	296 140.91 Nd neodymium	297 143.92 Pm promethium	298 150.36 Sm samarium	299 151.96 Eu europium	300 157.25 Gd gadolinium
301 132.91 Cs cesium	302 132.91 Ba barium	303 137.33 La lanthanum	304 138.91 Ce cerium	305 140.12 Pr praseodymium	306 140.91 Nd neodymium	307 143.92 Pm promethium	308 150.36 Sm samarium	309 151.96 Eu europium	310 157.25 Gd gadolinium
311 132.91 Cs cesium	312 132.91 Ba barium	313 137.33 La lanthanum	314 138.91 Ce cerium	315 140.12 Pr praseodymium	316 140.91 Nd neodymium	317 143.92 Pm promethium	318 150.36 Sm samarium	319 151.96 Eu europium	320 157.25 Gd gadolinium
321 132.91 Cs cesium	322 132.91 Ba barium	323 137.33 La lanthanum	324 138.91 Ce cerium	325 140.12 Pr praseodymium	326 140.91 Nd neodymium	327 143.92 Pm promethium	328 150.36 Sm samarium	329 151.96 Eu europium	330 157.25 Gd gadolinium
331 132.91 Cs cesium	332 132.91 Ba barium	333 137.33 La lanthanum	334 138.91 Ce cerium	335 140.12 Pr praseodymium	336 140.91 Nd neodymium	337 143.92 Pm promethium	338 150.36 Sm samarium	339 151.96 Eu europium	340 157.25 Gd gadolinium
341 132.91 Cs cesium	342 132.91 Ba barium	343 137.33 La lanthanum	344 138.91 Ce cerium	345 140.12 Pr praseodymium	346 140.91 Nd neodymium	347 143.92 Pm promethium	348 150.36 Sm samarium	349 151.96 Eu europium	350 157.25 Gd gadolinium
351 132.91 Cs cesium	352 132.91 Ba barium	353 137.33 La lanthanum	354 138.91 Ce cerium	355 140.12 Pr praseodymium	356 140.91 Nd neodymium	357 143.92 Pm promethium	358 150.36 Sm samarium	359 151.96 Eu europium	360 157.25 Gd gadolinium
361 132.91 Cs cesium	362 132.91 Ba barium	363 137.33 La lanthanum	364 138.91 Ce cerium	365 140.12 Pr praseodymium	366 140.91 Nd neodymium	367 143.92 Pm promethium	368 150.36 Sm samarium	369 151.96 Eu europium	370 157.25 Gd gadolinium
371 132.91 Cs cesium	372 132.91 Ba barium	373 137.33 La lanthanum	374 138.91 Ce cerium	375 140.12 Pr praseodymium	376 140.91 Nd neodymium	377 143.92 Pm promethium	378 150.36 Sm samarium	379 151.96 Eu europium	380 157.25 Gd gadolinium
381 132.91 Cs cesium	382 132.91 Ba barium	383 137.33 La lanthanum	384 138.91 Ce cerium	385 140.12 Pr praseodymium	386 140.91 Nd neodymium	387 143.92 Pm promethium	388 150.36 Sm samarium	389 151.96 Eu europium	390 157.25 Gd gadolinium
391 132.91 Cs cesium	392 132.91 Ba barium	393 137.33 La lanthanum	394 138.91 Ce cerium	395 140.12 Pr praseodymium	396 140.91 Nd neodymium	397 143.92 Pm promethium	398 150.36 Sm samarium	399 151.96 Eu europium	400 157.25 Gd gadolinium
401 132.91 Cs cesium	402 132.91 Ba barium	403 137.33 La lanthanum	404 138.91 Ce cerium	405 140.12 Pr praseodymium	406 140.91 Nd neodymium	407 143.92 Pm promethium	408 150.36 Sm samarium	409 151.96 Eu europium	410 157.25 Gd gadolinium
411 132.91 Cs cesium	412 132.91 Ba barium	413 137.33 La lanthanum	414 138.91 Ce cerium	415 140.12 Pr praseodymium	416 140.91 Nd neodymium	417 143.92 Pm promethium	418 150.36 Sm samarium	419 151.96 Eu europium	420 157.25 Gd gadolinium
421 132.91 Cs cesium	422 132.91 Ba barium	423 137.33 La lanthanum	424 138.91 Ce cerium	425 140.12 Pr praseodymium	426 140.91 Nd neodymium	427 14			

Metal–silicate partition coefficients

How strongly does an element prefer metal vs. silicate?

$$D_i^{\text{met/sil}} \equiv \frac{C_i^{\text{metal}}}{C_i^{\text{silicate}}}$$

Element	$D^{\text{met/sil}}$	Behaviour
Ir, Os, Pt	10^4 – 10^6	Highly siderophile
W, Mo	10^2 – 10^4	Moderately siderophile
Ni, Co	10–100	Weakly siderophile
Fe	~10	Weakly siderophile
Si, Mg, O	$\ll 1$	Lithophile

Source: Rubie et al. (2015); McDonough (2003)

Earth's mantle is depleted ~1000× in Ir and Pt: direct evidence that iron-loving elements followed the metal into the core.


Why silicon is in the core

Partition coefficients depend strongly on pressure, temperature, and oxygen fugacity (an effective O_2 partial pressure setting the melt's redox state).

- At low pressure (<10 GPa): Si is strongly lithophile, $D_{Si}^{met/sil} \ll 1$
- At 40–60 GPa (deep magma ocean base): $D_{Si}^{met/sil}$ rises to ~1
- This implies Earth's core contains **several wt%** silicon
- Si in core helps explain the ~5–10% core density deficit (Lecture 8)
- Not the only candidate: S, O, C, and H are invoked alongside Si, in proportions set by the redox conditions of core formation


Deep magma ocean partitioning (40–60 GPa) is the standard model for Earth's core composition, matching observed siderophile element depletions.

Source: Rubie et al. (2015); Fischer et al. (2015); reviews: Hirose et al. (2021); Shahar et al. (2026)



Metal-silicate partition coefficients versus pressure. Course figure, after Siebert et al. (2013).

Reading the partitioning experiments



- Curves: laboratory $D^{\text{met/sil}}$ for Ni and Co at magma-ocean conditions. At low pressure $D \sim 10^3$ – 10^4 : the mantle should keep almost no Ni or Co.
- Grey band: $D \approx 24$ – 26 , required by the Ni and Co content actually observed in the mantle.
- Both curves reach the band only near 40–60 GPa (blue band): equilibration at 54 ± 5 GPa, the base of a 1000–1500 km deep magma ocean (Fischer et al. 2015).

The mantle's siderophile depletions remember the depth of the magma ocean in which the core formed.

Hf–W chronometry: when did cores form?

- $^{182}\text{Hf} \rightarrow ^{182}\text{W}$, half-life = 8.9 Myr
- Hf is **lithophile** (stays in mantle)
- W is **siderophile** (extracted to core)
- Early core formation $\rightarrow$ excess ^{182}W in mantle

Body	$\epsilon^{182}\text{W}$	Core formation timing
Earth	+2	two-stage age ~30 Myr (lower limit)
Mars	+0.4 (bulk)	complete within ~10 Myr
Moon	~Earth	Consistent with giant impact

Sources: Kleine et al. (2009); Nimmo & Kleine (2015); Kleine & Walker (2017)

How the Hf–W clock works

Two bodies diverging from chondritic composition:

Early core formation (Mars):

- Core extracts all W, stops producing ^{182}W
- Mantle keeps Hf, continues making ^{182}W
- Result: a clear mantle excess ($\epsilon \approx +0.4$ bulk; up to +1.8 in depleted shergottites, a martian meteorite class)


Late core formation (hypothetical):

- ^{182}Hf already decayed
- No divergence between core and mantle ^{182}W
- Mantle would have $\epsilon^{182}\text{W} \approx 0$ (chondritic)

Earth's moderate excess ($+2\epsilon$) gives a two-stage age of ~30 Myr, a lower limit: the bulk of core formation finished within ~100 Myr (Nimmo & Kleine 2015; Kleine & Walker 2017).


(a) parent and daughter separate

mantle: Hf (lithophile) stays




W (siderophile)
sinks into the core

(b) earlier core formation, larger excess




(a) Core formation separates parent (Hf, mantle) from daughter reservoir (W, core). (b) Mantle ^{182}W excess vs. time of core formation, two-stage model with Earth's mantle Hf/W: the measured $+2\epsilon$ gives ~ 28 Myr. Course figure.



(a) The first 5 Myr after CAI condensation (CAIs: calcium-aluminium-rich inclusions, the first nebular solids, define $t = 0$). (b) Core formation to 300 Myr. Course figure, after Lichtenberg et al. (2023); ages from Nimmo & Kleine (2015), Kleine & Walker (2017), Kruijer et al. (2017).

Reading the accretion timeline



- Panel (a): the meteoritic record, chondrules and planetesimals within ~4 Myr (details in Lecture 12).
- Panel (b), our focus here: planetesimal cores (iron meteorites) by ~1–3 Myr; Mars within ~10 Myr.
- Earth: two-stage age ~30 Myr is a lower limit, done within ~100 Myr; Moon impact after ~50 Myr.


Small bodies differentiate within 1–3 Myr; Earth needs tens of Myr: radiometric clocks resolve the whole hierarchy.

Magma ocean crystallisation

Core formation is not the end. As the magma ocean cools:


- The convecting melt follows an **adiabat**; the peridotite **solidus** and **liquidus** rise more steeply with pressure
- The cooling adiabat therefore first drops below the melting curves at the **base** of the mantle
- Crystallisation starts at the bottom; the crystallisation front migrates **upward**
- Deep mantle: high-pressure minerals such as **bridgmanite** (MgSiO_3) crystallise first
- Final residual melt at the top: enriched in incompatible elements
- An old idea: Kelvin (1862) already argued that a once-molten Earth solidifies from great depth upward, because pressure raises the melting point

After Elkins-Tanton (2012), *Annu. Rev. Earth Planet. Sci.*



Peridotite solidus and liquidus versus magma-ocean adiabats at successive times t_1 – t_4 .
Course figure, after Elkins-Tanton (2012).

Reading the crystallisation diagram



- The melting curves rise more steeply with pressure than the adiabats t_1 – t_4 ; between them, crystals and melt coexist (mush).
- First crystals where the adiabat meets the liquidus: at the base (t_2 , open circle).
- The adiabat–solidus intersection (dots) is the crystallisation front: it sits shallower at each later time.
- Caveat: dense Fe-rich melt can survive at the base as a **basal magma ocean** (early Earth: Labrosse et al. 2007; Mars: Samuel et al. 2023).

Deep magma oceans crystallise from the bottom up: the adiabat meets the melting curves at the base first.

Incompatible element enrichment

- Large ions (U^{4+} , Th^{4+} , K^{+}) do not fit well into crystal lattices
- Progressively concentrated in the shrinking residual liquid
- Eventually trapped in the **crust**
- Continental crust is ~50–100× enriched in U, Th, K vs. upper mantle
- Roughly **half** of Earth's radiogenic heat is produced in the crust, even though the crust is <1% of Earth's mass

Incompatible element enrichment is why Earth's crust is both radioactive (heat source) and geochemically distinct from the mantle below.

Mantle reservoirs

Magma ocean crystallisation and billions of years of processing leave distinct mantle reservoirs:

DMM Depleted MORB Mantle: upper mantle source of mid-ocean ridge basalts; depleted by partial melting and crustal extraction.

EM/OIB Enriched Mantle / Ocean Island Basalt source: deep plumes sample reservoirs with preserved incompatible elements; may represent primordial mantle or recycled oceanic crust.

HIMU “High- μ ” reservoir: high U/Pb, likely very old subducted oceanic crust.


Isotopic diversity of ocean island basalts (Hawaii, Iceland, Samoa) is direct evidence for long-lived mantle heterogeneity.

The lunar magma ocean: direct evidence

The Moon shows the clearest record of magma ocean differentiation:


- The concept was born from Apollo 11: anorthosite fragments in the Mare Tranquillitatis soil led Wood et al. (1970) to propose a deep molten shell
- **Anorthositic highlands:** low-density plagioclase feldspar floated to the top of the crystallising lunar magma ocean ~4.4 Ga
- Apollo samples of anorthosite date to 4.36–4.44 Ga (within 200 Myr of solar system formation)
- **KREEP** layer: the last dregs of crystallisation, rich in **K**, **R**are **E**arth **E**lements, **P**hosphorus; shallow under the nearside maria
- Lunar crust thickness varies from ~30 km (nearside) to ~70 km (farside)

The Moon is the textbook case of magma ocean differentiation; its exposed anorthosite confirms the theoretical crystallisation sequence.



End-state of the lunar magma ocean; thin layers exaggerated. Course figure, after Elkins-Tanton (2012).

Reading the lunar cross-section



- **Outer pale shell:** the 40–50 km anorthosite flotation crust; beneath it, the yellow urKREEP layer is the last ~1% of liquid to solidify.
- Light green: pyroxenite and dunite cumulates; the dark green ilmenite-rich layer formed last and is denser than everything below it.
- The ilmenite layer sinks (mantle overturn) in the first ~500 Myr; an iron core sits at the centre.

Apollo samples match this predicted stratigraphy in detail: the Moon is the clearest test case of magma-ocean theory.

Volatile delivery: classical vs. modern view


Classical picture:

- Inner solar system forms dry, inside the water snow line (radius beyond which water condenses as ice)
- Volatiles delivered **late** by comets and outer-belt asteroids


Revised picture (post-2010):

- All chondrite groups accreted some water ice; ice reached the inner disk via the early outward-sweeping snow line, grain reactions with nebular gas, and scattered icy bodies
- Jupiter's growth blocked the icy pebble drift and created the NC/CC isotopic bifurcation
- The growing protoplanets then dried out again: ^{26}Al melting, impact devolatilisation, and pebble ablation in primordial atmospheres
- Most of Earth's remaining water was already present at accretion, not delivered afterwards

(a) classical view: dry inner solar system, water arrives late




(b) revised view: ice arrives early, Jupiter splits the disk




Classical versus revised picture of volatile delivery. Course figure, after Alexander et al. (2019), Lichtenberg et al. (2021), Grewal et al. (2024).

Reading the volatile-delivery schematic

(a) classical view: dry inner solar system, water arrives late



(b) revised view: ice arrives early, Jupiter splits the disk



- (a) Classical: a fixed snow line keeps the inner disk dry; water arrives late, by asteroids and comets scattered inward.
- (b) Revised: the snow line sweeps outward, then retreats; the earliest inner planetesimals form beyond it and accrete ice (blue bodies).
- Jupiter's growth blocks the inward pebble drift and splits the disk into the NC and CC meteorite reservoirs.

When a planetesimal forms matters as much as where: timing sets its water budget.

Outgassing from the magma ocean

During magma ocean cooling, dissolved volatiles **outgas** into the overlying atmosphere.

Speciation depends on **oxygen fugacity** (f_{O_2}), set by the mantle redox state:

Reducing (low f_{O_2}): H_2 , CO , N_2 , CH_4

Oxidising (high f_{O_2}): H_2O , CO_2 , N_2 , SO_2

The oxidation state of the mantle at magma ocean solidification sets the initial atmosphere, which then conditions habitability (Lectures 5–6).

Volatile solubility in silicate melt


Solubility is a chemical property: each gas dissolves through its own reaction with the melt, and depends on melt composition and redox state.

- H₂O enters the melt as hydroxyl groups; CO₂ as carbonate ions, ~100× less soluble
- N barely dissolves when oxidised, forms soluble nitrides when reduced; S dissolves as sulfide or sulfate
- Water: each molecule splits into two OH groups, giving a square-root law

$$X_{\text{H}_2\text{O}} \propto p_{\text{H}_2\text{O}}^{1/2}$$


A thicker steam atmosphere lets the magma retain more water in solution: a self-limiting feedback, so a planet cannot outgas its entire water budget.

Source: Hirschmann (2012), *Earth Planet. Sci. Lett.*



Volatile partitioning between magma ocean and atmosphere vs. oxygen fugacity, computed with an equilibrium outgassing model (Nicholls et al. 2024) for the bulk silicate Earth inventory of Krijt et al. (2023).

Reading the outgassing diagram



- (a) The melt keeps almost the whole water inventory dissolved ($\sim 5 \times 10^{21}$ kg, over 3 present-day oceans) at all but the most reducing conditions.
- (b) Carbon is far less soluble: CO_2 and CO fill the atmosphere at hundreds to thousands of bar, while steam stays at a few bar.
- Redox controls speciation: H_2 , CO , CH_4 when reduced; CO_2 , then SO_2 , when oxidised.


During the magma ocean stage most of the planet's water sits in the magma, not in the atmosphere.

Impact erosion: two regimes

- **Local:** an ordinary impactor blows off at most the air above the tangent plane at the impact point: a fraction $H/2R \approx 6 \times 10^{-4}$ of Earth's atmosphere per event
- Erosion by ordinary impacts is therefore cumulative; km-scale planetesimals are the most efficient eroders per unit impactor mass
- **Global:** a giant impact shakes the whole planet; air is lost wherever the ground approaches v_{esc}
- Even the Moon-forming impact removes only ~20%; a surface ocean raises the loss substantially

No single impact strips an atmosphere: erosion is the integral of many small events plus rare giant shakes.

Source: Melosh & Vickery (1989); Schlichting et al. (2015); Genda & Abe (2003, 2005)



Atmospheric loss in a giant impact: ejecta plumes drive near-field loss at the impact site, while the shock breaking out on the far side drives far-field loss; green particles are lost.

Roche et al. (2025), CC BY 4.0.

Delivery versus erosion

- The same planetesimals that erode also deliver volatiles; the net budget depends on impactor sizes, velocities, and volatile content
- Venus keeps $\sim 50\times$ more ^{36}Ar than Earth: oceans on proto-Earth enhanced its loss, while dry proto-Venus kept its primordial noble gases
- Any primordial H_2 -He envelope died by attrition: repeated impacts plus XUV-driven thermal escape (Lecture 5)

Earth's air and oceans are the ledger of delivery minus erosion, closed by magma-ocean outgassing.

Source: Genda & Abe (2005); Schlichting et al. (2015)

Meteoritic dichotomy: NC vs. CC


Meteorites split into two isotopic groups: **non-carbonaceous (NC)** and **carbonaceous (CC)**.

- NC and CC have distinct ^{50}Ti , ^{54}Cr , ^{48}Ca ratios
- They reflect two separate reservoirs in the early solar nebula
- Interpretation: **Jupiter** formed early, blocking mixing between inner and outer disk
- Both reservoirs contributed to Earth's accretion, but in changing proportions over time

Source: Kruijer et al. (2017); Lichtenberg et al. (2021)

This dichotomy is one of the strongest meteoritic constraints on the timing of giant planet formation.

Break



2. Planetary dynamos



Magnetic fields and habitability

A magnetic field is more than a compass-maker. It shapes a planet's long-term viability.

- Deflects the solar wind $\Rightarrow$ reduces ion sputtering (atoms knocked loose by particle impacts) of the upper atmosphere
- Limits cosmic ray flux to the surface $\Rightarrow$ lower radiation dose
- Protects the ozone layer from charged-particle destruction
- On Venus, the lack of a field is one likely contributor to hydrogen escape
- On Mars, the loss of the field preceded the loss of ~ 0.8 bar of atmosphere (MAVEN)
- Debated: how essential is a global field for a habitable surface?

Source: Lammer et al. (2003); Jakosky et al. (2018)

Why magnetic fields matter

- **Atmospheric shielding:** deflects solar wind, reduces erosion
Mars lost its field ~4 Ga → lost most of its atmosphere
- **Surface habitability:** reduces radiation dose at the surface
- **Geological record:** magnetic minerals record field history (paleomagnetism)
- **Interior probe:** field properties constrain the deep interior

The induction equation

Magnetic fields in a conducting fluid obey:

$$\frac{\partial \vec{B}}{\partial t} = \underbrace{\nabla \times (\vec{v} \times \vec{B})}_{\text{advection}} + \underbrace{\eta \nabla^2 \vec{B}}_{\text{diffusion}}$$

- **Advection:** fluid motions amplify the field (create flux)
- **Diffusion:** ohmic resistance destroys the field (decay)
- $\eta = 1/(\mu_0 \sigma) = \text{magnetic diffusivity}$

Which term wins determines whether a dynamo can operate.

Three requirements for a dynamo

1. **Electrically conducting fluid**

Terrestrial planets: liquid iron alloy in the core

Gas giants: metallic hydrogen

2. **Convection**


Thermal buoyancy (core hotter than mantle)

Compositional buoyancy (inner core crystallisation releases light elements)

3. **Sufficient flow vigour**

Advection must dominate over diffusion

Quantified by the **magnetic Reynolds number** $R_m \gtrsim 10\text{--}100$

A photograph of the Aurora australis (Southern Lights) in a dark night sky. The aurora appears as a bright green, jagged, and elongated light structure. It has a main vertical column on the right side and a horizontal, diffuse band extending to the left. The background is a deep black with some faint, wispy green light trails.

3. Blackboard derivation: magnetic Reynolds number

Deriving the magnetic Reynolds number

Goal: From the induction equation, derive $R_m = UL/\eta$ and show that advection dominates in Earth's core.

Dimensional analysis of each term:

Advection: $|\nabla \times (\vec{v} \times \vec{B})| \sim UB/L$

Diffusion: $|\eta \nabla^2 \vec{B}| \sim \eta B/L^2$

The magnetic Reynolds number

$$R_m = \frac{\text{advection}}{\text{diffusion}} = \frac{UB/L}{\eta B/L^2}$$

The field strength B **cancels**:

$$R_m = \frac{UL}{\eta}$$

- $R_m \gg 1$: advection dominates, field “frozen in” to the flow
- $R_m \ll 1$: diffusion wins, field decays
- Critical value: $R_{m_c} \sim 10\text{--}100$

Key insight: R_m depends on the **flow**, not the field.

Application: Earth's outer core


Parameter	Symbol	Value
Core radius	L	$3.5 \times 10^6 \text{ m}$
Flow velocity	U	$5 \times 10^{-4} \text{ m s}^{-1}$
Magnetic diffusivity	η	$\sim 1 \text{ m}^2 \text{ s}^{-1}$

$$Rm_{\oplus} = \frac{5 \times 10^{-4} \times 3.5 \times 10^6}{1} \approx 1750$$

$Rm \approx 1750 \gg Rm_c$: advection overwhelmingly dominates.

Ohmic diffusion timescale: $\tau_{\text{ohm}} \sim L^2 / \eta \approx 400,000 \text{ yr} \ll \text{Earth's age} \Rightarrow \text{field must be **continuously regenerated**}.}$

Dynamo scaling laws




Credit: Course figure, after Christensen & Aubert (2006)

Dynamo simulations collapse onto simple power laws:

- Field strength $\propto$ (convective power) $^{1/3}$
- **Independent of rotation rate**
- Predicts Earth, Jupiter, Mercury, Ganymede from first principles
- Works across ~4 orders of magnitude in parameters

Planetary magnetic field strength is set by the **available convective power**, not by rotation rate or geometry.

Reading the dynamo scaling law



- Horizontal axis: a combination of convective buoyancy flux, shell thickness, and core density; vertical: the observed or simulated dipole field.
- Grey crosses illustrate the collapse of numerical dynamos onto one line over ~4 orders of magnitude.
- Earth, Jupiter, Saturn, Mercury, and Ganymede sit on the same regression within factors of order unity; Venus is absent, its dynamo off.

$B_{\text{dip}} \propto \sqrt{\rho \mu_0} (F_q D)^{1/3}$: field strength is bought with convective power, not rotation.

Hf-W chronometry: beyond the first estimate

Modern refinements of the core formation timescale.

- Pressure-dependent partitioning: $D_W^{\text{met/sil}}$ decreases at high P
- At deep magma ocean pressures (~60 GPa), D_W is ~10 instead of ~40 (surface)
- This widens the permitted window to ~30–100 Myr, depending on how much metal re-equilibrates
- Complementary isotope systems (Pd–Ag, Pt–Os) cross-check the timing
- Evidence for **disequilibrium core formation**: not all metal re-equilibrates with silicate before sinking
- Earth's core likely formed in episodic accretion events, not continuous

Source: Rudge et al. (2010); Kleine et al. (2017)

The Archean-Proterozoic paleomagnetic record

Geologists trace Earth's dynamo history through preserved magnetic signals.

- **Archean** (>2.5 Ga): paleomagnetic records in komatiites and banded iron formations
- Field has been **continuous** from ≥ 3.4 Ga to today (confirmed at multiple sites)
- **Proterozoic** (2.5–0.54 Ga): Apparent Polar Wander (APW) paths reveal plate motions and supercontinent cycles
- Paleointensity from Thellier experiments: early field strength similar to or slightly stronger than modern
- Reversal statistics: Archean reversal rate poorly constrained; modern rate set in the last ~500 Myr
- **Singularity puzzle**: some Proterozoic intervals show very frequent reversals followed by superchrons (long intervals with no reversal)

Source: Biggin et al. (2015); Tarduno et al. (2020)

How old is Earth's magnetic field?


Paleomagnetism records the field in ancient rocks.

- Oldest confirmed field: ~**3.4–3.5 Gyr** (Archean South Africa, Tarduno 2010)
- Single-crystal zircon paleomagnetism pushes this to **4.2 Gyr** (Tarduno et al. 2015)
- Implies Earth's dynamo has been continuously active for most of its history
- Constrains the core cooling rate and the earliest habitability of the surface


The dynamo was active during the Hadean eon, shielding Earth's atmosphere during the time when life was emerging.

Source: Tarduno et al. (2010, 2015), *Science*

4. Earth's dynamo & comparative magnetospheres



Earth's geodynamo



Credit: Wikimedia, CC BY-SA 4.0

- **Outer core:** liquid Fe alloy (1220–3480 km from centre)
- Two driving mechanisms:
 1. Thermal convection (core hotter than mantle)
 2. Compositional convection (inner core crystallisation)
- Predominantly **dipolar** field
- Tilted $\sim 11^\circ$ from rotation axis
- Surface field: 25–65 μT

Earth's core: composition & growth

Inner core (0–1220 km):


- Solid iron–nickel alloy
- Growing at ~0.5 mm/yr
- Age: ~0.5–1 Gyr (far younger than Earth)
- Latent heat of crystallisation powers the dynamo today

Outer core (1220–3480 km):


- Liquid Fe–Ni alloy + light elements
- ~5–10% less dense than pure Fe
- Light elements: S, Si, O, C, H
- Vigorous convection $\Rightarrow$ dynamo

Compositional convection (inner core crystallisation ejects light elements) is the **dominant** driver of today's geodynamo.


(a) thermal convection: the core cools



(b) compositional convection: the inner core grows




The two buoyancy sources of the geodynamo, in a quarter-section cutout (layer radii to scale): (a) the mantle extracts heat across the core-mantle boundary, so the core cools from above and convects; (b) inner-core growth rejects light elements (S, Si, O) and releases latent heat, driving compositional convection. Course figure.




Cross-section of Earth's interior. Credit: NASA/JPL-Caltech/SwRI (PIA25063).

(a) S waves: stopped by the liquid outer core




(b) P waves: refracted by the core




How *P* and *S* waves constrain the core: *S* waves stop at the liquid outer core (no arrivals beyond 103°); *P* waves are refracted into a shadow zone between 103° and 143°; PKIKP arrivals inside it reveal the solid inner core (phase letters name the ray legs: *P* mantle, *K* outer core, *I* inner core). Course figure.

The shadow-zone argument




- S waves cannot cross a **liquid**: no direct S beyond 103° $\Rightarrow$ the outer core is **liquid** (Oldham 1906)
- V_p drops $\sim 13.7 \rightarrow 8$ km/s at the CMB: rays refract steeply $\Rightarrow$ P shadow zone from 103° to 143°
- Refracted core phases (PKP) emerge beyond 143°
- Phase letters spell the ray path, one per leg: P mantle, K liquid outer core, I solid inner core
- Weak arrivals *inside* the shadow (PKIKP): refracted again by a **solid inner core** (Lehmann 1936)



PREM seismic velocities and density versus depth. Course figure, after Dziewonski & Anderson (1981).

Reading the PREM profile



- V_s (red) vanishes between 2890 and 5150 km depth: the outer core transmits no shear waves, so it is liquid.
- Density (green) jumps at the core-mantle boundary and again at the inner-core boundary: the metal-silicate and the solid-liquid interfaces.
- The liquid-shell geometry read off this curve is exactly the arena the dynamo needs.

Seismology hands the dynamo its arena: a 2260 km thick shell of convecting liquid iron.

Secular variation of the field

Earth's field is not static. It varies on many timescales:

- **Secular variation** (decades to centuries): non-dipole components grow, decay, drift
- **Westward drift**: non-dipole features migrate westward at $\sim 0.2^\circ/\text{yr}$, implying differential rotation between core and mantle
- **South Atlantic Anomaly (SAA)**: a $\sim 30\%$ reduction in field strength over South America and the South Atlantic
- SAA is the surface expression of a reversed-flux patch at the core–mantle boundary
- Expanding and weakening over the past century; potential precursor to a reversal (but unconfirmed)

Source: Finlay et al. (2020), Swarm satellite constellation data

Magnetic field reversals

- N and S magnetic poles **swap**
- Current rate: ~4–5 per Myr
- Each reversal takes
~1000–10,000 yr
- Field weakens, becomes multipolar,
re-establishes
- **Magnetic stripes** on ocean floor:
key evidence for seafloor spreading

Credit: Course figure; chron ages after Gradstein et al. (2020)

The Cretaceous Normal Superchron

Not all times have frequent reversals.

- **Cretaceous Normal Superchron (CNS): ~121–83 Ma**
- The field kept a **single polarity for ~38 Myr**
- Also a Permo-Carboniferous reversed superchron (~312–262 Ma)
- Superchrons may reflect changes in core–mantle boundary heat flow, controlled by slow mantle convection patterns


Reversal rate is not constant; it is modulated by the thermal evolution of the core–mantle boundary over hundreds of millions of years.

Source: Biggin et al. (2012); Olson & Amit (2015)

Magnetic fields across the solar system


Body	Field type	Surface field	Rel. dipole
Earth	Active dynamo	25–65 μT	1
Mercury	Active (weak)	$\sim 0.3 \mu\text{T}$	5×10^{-4}
Venus	None detected	$< 0.01 \mu\text{T}$	$< 10^{-5}$
Mars	Remnant crustal	up to 1500 nT	n/a
Jupiter	Active (strong)	400–1400 μT	20,000
Saturn	Active	$\sim 20 \mu\text{T}$	~ 600
Ganymede	Active	$\sim 0.7 \mu\text{T}$	1.5×10^{-3}

Jupiter's field is $\sim 20,000\times$ Earth's, generated in metallic hydrogen.




Surface radial magnetic fields of six solar-system dynamos. Soderlund & Stanley (2020).

Reading the field morphologies




- Earth, Jupiter, and Saturn: dipoles roughly aligned with the rotation axis; Saturn almost perfectly axisymmetric.
- Uranus and Neptune: multipolar, tilted $\sim 59^\circ$ and $\sim 47^\circ$, offset from the planet centre.
- Mercury: weak and offset ~ 500 km northward; the per-panel colour scales span four orders of magnitude.

One mechanism, many faces: geometry and shell structure sculpt what the dynamo shows at the surface.



Dipolar versus multipolar dynamo regimes. Olson & Christensen (2006).

Reading the regime diagram



- Axes: dimensionless dipole moment versus local Rossby number (a measure of inertia relative to rotation); dark region dipolar, light multipolar, calibrated on numerical dynamo runs.
- Earth, Jupiter, Saturn, and Ganymede sit in the dipolar regime; Mercury, Uranus, and Neptune lie in or near the multipolar one.
- Two control parameters predict the field morphology without simulating each planet individually.

Whether a planet wears a clean dipole is set by its Rossby number: rapidly rotating, thick-shell dynamos stay dipolar.

The Venus magnetic field puzzle

Venus: Earth's twin in size and composition, yet **no magnetic field**.

- Slow rotation (243-day period)? Simulations say this alone is insufficient.
- **No inner core?** Without compositional convection, thermal convection alone may be too weak.
- **Stagnant lid tectonics?** Mantle may not extract core heat efficiently enough.

Most likely: **combination** of reduced core cooling (no plate tectonics) and possibly no inner core nucleation.


Mercury: an offset dynamo

Mercury's magnetic field is weak but intriguing.

- Surface field: $\sim 0.3 \mu\text{T}$, 1% of Earth's
- Field is **offset** **$\sim 500 \text{ km north}$** of the geometric centre
- Dominated by the dipole component (g_{10}), weak higher harmonics
- MESSENGER (2011–2015) mapped this asymmetry
- Interpretation: thin liquid outer shell above a partially solidified deeper core
- Limits convective vigour and field strength

Source: Anderson et al. (2012), *Journal of Geophysical Research*

Mars: a fossil magnetic field



Credit: NASA/GSFC

- No global field today
- Intense **crustal magnetisation** in southern highlands
- Absent in young northern lowlands and large basins

This tells us:


1. Mars **had** an active dynamo
2. Dynamo died between ~4.1 and 3.7 Ga
3. Atmosphere left unshielded → erosion by solar wind

Mars dynamo: a more recent shutdown?

The classical Acuña (1999) interpretation placed the dynamo death at ~4.1 Ga. Recent work has revised this:


- Mittelholz et al. (2020): magnetisation in the ~3.7 Ga Lucus Planum lavas
- Implies Mars dynamo was **intermittent** or shut down **later** than thought
- Consistent with core cooling becoming insufficient over 4.1–3.7 Ga
- MAVEN: measured present-day atmospheric loss rate of ~100 g/s (O^+)
- Extrapolated over 4 Gyr: consistent with Mars losing ≥ 0.8 bar of CO_2

Source: Mittelholz et al. (2020), *Science Advances*; Jakosky et al. (2018)



InSight's Mars core: radius and density posteriors. Stähler et al. (2021).

Reading the Mars core inversion



- Independent inversions converge on $R_{\text{core}} \approx 1830$ km; the mean core density comes out near 6 g cm^{-3} .
- The purple bands (lines of fixed composition) demand several wt% of light elements: the core is too light for pure iron.
- A sulfur-rich core may never nucleate an inner core: no compositional buoyancy, an early-dying dynamo.

InSight's big, light core explains the fossil field: without inner-core growth there is no compositional buoyancy to keep the dynamo alive.

Jupiter: dynamo in metallic hydrogen

Jupiter has the strongest magnetic field in the solar system, $\sim 20,000\times$ Earth's dipole moment.

- Molecular H_2 becomes **metallic** above ~ 100 GPa
- Transition at ~ 0.85 Jupiter radii from the centre
- Metallic H has conductivity comparable to liquid metals
- Dynamo operates in the deep metallic-H layer
- Juno gravity measurements show the transition is **gradual**, not sharp

Juno has revealed a complex, asymmetric surface field with a “Great Blue Spot” (reversed-flux patch in the northern hemisphere).

Source: Connerney et al. (2022); Bolton et al. (2017)

Ganymede: the only moon with a dynamo

Ganymede is the **only moon** in the solar system with an intrinsic magnetic field.

- Moon discovered by Galileo Galilei (1610); intrinsic field detected by NASA *Galileo* spacecraft (1996): surface field $\sim 0.7 \mu\text{T}$
- Dipole tilted $\sim 176^\circ$ (nearly anti-parallel to Jupiter's field)
- Surprising: Ganymede is small ($R = 2634 \text{ km}$), should be thermally dead
- Possible explanations:
 - Tidal heating from Laplace resonance sustains core convection
 - Slow freezing of the outer core releases compositional buoyancy
- JUICE (arriving 2031) will map Ganymede's field in detail

Source: Kivelson et al. (1996); Bland et al. (2008)

Magnetopause standoff distance


Pressure balance at the magnetopause:

$$\underbrace{\frac{1}{2} \rho_{\text{sw}} v_{\text{sw}}^2}_{\text{solar wind ram pressure}} = \underbrace{\frac{B_{\text{mp}}^2}{2\mu_0}}_{\text{magnetic pressure}}$$

For a dipole field, $B \propto r^{-3}$, so the standoff distance scales as:


$$r_{\text{mp}} \propto \left(\frac{B_0^2}{\mu_0 \rho_{\text{sw}} v_{\text{sw}}^2} \right)^{1/6}$$

- Earth under typical solar wind: $r_{\text{mp}} \approx 10 R_{\oplus}$
- During a CME: compressed to $\sim 6 R_{\oplus}$
- Jupiter (much stronger field): $\sim 50\text{--}100 R_J$



Anatomy of Earth's magnetosphere: bow shock, magnetosheath, and magnetopause. Credit: ESA, CC BY-SA 3.0 IGO.

Reading the magnetosphere anatomy



- The solar wind arrives from the left; the bow shock slows and deflects it, and the shocked plasma fills the magnetosheath.
- The magnetopause sits where ram pressure equals magnetic pressure, at $r_{mp} \approx 10 R_{\oplus}$ under typical solar wind.
- The compressed dayside field and the stretched nightside tail both follow from that single balance.

One pressure equality fixes the size of the whole dayside magnetosphere.

Magnetotail reconnection

- On the night side, Earth's field is stretched into a long tail ($>200 R_{\oplus}$)
- Opposing field lines in the magnetotail can **reconnect**, releasing stored magnetic energy
- Reconnection accelerates particles toward both Earth and downstream
- Earthward-moving particles precipitate into the polar atmosphere → **aurorae**
- Downstream particles are ejected as **plasmoids** (self-contained blobs of magnetised plasma)
- Reconnection drives **magnetospheric substorms** observed every few hours during active solar conditions

Magnetic reconnection is the engine of space weather, converting solar wind kinetic energy into particle acceleration and auroral emission.

The Van Allen radiation belts


Earth's magnetosphere traps energetic particles in two toroidal belts:

Inner belt $\sim 1.5 R_{\oplus}$: high-energy protons (10–100 MeV), produced by cosmic-ray interaction with the atmosphere (CRAND process)

Outer belt $\sim 4\text{--}5 R_{\oplus}$: high-energy electrons (0.1–10 MeV), injected during geomagnetic storms and accelerated by wave–particle interactions

- Discovered by Explorer 1 (1958)
- Major hazard to spacecraft (electronics) and crewed missions
- Van Allen Probes (2012–2019): discovered a third, transient belt

Source: de Pater & Lissauer (2010); Baker et al. (2013)



Cross-section of Earth's Van Allen radiation belts. Credit: Booyabazooka / NASA, public domain.

Aurorae: light from the magnetosphere



Credit: NASA/ISS

Solar wind particles channelled along field lines to polar regions:

- **Green:** excited O at 100–200 km (557.7 nm)
- **Red:** excited O at 200–400 km (630.0 nm)
- **Blue/violet:** excited N_2 bands

Observed on Jupiter, Saturn, and Uranus: any magnetised planet.

Why aurorae happen at the poles


- Earth's dipole focuses incoming particles toward the magnetic poles
- **Cusps** near the poles are regions where solar wind has direct access to the atmosphere
- Reconnection in the magnetotail releases stored energy
- Accelerated electrons precipitate down field lines into the atmosphere
- Energies of 1–100 keV; penetrate to 100–400 km altitude
- Emission lines and bands are fingerprints of the excited species

Aurorae map the coupling between the solar wind and the magnetosphere, lit up by atmospheric atoms.

Jupiter's aurorae: the most powerful in the solar system

- Total UV power: 10^{13} – 10^{14} W (10^3 – $10^4\times$ Earth's auroral power)
- Powered by:
 - Jupiter's rapid rotation (~10 hr)
 - Mass loading from Io's volcanoes (~1 tonne/s of sulfur and oxygen)
 - Solar wind interaction (secondary)
- Io, Europa, and Ganymede each create **auroral footprints** by electromagnetic coupling
- Observed in UV by HST, Juno-UVS; infrared from ground-based telescopes

Source: Connerney et al. (2022); Mauk et al. (2017) on Juno plasma observations



Jupiter's ultraviolet aurora over an optical image, Hubble Space Telescope. Credit: NASA, ESA, J. Nichols (University of Leicester).


A photograph of the Aurora australis (Southern Lights) in a dark night sky. The aurora appears as a bright green, jagged, and elongated light structure. It has a main vertical column on the right side and a horizontal, diffuse band extending towards the left. The background is a deep black, with some faint, wispy clouds visible on the left side.

5. Recent advances & outlook

Juno: Jupiter's asymmetric dynamo


- Juno (2016–) has measured Jupiter's gravity and magnetic field in unprecedented detail
- Surface field dramatically **non-dipolar**
- Northern hemisphere: intense concentrated “flux lobe”
- “Great Blue Spot”: a patch of reversed polarity near the equator
- Magnetic equator offset from the rotational equator
- Implies dynamo operates at relatively shallow depths in the stably-stratified dilute core (Lecture 8)

Source: Connerney et al. (2022), *Journal of Geophysical Research*



Jupiter's surface radial field after Juno's prime mission (top) and before Juno (bottom). Credit: NASA/JPL-Caltech/SwRI (PIA25040).

Reading Jupiter's magnetic map



- Yellow and red mark outward flux, blue inward: the surface field is strongly non-dipolar.
- The Great Blue Spot: an intense localised patch of inward flux near the equator, invisible in the low-resolution pre-Juno map (bottom).
- The morphology places the dynamo in a deep but not razor-thin shell of metallic hydrogen.

Juno turned Jupiter's field from a textbook dipole into a weather map of the deep interior.

BepiColombo: new eyes on Mercury

- ESA/JAXA joint mission, launched 2018
- Multiple Mercury flybys (2021–2025) en route to orbit insertion
- Already confirmed the weak but active dipolar field
- Revealed sodium ion cyclotron waves in Mercury's magnetosphere
- Will map the offset field with unprecedented precision from polar orbit
- Two orbiters: Mercury Planetary Orbiter (MPO) + Mercury Magnetospheric Orbiter (MMO)

Source: Benkhoff et al. (2021), *Space Science Reviews*

JUICE: mapping Ganymede's dynamo

- Jupiter Icy Moons Explorer, launched April 2023
- Arrives at Jupiter in 2031; Ganymede orbit insertion 2034
- First orbiter of a moon other than Earth's
- Will map Ganymede's intrinsic field at high resolution
- Separate intrinsic field from induced field (ocean signature)
- Tests the hypothesis that Ganymede has a **subsurface ocean**
- Bridges Lecture 4 (dynamo physics) and Lecture 8 (interiors)

Source: Grasset et al. (2013) on JUICE science goals

MAVEN and the escape of Mars' atmosphere

- NASA MAVEN (2014–) measures Mars atmospheric loss in situ
- Total escape rate: ~2–3 kg/s, dominated by H and O
- Dominated by photochemical escape of oxygen
- Escape accelerates during solar storms
- Extrapolated over 4 Gyr: ~0.8 bar of CO₂ lost, consistent with a warmer early Mars
- **Directly ties** the loss of the dynamo (Lecture 4) to the loss of the atmosphere (Lecture 5)

Source: Jakosky et al. (2018), *Icarus*; Lillis et al. (2017)

Psyche: visiting a naked core

NASA Psyche launched in October 2023, arrives at asteroid 16 Psyche in 2029.

- Target: a ~220 km asteroid, one of the largest M-type
- Long suspected to be the exposed metallic core of a differentiated planetesimal
- Bulk density $\sim 4 \text{ g/cm}^3$ (higher than stony, lower than pure iron)
- Instruments: multispectral imager, magnetometer, gamma-ray spectrometer
- Will test whether Psyche has remnant magnetisation (ancient dynamo relic)
- A direct sample of the kind of body L4's theory describes

Source: Elkins-Tanton et al. (2017), *J. Geophys. Res. Planets*

Open questions in planetary magnetism

Despite decades of work, key questions remain:

- **When did the inner core nucleate?** (Most estimates now 0.5–1 Gyr ago)
- **How did Earth sustain a dynamo before inner core formation?**
- **Why does Venus have no detectable field?** Inner core, rotation, or tectonics?
- **Why is Mercury's field so weak yet active?** What powers it?
- **What drives Ganymede's dynamo?** (A moon should be thermally dead)
- **How do Uranus and Neptune generate multipolar fields?** (Exotic dynamo physics)
- **Will Earth reverse soon?** (No reliable prediction method exists)
- **How important are magnetic fields for habitability?** (Crucial or secondary?)

Future missions (JUICE, BepiColombo, Europa Clipper, MSR) will address several of these.

Looking ahead to Lecture 5

This lecture ended with a planet that has a core, a mantle, a magnetic field, and a freshly outgassed secondary atmosphere.

- Lecture 5 (dr. Mara Attia) picks up that atmosphere: primary, secondary, and tertiary atmospheres
- Pressure and temperature with height: hydrostatic equilibrium
- The greenhouse effect and planetary energy balance
- How atmospheres are lost to space; magnetic shielding is one control on escape, but thermal escape proceeds regardless

Differentiation builds the planet; the atmosphere it outgasses is where climate and habitability are decided.

Summary

1. Magma oceans drive **metal–silicate separation**: iron sinks, silicates float
2. Hf–W chronometry gives a two-stage age of ~30 Myr for Earth's core formation, a lower limit; protracted accretion and partial equilibration place completion within ~100 Myr
3. Magma ocean crystallisation creates chemically distinct layers (crust, mantle reservoirs)
4. Dynamos require: conducting fluid + convection + sufficient R_m (≥ 10 –100)
5. Earth's $R_m \approx 1750$: vigorous dynamo action
6. Mars lost its dynamo between ~4.1 and 3.7 Ga (possibly intermittent, Mittelholz 2020) → atmospheric erosion
7. Venus has no detectable field: likely a combination of stagnant-lid tectonics, slow rotation, and/or absence of an inner core

Questions?

Next: Lecture 5, Atmospheres I: composition, structure & energy balance (Mara Attia)

Mon 21 Sep, 15:00–17:00, room 5419.0161 (Collegezaal)

Next tutorial: Fri 25 Sep, 13:00–15:00, room 5419.0161 (Collegezaal)

<https://ips.formingworlds.space>