

Lecture 5: Atmospheres I

Composition, Structure, & Energy Balance

Planetary Systems

A horizontal, slightly curved blue line representing the Earth's atmosphere as seen from space. The line is thin and has a soft, glowing appearance, especially in the center where it seems to be illuminated from below.

Tim Lichtenberg
Kapteyn Astronomical Institute
University of Groningen
2026

Learning Objectives

By the end of this lecture, you will be able to:

- Classify atmospheric types (primary, secondary, tertiary)
- Derive pressure profiles from hydrostatic equilibrium
- Compute the atmospheric scale height and apply it across the solar system
- Explain how the greenhouse effect raises surface temperature above T_{eff}
- Evaluate atmospheric escape mechanisms and retention criteria

Recap: Lecture 4

- Metal–silicate separation driven by magma oceans
- Hf–W chronometry: Earth's core formed in ~30–60 Myr
- Goldschmidt classification: siderophile, lithophile, chalcophile, atmophile
- Dynamo requirements: conducting fluid + convection + $R_m \gtrsim 10\text{--}100$
- Mars lost its dynamo ~4.1 Ga → atmospheric erosion

Today: The thin gaseous envelope that controls a planet's surface conditions.



1. Atmospheric Composition

Captured directly from the protoplanetary disk.

- Dominated by H_2 and He (solar composition)
- Requires mass $\gtrsim 5\text{--}10 M_{\oplus}$ to capture nebular gas
- Disk disperses in $\sim 3\text{--}10$ Myr (Lecture 2)
- **Gas giants:** Jupiter and Saturn, massive H_2/He envelopes
- **Ice giants:** Uranus and Neptune, envelopes only 10–20% of total mass
- Terrestrial planets too small $\rightarrow$ any primordial H quickly lost

Secondary Atmospheres

Produced by outgassing from the interior.

- Volcanism and magma ocean degassing release dissolved volatiles
- Speciation depends on **oxygen fugacity** (Lecture 4):
 - Oxidising: CO_2 , H_2O , N_2
 - Reducing: H_2 , CO , N_2
- **Venus:** CO_2 -dominated (96.5%), 92 bar
- **Mars:** CO_2 -dominated (95%), but only 6 mbar
- **Titan:** thick N_2 (from NH_3 conversion), 1.5 bar

Where Do Atmospheres Come From?

A pictorial summary of the three classes.

Primary Directly captured nebular gas; H_2/He dominated.
Requires $M \gtrsim 5\text{--}10 M_{\oplus}$ before disk dispersal.

Secondary Outgassed from the interior during accretion + cooling.
Composition set by f_{O_2} of the mantle.

Tertiary Modified by surface processes, photochemistry, and (for Earth) biology.

Over time, a planet's atmospheric inventory can be **replaced** or **re-equilibrated** entirely: the present composition may not reflect the initial inventory.

Tertiary Atmospheres

Modified by surface processes, photochemistry, or biology.

- Earth's original outgassed atmosphere: likely $\text{CO}_2 + \text{N}_2$ (similar to Venus)
- Oxygenic photosynthesis transformed it over billions of years
- Present atmosphere: N_2 (78%) + O_2 (21%)
- O_2 **is entirely biogenic**, and would disappear in ~Myr without life


Earth's atmosphere is a **tertiary atmosphere**: the product of billions of years of biological modification.

Comparative Atmospheric Properties

	Venus	Earth	Mars	Jupiter	Titan
P_{surf} (bar)	92	1.0	0.006	n/a	1.5
T_{surf} (K)	735	288	210	n/a	94
Dominant gas	CO ₂	N ₂	CO ₂	H ₂	N ₂
Secondary gas	N ₂	O ₂	N ₂	He	CH ₄
μ	43.4	29.0	43.3	2.2	28.6
Type	2°	3°	2°	1°	2°

Staggering diversity: surface pressures spanning five orders of magnitude, temperatures from 94 to 735 K.

Venus: A Secondary Atmosphere in Extremis



Credit: USSR/NASA NSSDC, public domain

- *Venera 13* lander, 1 March 1982
- Surface pressure: 92 bar
- Surface temperature: 735 K
- Orange sky: thick CO₂ atmosphere scatters light
- Lander survived 127 minutes

A vivid demonstration of how a massive secondary atmosphere transforms surface conditions.



2. Hydrostatic Equilibrium

Hydrostatic Equilibrium

Consider a thin slab of atmosphere at height z , thickness dz , area A :

- Pressure from below (upward): $P(z) \cdot A$
- Pressure from above (downward): $P(z + dz) \cdot A$
- Weight (downward): $\rho(z) g A dz$

Force balance:

$$P(z) A - P(z + dz) A = \rho g A dz$$

$$\frac{dP}{dz} = -\rho g$$

Pressure decreases with altitude: each layer supports the weight above it.

The Ideal Gas Law (Atmospheric Form)

To solve hydrostatic equilibrium, we need $\rho(P)$:

$$P = n k_B T = \frac{\rho k_B T}{\mu m_u}$$

n number density

k_B Boltzmann constant ($1.381 \times 10^{-23} \text{ J K}^{-1}$)

μ mean molecular weight (dimensionless, in amu)

m_u atomic mass unit ($1.661 \times 10^{-27} \text{ kg}$)

Rearranging: $\rho = P \mu m_u / (k_B T)$

The Barometric Formula

Substitute $\rho = P\mu m_u / (k_B T)$ into hydrostatic equilibrium:

$$\frac{dP}{dz} = -\frac{\mu m_u g}{k_B T} P$$

For an **isothermal** atmosphere (constant T , constant g), this separates:

$$P(z) = P_0 \exp\left(-\frac{z}{H}\right)$$

where $H = k_B T / (\mu m_u g)$ is the **pressure scale height**.

- Pressure drops by $e \approx 2.718$ every scale height
- 99% of the atmospheric mass lies below $\sim 5H$

3. Blackboard Derivation: Scale Height

Deriving the Scale Height

Goal: Derive $H = k_B T / (\mu m_u g)$ from hydrostatic equilibrium + ideal gas law, and compute H for solar system bodies.

Start from:

$$\frac{dP}{dz} = -\rho g \quad \text{and} \quad P = \frac{\rho k_B T}{\mu m_u}$$

Express ρ in terms of P :

$$\rho = \frac{P \mu m_u}{k_B T}$$

Substitute:

$$\frac{dP}{dz} = -\frac{\mu m_u g}{k_B T} P \equiv -\frac{P}{H}$$

The Atmospheric Scale Height

$$H = \frac{k_B T}{\mu m_u g}$$

Physical interpretation:

- **Higher T** $\rightarrow$ larger H : hotter gas extends further
- **Heavier molecules** (larger μ) $\rightarrow$ smaller H : harder to loft
- **Stronger gravity g** $\rightarrow$ smaller H : more compressed

Key insight: H encodes the competition between thermal energy and gravitational binding.

Worked Example: Earth's Scale Height

$$H_{\oplus} = \frac{k_B T}{\mu m_u g} = \frac{1.381 \times 10^{-23} \times 288}{28.97 \times 1.661 \times 10^{-27} \times 9.81}$$
$$= \frac{3.98 \times 10^{-21}}{4.72 \times 10^{-25}} \approx 8400 \text{ m} \approx 8.4 \text{ km}$$

- Commercial aircraft: 10–12 km (~ 1.2 – $1.4 H$), pressure ~ 0.2 – 0.3 atm
- Top of Everest: 8.8 km ($\sim 1 H$), pressure ~ 0.3 atm
- 99% of mass below ~ 42 km ($\sim 5 H$)

Scale Heights Across the Solar System


Body	T (K)	μ	g (m s^{-2})	H (km)
Venus	735	43.4	8.87	15.8
Earth	288	29.0	9.81	8.5
Mars	215	43.3	3.72	11.1
Jupiter	165	2.2	24.8	27
Titan	94	28.6	1.35	21

- Jupiter: large H despite strong gravity, because H_2 is very light ($\mu = 2.2$)
- Titan: large H despite low T , thanks to very weak gravity ($g = 1.35 \text{ m s}^{-2}$)
- Venus: hot but heavy molecules + decent gravity $\rightarrow$ moderate H

4. Vertical Structure



Atmospheric Layers



Credit: Wikimedia, public domain

Layers defined by the sign of $\frac{dT}{dz}$:

1. **Troposphere** (0–12 km): T decreases with z
Heated from below; convective
2. **Stratosphere** (12–50 km): T increases
UV absorbed by ozone layer
3. **Mesosphere** (50–85 km): T decreases
Coldest point at mesopause (~190 K)
4. **Thermosphere** (>85 km): T increases
EUV absorption; $T > 1000$ K but gas is rarefied

The Adiabatic Lapse Rate

In the troposphere, convection sets the temperature gradient:

$$\Gamma_d = -\frac{dT}{dz} = \frac{g}{c_p}$$

- **Dry adiabatic lapse rate** for Earth: $\Gamma_d = 9.81/1004 \approx 9.8 \text{ K km}^{-1}$
- Observed average: $\sim 6.5 \text{ K km}^{-1}$ (lower due to latent heat from water condensation)
- Physical picture: rising air expands, does work on surroundings, cools

Tropopause height: $\sim 8 \text{ km}$ (poles) to $\sim 17 \text{ km}$ (equator).

Comparative Vertical Structures


Venus: Massive troposphere (~65 km). No ozone layer → no stratospheric inversion. Cloud deck at 48–70 km.


Mars: Thin troposphere (~40 km), directly into thermosphere. No significant ozone. Dust dominates atmospheric heating.

Jupiter: Troposphere extends hundreds of km deep. Stratosphere heated by CH_4 . No solid surface; pressure increases continuously.

Titan: Thick troposphere (~40 km). Stratosphere heated by organic haze. Extended thermosphere reaches ~1500 km (weak gravity).

Break





Three Interactions: Radiation and Matter

When radiation passes through an atmosphere:

Absorption: Photon energy $\rightarrow$ internal energy (vibrational, rotational)

Key absorbers: CO_2 , H_2O , CH_4 , O_3 , N_2O

Emission: Warm gas radiates at wavelengths where it absorbs (Kirchhoff's law)


This is how the atmosphere radiates energy to space

Scattering: Photons redirected, not absorbed

Rayleigh ($\propto \lambda^{-4}$): blue sky, red sunsets

Mie (larger particles): less wavelength-dependent

Stellar vs. Planetary Emission



Credit: Wikimedia, public domain

- Sun (~5800 K): peak at ~0.5 μm (visible)
- Planet (~300 K): peak at ~10 μm (thermal IR)

This **wavelength separation** is the physical basis of the greenhouse effect: the atmosphere can be transparent at visible wavelengths but opaque in the infrared.

Optical Depth

The cumulative absorption along a path through the atmosphere:

$$\tau = \int_0^s \kappa \rho \, ds'$$

- κ = mass absorption coefficient ($\text{m}^2 \text{kg}^{-1}$)
- τ is dimensionless: counts “e-folding lengths” of absorption

Optically thin ($\tau \ll 1$):

Most radiation passes through.

Optically thick ($\tau \gg 1$):

Radiation strongly absorbed; only photons from near the “surface” escape.

The Classical Habitable Zone

Where around a star can a planet maintain liquid water?

- Inner edge: runaway greenhouse limit (water fully evaporates)
- Outer edge: CO₂ condenses into ice clouds (Maximum Greenhouse limit)
- For the Sun: classical HZ is ~0.95–1.67 AU (Kopparapu et al. 2013)
- Assumes an Earth-like atmosphere with CO₂/H₂O feedback
- Does not account for tidal heating, magnetic fields, or greenhouse gas diversity
- A starting point, not the last word

Source: Kasting et al. (1993); Kopparapu et al. (2013)

The Moist Adiabatic Lapse Rate

When a parcel rises and condensation occurs, latent heat release warms it relative to the dry adiabat.

$$\Gamma_{\text{moist}} = \Gamma_{\text{dry}} \cdot \frac{1 + \frac{L_v r_s}{R_d T}}{1 + \frac{L_v^2 r_s}{c_p R_v T^2}}$$

- r_s : saturation mixing ratio; L_v : latent heat of vaporisation
- Earth's lower troposphere: $\Gamma_{\text{moist}} \approx 6 \text{ K/km}$ (compared to 9.8 K/km dry)
- Moist convection is **deeper and more vigorous** than dry; drives thunderstorms and hurricanes
- Latent heat release in rising parcels provides the buoyancy that makes tropical convection reach the tropopause
- Runaway greenhouse reduces Γ_{moist} further, letting more IR escape to

Deriving the Scale Height (Live)

Template for the blackboard derivation.

1. Start: $\frac{dP}{dz} = -\rho g$ (hydrostatic equilibrium)

2. Ideal gas: $P = \frac{\rho k_B T}{\mu m_u}$

3. Rearrange for ρ : $\rho = \frac{\mu m_u P}{k_B T}$

4. Substitute: $\frac{dP}{dz} = -\frac{\mu m_u g}{k_B T} P$

5. Separate variables and integrate:

$$P(z) = P_0 \exp\left(-\frac{z}{H}\right), \quad H = \frac{k_B T}{\mu m_u g}$$

6. Numerical check: Earth ($T = 288 \text{ K}$, $\mu = 29$, $g = 9.81$) gives $H \approx 8.4 \text{ km}$

Transmission Spectroscopy: Preview

During a transit, starlight passes through the planet's **limb**.

- Absorbing molecules imprint dark features in the transmission spectrum
- Path length $\sim 2\sqrt{2R_p H}$ through a layer of scale height H
- Signal: $\Delta(R_p/R_\star)^2 \sim 2R_p H/R_\star^2$
- Bigger signal for puffy atmospheres (large H) around small stars
- JWST has detected H_2O , CO_2 , CH_4 , SO_2 in multiple exoplanet atmospheres
- Full treatment in Lecture 13 (Exoplanet Atmospheres)

Source: Seager & Sasselov (2000); Kreidberg et al. (2014)


The Beer–Lambert Law

For a beam of initial intensity I_0 through a medium of optical depth τ :

$$I = I_0 e^{-\tau}$$

- Intensity drops by factor e per unit optical depth
- **Atmospheric photosphere:** at $\tau \approx 1$, photons escape to space
- Below $\tau = 1$: photons reabsorbed before escaping
- Above $\tau = 1$: photons escape freely
- Temperature at the $\tau \approx 1$ level sets T_{eff}

Earth's Atmospheric Absorption Spectrum



6. Energy Balance & Greenhouse Effect



Radiative-Convective Equilibrium

A realistic atmosphere combines two regimes:

- **Troposphere:** strong greenhouse absorption → unstable to convection
 - Temperature follows the adiabatic lapse rate
 - Most of the atmospheric mass
 - Clouds and weather occur here
- **Stratosphere:** optically thin → radiative equilibrium
 - Temperature set by the balance of absorbed vs. emitted radiation
 - Stable against convection
 - On Earth, warmed by ozone UV absorption

The boundary (the tropopause) occurs where the radiative gradient becomes less steep than the adiabat.

Manabe and Strickler (1964)

The first realistic 1D radiative-convective model of Earth's atmosphere.

- Solved for the vertical temperature profile with absorbing gases (H_2O , CO_2 , O_3)
- Reproduced the surface temperature and the tropopause self-consistently
- Quantified the effect of doubling CO_2 : ~ 2 K warming (first “climate sensitivity” estimate)
- Foundation of modern climate modelling
- Manabe received the 2021 Nobel Prize in Physics for this work

Source: Manabe & Strickler (1964), *J. Atmos. Sci.*

Planetary Energy Balance

Absorbed stellar power = emitted thermal radiation:

$$(1 - A) \frac{L_{\star}}{4\pi d^2} \pi R_p^2 = 4\pi R_p^2 \sigma T_{\text{eff}}^4$$

- A = Bond albedo (fraction of light reflected)
- πR_p^2 = cross-sectional area (intercepts light)
- $4\pi R_p^2$ = total surface area (emits isotropically)

Solving:

$$T_{\text{eff}} = \left[\frac{(1 - A) L_{\star}}{16\pi\sigma d^2} \right]^{1/4}$$


Effective vs. Surface Temperature

Body	A	d (AU)	T_{eff} (K)	T_{surf} (K)	ΔT (K)
Venus	0.77	0.72	227	735	+508
Earth	0.30	1.00	255	288	+33
Mars	0.25	1.52	210	210	~0
Jupiter	0.34	5.20	110	165*	+55

*Jupiter 1-bar level; excess partly from internal heat (Lecture 3).

$\Delta T = T_{\text{surf}} - T_{\text{eff}}$ reveals the **greenhouse effect**.
Venus: 508 K. Earth: 33 K. Mars: ~0 K (too thin).

The Greenhouse Mechanism



Credit: Wikimedia, public domain

1. Sunlight (visible, $\sim 0.5 \mu\text{m}$) reaches the surface
2. Surface emits thermal IR ($\sim 10\text{--}15 \mu\text{m}$)
3. Greenhouse gases absorb outgoing IR
4. Absorbing layer re-emits: half up, half down
5. Downward emission adds energy to surface

Result: $T_{\text{surf}} > T_{\text{eff}}$

One-Layer Greenhouse Model: Setup

Single isothermal layer, emissivity ε in IR, transparent at visible:

Atmospheric balance:

$$\varepsilon \sigma T_s^4 = 2 \varepsilon \sigma T_a^4 \quad \Rightarrow \quad T_a^4 = \frac{T_s^4}{2}$$

Surface balance:

$$(1 - A) \frac{F_\star}{4} + \varepsilon \sigma T_a^4 = \sigma T_s^4$$

Top-of-atmosphere:

$$(1 - A) \frac{F_\star}{4} = \sigma T_{\text{eff}}^4$$

One-Layer Greenhouse Model: Result

Combining the three balance equations:


$$T_s = T_{\text{eff}} \left(\frac{2}{2 - \varepsilon} \right)^{1/4}$$

$$\varepsilon = 0 \text{ (no greenhouse): } T_s = T_{\text{eff}}$$

$$\varepsilon = 1 \text{ (perfect absorber): } T_s = 2^{1/4} T_{\text{eff}} \approx 1.19 T_{\text{eff}}$$

For Earth: $T_s \approx 1.19 \times 255 \approx 303$ K. Reasonable first estimate.

Next lecture: What happens when T_s rises so high that outgoing radiation saturates? → *Runaway greenhouse* (Lecture 6).



Thermal (Jeans) Escape

Molecules have a Maxwell–Boltzmann velocity distribution. The high-velocity tail extends beyond escape speed.

The **Jeans escape parameter** at the exobase:

$$\lambda_J = \frac{GM_p m}{k_B T_{\text{exo}} r_{\text{exo}}} = \frac{v_{\text{esc}}^2}{v_{\text{th}}^2}$$

- $\lambda_J \gg 1$: escape negligible (tail too depleted)
- $\lambda_J \lesssim 2\text{--}3$: rapid atmospheric erosion
- Escape flux: $\Phi_J \propto (1 + \lambda_J) e^{-\lambda_J}$, exponentially sensitive

Jeans Parameter for Earth and Mars

Exobase temperatures: 1000 K (Earth), 300 K (Mars)

Species	m (u)	λ_j (Earth)	λ_j (Mars)
H	1	7.5	5.4
H ₂	2	15	11
He	4	30	22
N ₂	28	210	150
CO ₂	44	330	240

Heavy species (N₂, CO₂): λ_j so large that Jeans escape is negligible.
Atomic H: moderate $\lambda_j \rightarrow$ both Earth and Mars lose hydrogen to space.

Hydrodynamic Escape

When stellar EUV heating is intense, escape transitions from molecule-by-molecule to a bulk **atmospheric wind**:

- Most important in the first ~100 Myr (young stars: 10–100× more EUV)
- Can strip H₂-rich primary atmospheres from planets up to several $M_{\oplus}$
- Outflowing H can **drag along heavier species** (He, C, N, O)
- Leading explanation for the exoplanet **radius valley** (deficit at ~1.5–2 $R_{\oplus}$; Lecture 13)

Stripping Primary Atmospheres

Young stars are highly active in EUV and X-ray; primordial H/He envelopes can evaporate.

- EUV flux is $\sim 100\times$ higher in the first 100 Myr than today
- Mass loss rate for hydrodynamic regime: $\dot{M} \propto F_{\text{EUV}} R_p^3 / (GM_p)$
- For sub-Neptunes ($M_p \lesssim 10 M_\oplus$), atmospheres can be fully stripped in a few hundred Myr
- For Neptune-mass planets and above, envelopes are retained
- Produces the “evaporation valley” at $\sim 1.8 R_\oplus$ in Kepler data

Source: Owen & Wu (2013); Lopez & Fortney (2013); Fulton et al. (2017)

Energy-Limited Escape Rate

A convenient estimate of mass loss:

$$\dot{M}_{\text{EL}} = \eta \frac{\pi F_{\text{EUV}} R_p^3}{G M_p}$$

- F_{EUV} : incident EUV flux
- η : heating efficiency, typically 0.1–0.2
- Scales as R_p^3 : large, fluffy atmospheres lose mass fastest
- Scales inversely with M_p : low gravity eases escape
- Valid when recombination is negligible (“energy-limited regime”)

Source: Watson et al. (1981); Owen (2019)

Non-Thermal Escape Mechanisms

Sputtering: Solar wind ions impact upper atmosphere, eject molecules.
Significant for Mars (no global magnetic field).


Photochemical: UV dissociates $\text{H}_2\text{O} \rightarrow \text{H} + \text{OH}$; fast H escapes.
Key for Venus and Mars water loss.

Ion pickup: Atmospheric atoms ionised by UV, swept away by solar wind B -field.
Effective at unmagnetised planets.

Impact erosion: Large impacts can eject substantial atmospheric mass in a single event.

MAVEN at Mars: present loss rate $\sim 100 \text{ g s}^{-1}$ (O^+), dominated by solar wind stripping.

Atmospheric Retention: v_{esc} vs. T Diagram




Rule of thumb for long-term retention:

$$v_{\text{esc}} \gtrsim 6 v_{\text{th}}$$

- Gas giants: retain **everything**
- Earth/Venus: retain heavy species, lose H
- Mars: marginal; non-thermal losses dominate
- Titan: cold $\rightarrow$ retains N_2 despite low v_{esc}

Mars: A Lost Atmosphere



Credit: NASA/JPL (Viking 1, 1976)

- Present surface pressure: only 6 mbar
- Too thin for liquid water or significant greenhouse warming
- Geological evidence: ancient rivers, lakes, possibly ocean
- Mars had a much thicker atmosphere >4 Gyr ago
- Dynamo loss (~4.1 Ga, Lecture 4) → unshielded solar wind erosion



8. Recent Advances & Outlook

Recent Advances

- **JWST / TRAPPIST-1 b:** Thermal emission consistent with bare rock, no significant atmosphere. Inner rocky planets around active M dwarfs may be stripped.
- **JWST / TRAPPIST-1 c:** Similar result, reinforcing theoretical predictions of enhanced atmospheric escape around low-mass stars.
- **MAVEN at Mars:** Quantified loss rates for multiple species. Ion escape (solar wind stripping) dominates over Jeans escape. Integrated losses can account for a substantial fraction of Mars's early atmosphere.

Central theme: atmospheric retention depends on mass, temperature, stellar radiation, magnetic shielding, and geological activity.

Impact Erosion of Atmospheres

Giant impacts can eject substantial atmospheric mass in a single event.

- Shockwaves from a >km-sized impactor accelerate upper-atmospheric gas above escape velocity
- Efficiency depends on impact angle, velocity, and atmosphere mass
- A grazing giant impact can eject the entire atmosphere above the impact horizon
- Key process during the first ~100 Myr of solar system history
- May explain why Earth's early atmosphere was lost despite adequate gravity
- Signature: noble gas isotope ratios ($^{36}\text{Ar}/^{38}\text{Ar}$) record impact-driven loss

Source: Schlichting & Mukhopadhyay (2018); Genda & Abe (2005)

Outgassing vs. Escape: A Dynamic Balance

An atmosphere is not a static inheritance; it is a reservoir in flow.

- **Supply:** volcanic outgassing, impact delivery, biological fluxes
- **Loss:** thermal escape, non-thermal escape, chemical weathering, sequestration in oceans/crust
- The **steady state** depends on the balance of these fluxes
- Earth's active volcanism continually replaces CO_2 on the carbonate-silicate timescale (~ 0.5 Myr)
- Mars lost its active volcanism and dynamo, so escape was not replenished
- Venus has active (perhaps episodic) volcanism, but no oceans to absorb CO_2

Atmospheric longevity is as much about **replacement** as about initial inventory.

MAVEN at Mars: Quantifying Escape

NASA's MAVEN (2014–) has measured Mars' atmospheric loss in situ.

- Present-day total escape rate: ~1–2 kg/s (all species)
- Dominant loss channel: photochemical escape of oxygen
- Enhanced during solar storms (~10× increase)
- Integrated loss over 4 Gyr: consistent with 0.5–0.8 bar of CO₂ lost
- Combined with Jeans escape of hydrogen: ~15–25 m global equivalent layer of water lost

Source: Jakosky et al. (2018), *Icarus*; Lillis et al. (2017)

JWST and the Fate of Rocky Atmospheres

JWST has begun to directly test whether rocky planets around M dwarfs retain atmospheres.

- TRAPPIST-1 b: MIRI thermal emission consistent with a bare rock (dayside 508 K)
- TRAPPIST-1 c: MIRI shows low heat redistribution $\Rightarrow$ thin or no atmosphere
- LHS 3844 b: similar result using Spitzer data
- Implications: inner rocky planets around active M dwarfs lose their atmospheres
- Open question: whether **outer** TRAPPIST-1 planets retain atmospheres
- JWST will target TRAPPIST-1 e, f during Cycle 3–4

Source: Greene et al. (2023); Zieba et al. (2023), *Nature*

Runaway Greenhouse: Preview for Lecture 6

What happens when a water-rich atmosphere gets too hot?

- More water vapour → stronger greenhouse → hotter surface
- Positive feedback: hotter surface → more evaporation → more greenhouse
- At $\sim 360 \text{ W/m}^2$ absorbed, the **Simpson-Nakajima limit** is reached
- Beyond this, oceans fully evaporate and the runaway becomes unstoppable
- Hydrogen then escapes hydrodynamically, leaving a dry CO_2 atmosphere
- Venus' modern state is the end point of this process (Nakajima 1992; Goldblatt 2012)

Full treatment in **Lecture 6: Atmospheres II**.

Summary

1. Three atmosphere types: **primary** (captured H_2/He), **secondary** (outgassed), **tertiary** (biologically modified)
2. Hydrostatic equilibrium: $\frac{dP}{dz} = -\rho g$; leads to exponential pressure decay
3. Scale height: $H = k_B T / (\mu m_u g)$; encodes thermal energy vs. gravity
4. Vertical structure: troposphere, stratosphere, mesosphere, thermosphere
5. Beer–Lambert law: $I = I_0 e^{-\tau}$; $\tau \approx 1$ defines the emission level
6. Energy balance: $T_{\text{eff}} \propto [(1 - A)L_\star / d^2]^{1/4}$; greenhouse raises T_s above T_{eff}
7. Escape: Jeans (λ_j), hydrodynamic, non-thermal; retention requires $v_{\text{esc}} \gtrsim 6 v_{\text{th}}$

Questions?

Next: **Lecture 6. Atmospheres II: Clouds, Weather, & Climate**

Lecture notes: formingworlds.github.io/IntroductionToPlanetaryScience