


A global map of Mercury's topography in a pseudocylindrical projection. The map uses a color scale from dark blue (-5 km) to dark red (3 km). The equatorial region is predominantly red and orange, indicating higher elevations, while the polar regions are mostly blue and green, indicating lower elevations. Numerous impact craters of various sizes are visible across the surface. A vertical color bar on the right side of the map provides the elevation scale in kilometers.

Lecture 10: Rocky Planets II

Mercury & Mars

Planetary Systems

Tim Lichtenberg

Kapteyn Astronomical Institute
University of Groningen

2026

Learning Objectives

By the end of this lecture, you will be able to:

- Explain Mercury's large iron core, its 3:2 spin-orbit resonance and its weak offset dynamo
- Describe Mars' interior from the InSight seismic results and its wet-to-dry surface record
- Derive the Jeans escape flux, apply it to H, H₂ and CO₂ on Mars, and distinguish thermal from non-thermal escape

Recap: Lecture 9



Credit: NASA/JPL

- Earth and Venus: similar bulk parameters, radically divergent evolution
- Simpson-Nakajima runaway limit: maximum OLR of a saturated water atmosphere $\sim 280 \text{ W m}^{-2}$
- Plate tectonics, dynamo, and biosphere form an Earth-only coincidence

Today: Mercury and Mars, the smaller siblings at opposite evolutionary limits.



1. Mercury Overview and Surface


Caloris Basin, Mercury. MESSENGER (NASA/JHUAPL)

Mercury at a Glance

Parameter	Value	Note
Mass	$0.0553 M_{\oplus}$	smallest planet
Radius (mean)	$0.383 R_{\oplus}$	2440 km
Mean density	5429 kg m^{-3}	uncompressed $\sim 5300 \text{ kg m}^{-3}$
Semimajor axis	0.387 AU	innermost
Orbital period	87.97 d	
Rotation period	58.65 d	3:2 spin-orbit resonance
Eccentricity	0.206	largest of any planet
T_{surf} (day/night)	700/100 K	no atmosphere to redistribute
Magnetic field	$\sim 1\%$ Earth's	weak but global

Source: NASA Mercury Fact Sheet (NSSDC)

Mercury's interior structure (post-MESSENGER model)




Mercury's internal structure from MESSENGER: crust, thin silicate mantle, liquid outer core and solid inner core. Credit: Course-original figure.

A crust of 30–50 km and a 400 km mantle overlie a metallic core that fills 83% of the radius.


The 3:2 Spin-Orbit Resonance

Mercury rotates exactly three times for every two orbits.

- Discovered by radar (Pettengill & Dyce 1965); stabilised by the high eccentricity ($e = 0.206$)
- The solar day lasts 176 Earth days, exactly 2 Mercury orbits
- The solar tidal torque on a permanent quadrupole moment locks the resonance




MESSENGER MLA global topography: about 10 km of relief across ancient cratered highlands and younger volcanic plains (3.7–3.9 Ga). Credit: NASA/JHUAPL/CIW.




Caloris Basin, about 1550 km across, formed about 3.8 Ga, with its concentric rings and lava-flooded floor. Credit: NASA/JHUAPL (MESSENGER).

The largest well-preserved basin on Mercury and its chronostratigraphic reference; the focused seismic shock built the antipodal “weird terrain”.




A lobate scarp: a compressional thrust fault with up to 3 km of relief and over 1000 km of length. Credit: NASA/JHUAPL (MESSENGER); Byrne et al. (2014).

Secular cooling shrank Mercury by $\Delta R \approx 7$ km since the late bombardment; the area loss $\Delta A/A = 2\Delta R/R \approx 0.6\%$ is taken up by thrust scarps across one unbroken lid.



North polar topography and reflectance from MESSENGER MLA with modelled illumination and temperature: the permanently shadowed craters coincide. Credit: Neumann et al. (2013).

Crater floors below 100 K are cold traps; neutron spectrometry confirms water ice, delivered by comet and asteroid impacts, on the hottest planet.




Surface composition of Mercury from the MESSENGER X-ray and gamma-ray spectrometers.
Credit: Nittler et al. (2020).

High S, K, Cl and Na despite 700 K days, FeO below 2% and a high Mg/Si ratio: a highly reducing mantle formed under low oxygen fugacity.

Exosphere and Magnetosphere


A surface-bounded exosphere fed by sputtering, photon desorption and micrometeoroid vaporisation.

- Surface pressure $\sim 10^{-12}$ bar; detected species include Na, K, Ca, Mg, He, H
- The sodium tail extends $\sim 10^6$ km anti-sunward under radiation pressure
- A miniature magnetosphere stands off the solar wind at $\sim 1.5 R_{Hg}$ subsolar distance

A grayscale image of the Mercury surface, showing a dense field of craters of various sizes. A prominent, dark, linear feature, possibly a fault or a deep crater, runs diagonally from the upper right towards the lower center. The surface texture is highly irregular due to the numerous impact craters.


2. Mercury Interior and Dynamo

Mercury surface detail. MESSENGER (NASA/JHUAPL)





Mercury's layers from the libration and gravity measurements. Credit: Margot et al. (2018).

A crust of ~ 35 km and a ~ 400 km silicate mantle above a liquid outer core below ~ 2020 km radius (~ 83% of the planet) possibly capped by a thin solid Fe-S layer, and an unresolved solid inner core: core mass fraction ~ 74%.




Radar speckle interferometry measures Mercury's 88-day forced libration. Credit: Margot et al. (2007), Fig. 3.

The amplitude of 35.8 ± 2 arcseconds gives $C_m/C \approx 0.5$: only the silicate shell librates, so the mantle floats on a liquid core.




Monte Carlo distributions of the mantle's share of the moment of inertia, C_m/C , from the libration amplitude and C_{22} , for C/MR^2 of 0.325 (red) and 0.380 (blue). Reproduced from Margot et al. (2007).

Every distribution peaks near $C_m/C \approx 0.5$, far from the value of 1 a fully solid Mercury requires; MESSENGER later gave $C/MR^2 = 0.346$, between the two cases.






Interiors of the terrestrial planets to scale: Mercury's core is out of proportion. Credit: NASA Solar System Exploration.

Giant impact mantle stripping leads; **thermal vaporisation** is disfavoured by the retained volatiles (Na, S, K) and **selective condensation** of metal before silicate is no longer favoured.



Remnant mass against semimajor axis in N-body terrestrial-formation simulations for surface-density slopes $x = 0.5$ – 5.5 ; triangles mark the real planets. Credit: Franco et al. (2022), Fig. A1.

A Mercury-mass, iron-rich remnant at 0.39 AU appears in $\ll 1\%$ of trials: a single-impact origin is dynamically rare.



Mercury's dipole, offset $\sim 0.2 R_{Hg}$ northward, in a dynamo model with a stably stratified layer. Credit: Wicht & Heyner (2017).

Convection in a thin outer-core shell gives a surface field of only $\sim 200\text{--}400$ nT (Earth $\sim 50,000$ nT), and the stratified liquid layer above suppresses the high-degree harmonics.

A grainy, black and white photograph of the Martian surface, showing numerous craters of various sizes. The image is framed by a dark border with white tick marks, resembling a film strip. The text "3. Mars Overview and Moons" is overlaid in white.


3. Mars Overview and Moons

Mariner 4 view of Mars (1965). NASA/JPL

Mars at a Glance

Parameter	Value	Note
Mass	$0.107 M_{\oplus}$	one tenth of Earth
Radius (mean)	$0.532 R_{\oplus}$	3389 km
Mean density	3934 kg m^{-3}	lowest of the rocky planets
Semimajor axis	1.524 AU	
Orbital period	686.97 d	
Rotation period	24.62 h	similar to Earth
Axial tilt	25.2°	seasons like Earth's
Eccentricity	0.0934	strong climate forcing
T_{surf} (day/seasonal)	-130 to $+30^{\circ}\text{C}$	
Surface pressure	$\sim 6 \text{ mbar}$	0.6% of Earth's
Magnetic field	none global	local crustal anomalies

Source: NASA Mars Fact Sheet (NSSDC)




The first close-up of Mars, Mariner 4 on 15 July 1965: cratered terrain ended the speculation about canals and vegetation. Credit: NASA/JPL.

Phobos and Deimos


Two small rubble-pile moons: captured carbonaceous asteroids, or re-accreted from a giant-impact debris disk.

- **Phobos:** diameter 22 km, orbital period 7.7 h at 9376 km radius
- **Deimos:** diameter 12 km, orbital period 30.3 h at 23463 km radius
- Phobos orbits below the synchronous radius and will tidally disrupt in 30–50 Myr



SPH simulation of a Borealis-scale impact on Mars that forms an extended debris disk. Credit: Hyodo et al. (2017).

Phobos and Deimos accrete from the disk, which reproduces their orbits and predicts a fraction of Mars-derived material in both moons; the composition (Mars mantle or D-type) is still debated.




Operation plan of JAXA's Martian Moons eXploration: a three-year stay at Mars in five phases, drawn for the earlier 2024 launch baseline. Credit: Kuramoto et al. (2022), Fig. 4.

Launch now 2026, touchdowns on Phobos and a Deimos flyby between the Mars eclipses, sample return 2031: the samples will decide between capture and impact origin.

The image displays four terrestrial planets in a row, scaled to show their relative sizes. From left to right: Mercury is a small, grey, cratered sphere; Venus is a large, pale yellowish-brown sphere with faint cloud bands; Earth is a large, blue and white sphere showing continents and swirling cloud patterns; and Mars is a medium-sized, reddish-brown sphere with darker patches. The text "4. Mars' Interior: The InSight Revolution" is centered over the planets in white.



4. Mars' Interior: The InSight Revolution

Terrestrial planets to scale. Credit: NASA/JPL-Caltech




Mars' interior from InSight SEIS: crust, mantle and core. Credit: Stähler / Khan / Knapmeyer-Endrun (2021).

Crust 24–72 km above a mantle to ~ 1560 km depth; the shear reflector at ~ 1830 km radius was read as the liquid core in 2021. The 2023 reanalyses place a molten silicate layer above a smaller core (~ 1650–1675 km, ~ 6500 kg m^{-3}).




Marsquake waveforms recorded by InSight on the transverse component: the direct S arrival and the core-reflected ScS window. Credit: Stähler et al. (2021).

ScS shear reflections confirm a liquid core; far-side quakes and impacts (2023) refine its radius to ~ 1650–1675 km; over 1300 events were recorded, the largest S1222a at $M = 4.7$.



A 3D thermal-evolution model of Mars' mantle consistent with the InSight constraints. Credit: Pleša et al. (2022).

A few long-wavelength upwellings and sluggish lower-mantle flow under a thick stagnant lid; the Tharsis plume persists to the present.




Present-day crustal thickness of Mars: the global mean spans 40.6 km (31 km at InSight) in the thin end-member to 71.4 km (49 km at InSight) in the thick one. Pleša et al. (2022).




5. Mars Geology and Surface Highlights

Olympus Mons, Viking Orbiter mosaic. Credit: NASA/JPL/USGS




The three classical periods of Mars' geological time scale. Credit: Tanaka et al. (2014).

Noachian (> 3.7 Ga): heavy bombardment, valley networks, warm and wet episodes; **Hesperian** (3.7–3.0 Ga): outflow channels and large-scale volcanism; **Amazonian** (< 3.0 Ga): cold, hyper-arid, thin atmosphere.




The global geological map of Mars. Credit: Tanaka et al. (2014).


Tharsis dominates the western hemisphere, Hellas, Argyre and Isidis are the giant impact basins, and the northern lowlands are young, smooth and partly buried.



Olympus Mons: a shield volcano ~ 21 km high and 600 km wide, built by stationary hotspot volcanism over a stagnant lid. Viking Orbiter mosaic, credit: NASA/JPL/USGS, public domain.




Valles Marineris: a graben system 4000 km long and up to 7 km deep, opened by crustal extension under the Tharsis load. Credit: NASA/JPL-Caltech.




Dendritic valley networks carved into Noachian terrain, imaged by the Viking Orbiters. Credit: NASA/JPL/USGS.

Confluences, meanders and tributaries confirm liquid-water erosion, with drainage densities near terrestrial arid networks, active for > 100 Myr in the late Noachian.




Late Noachian valley networks (~ 3.6 Ga and older, top) against late Hesperian to Amazonian fans (3.5–3 Ga, bottom); elevations –6 to +6 km. Credit: Kite et al. (2022), Fig. 1.

The early valleys sit on high ground and the later fans in a mid-latitude band: fluvial control shifts from **elevation** to **latitude** as the climate cools.




The OMEGA mineralogy time line: Phyllosian, Theiikian, Siderikian. Credit: Bibring et al. (2006), Fig. 5.

Phyllosian: phyllosilicates from neutral-pH water; **Theiikian:** sulfates from acidic, evaporating water; **Siderikian** (from ~ 3.5 Ga): anhydrous ferric oxides under hyper-arid conditions.




Global distribution of phyllosilicates (red), sulfates (blue) and other hydrated minerals (yellow). Credit: Bibring et al. (2006), Fig. 3.

Clays cluster in the ancient southern Noachian highlands, sulfates in the equatorial evaporative basins.



Widespread unweathered olivine in the Martian crust. Credit: Ehlmann & Edwards (2014).


Olivine alters quickly in liquid water to serpentine and smectite clays, so its survival rules out a persistent global ocean: water activity was episodic or localised.




Mars surface compositions from infrared remote sensing and rover instruments: basalts, hydrated silicates, sulfates, carbonates, hematite and opaline silica record a wide range of aqueous and igneous environments. Ehlmann & Edwards (2014).

Mars Sample Return Campaign Concept (Schematic)

On Mars: collection and ascent



Return to Earth: capture, entry, analysis




Note: Campaign architecture and schedule are being rebaselined after the 2024 cost review; concept shown is schematic.

Mars Sample Return concept: cache at Jezero, retrieval lander and ascent vehicle, orbital capture, Earth entry and laboratory analysis; under rebaselining after the 2024 cost review.

Course figure.


Break



6. Blackboard Derivation: Jeans Escape Flux



Goal and strategy



Goal: the rate at which thermal energy alone lets molecules escape, applied to Mars.

- **The exobase:** mean free path equals scale height; above it, ballistic flight
- $v_{\text{th}} = \sqrt{2k_B T / m}$ against
 $v_{\text{esc}} = \sqrt{2GM / r_{\text{exo}}}$
- Count the molecules in the fast tail that point outward

A molecule faster than v_{esc} leaves; a slower one falls back.

The Escape Parameter

$$\lambda \equiv \left(\frac{v_{\text{esc}}}{v_{\text{th}}} \right)^2 = \frac{G M m}{k_{\text{B}} T r_{\text{exo}}}$$

- $\lambda \gg 1$: molecules gravitationally bound; escape rare
- $\lambda \lesssim 1$: thermal energy comparable to escape energy; rapid loss
- Depends strongly on molecular mass m ; independent of magnetic field, ionisation and solar wind

Result: The Jeans Formula

Integrating upward velocity over the upper hemisphere with cutoff $v \geq v_{\text{esc}}$:


$$\Phi_J = n \langle v \rangle \frac{1}{4} (1 + \lambda) e^{-\lambda}$$

where $\langle v \rangle = \sqrt{8k_B T / (\pi m)}$ is the mean thermal speed.

Substituting and simplifying:

$$\Phi_J = n \sqrt{\frac{k_B T}{2\pi m}} (1 + \lambda) e^{-\lambda}$$

Jeans (1925). Limits: $\lambda \rightarrow 0$ gives the effusion flux $n\sqrt{k_B T / (2\pi m)}$; for $\lambda \gg 1$ the flux falls as $\lambda e^{-\lambda}$, so λ from 5 to 10 cuts it by ~ 80 .



Maxwell-Boltzmann speed distributions of atomic hydrogen at the Earth (1000 K) and Mars (270 K) exobases: only the tail above the escape velocity leaves. Credit: Course-original figure.


Apply to Mars

$M = 6.4 \times 10^{23} \text{ kg}$, $r_{\text{exo}} \approx 3.6 \times 10^6 \text{ m}$, $T_{\text{exo}} \approx 270 \text{ K}$.

Escape velocity at the exobase: $v_{\text{esc}} \approx 4.9 \text{ km/s}$

Species	λ	$e^{-\lambda}$
H	5.3	5×10^{-3}
H ₂	10.6	$2-3 \times 10^{-5}$
He	21.2	$\sim 10^{-9}$
O	84.8	10^{-37}
CO ₂	230	10^{-100}

Mars cannot lose CO₂ thermally. Only H and H₂ escape via Jeans; heavier species need non-thermal channels.




Mars before and during the 2018 global dust storm, imaged by the Mars Reconnaissance Orbiter: lofted dust hid the surface and blocked solar power. Credit: NASA/JPL-Caltech/MSSS, public domain.



7. Mars Atmosphere and Climate Loss

Dendritic channel networks in the cratered Martian highlands, imaged by the Viking Orbiters. Credit: NASA/JPL/USGS


cold baseline (icy highlands): snow collects on the southern highlands



The early Mars climate problem, part 1: the cold baseline. Course figure after Wordsworth (2016), Wordsworth et al. (2017, 2021) and Kite & Conway (2024).


At 4 Ga the Sun gave $\sim 75\%$ of its modern flux and a pure CO_2 - H_2O atmosphere yields $T \leq 225$ K, too cold for the valley networks that need liquid water.

transient warm excursion, repeated: snow melts, runoff cuts valleys




The early Mars climate problem, part 2: one transient warm excursion. Course figure after Wordsworth (2016), Wordsworth et al. (2017, 2021) and Kite & Conway (2024).

Episodic H₂ release adds H₂-CO₂ collision-induced absorption that melts highland snow; runoff carves valley networks during model excursions of 10⁵ to 10⁶ yr.




Long-term climate regimes of Mars: mean surface temperature (divider ~ 280 K) against surface H_2O inventory (divider ~ 200 m GEL), with cross-sections. Credit: Wordsworth (2016), Fig. 7.

The four quadrants separate ice-capped highlands from lowland water; the geomorphology favours a cold, dry baseline with episodic melting.




A coupled atmosphere-surface-interior model of Mars' carbon inventory through time; initial CO_2 of 0.32–0.50 bar (blue) and 0.25–0.29 bar (red). Credit: Hu et al. (2015).

Late Noachian carbon stays below about 1 bar for most solutions (1.8 bar only with extensive carbonate sequestration), then goes to escape and carbonate burial: the modern 6 mbar is the remnant.




Climate state against CO₂ loss and lost non-CO₂ greenhouse forcing, from the high early valleys to the later low fans. Credit: Kite et al. (2022), Fig. 6.

The transition cooled Mars by ≥ 10 K, driven primarily by **waning non-CO₂ forcing** rather than by the thinning of the CO₂ atmosphere.




MAVEN oxygen escape channels extrapolated back to ~ 3.5 Ga. Credit: Jakosky et al. (2018).

Dissociative recombination dominates modern O loss, **pick-up ions** show the steepest historical decline, **ion outflow** and solar-wind **sputtering** add the rest; the present total escape is $\sim 2 \text{ kg s}^{-1}$, and the heavy species require these non-thermal channels.




Hydrogen corona column density at Mars over a Mars year: the escape varies by an order of magnitude and peaks near perihelion, when water vapour reaches the altitudes where it is photolysed. Jakosky et al. (2018).




8. Mars Magnetism


Early Mars from Mariner 4 (1965). Credit: NASA/JPL



No global dipole today, but intense remnant magnetisation in the southern highlands requires a vigorous early dynamo that shut down; the northern lowlands and the major basins were demagnetised by impacts.



Basaltic crust recorded the field as it cooled, possibly through reversals; the pre-Borealis crust preserves the primordial dynamo, whose cessation is dated to ~ 4.1 Ga (refined to ~ 3.7 Ga).




Dynamo lifetimes of the rocky planets: Earth active, Mercury weak in a thin liquid shell, Mars for its first 400 to 800 Myr, Venus with no detectable field. Course figure.

A magnetic shield may suppress ion escape by up to an order of magnitude over Gyr, but whether, and by how much, a field protects an atmosphere is debated; an unmagnetised Mars-sized planet still loses its atmosphere on 10^9 -year timescales.

9. Comparative Payoff: Mercury vs Mars






Mercury's magnetosphere, the smallest in the solar system, with its subsolar magnetopause at $\sim 1.5 R_{Hg}$; BepiColombo (ESA/JAXA) enters orbit in late 2026 with two orbiters to map the offset field, the core and the exosphere. Credit: Wicht & Heyner (2017).

Size and Distance Set the Trajectory

- **Small bodies cool fast:** surface-area-to-volume ratio $\propto 1/R$; Mars cooled faster than Earth, so its dynamo shut off early
- **Mercury keeps a weak dynamo:** light elements (S, Si) keep a thin outer-core shell liquid; thermal stratification gives the weak, offset field
- **Lesson:** a dynamo persists only while the core keeps convecting; Mars cooled too fast and never froze an inner core, Mercury keeps a thin liquid shell through light elements, Earth has both mantle pacing and inner-core buoyancy




Size and distance set the trajectory (schematic); the shaded regime boundary is qualitative.
Course figure.

Larger bodies keep their atmospheres, and their dynamos, for longer; size and distance govern core cooling, dynamos and volatile retention.

Summary

1. **Mercury:** 3:2 spin-orbit resonance, a ~ 74% iron core from mantle stripping, lobate scarps from $\Delta R \approx 7$ km of **contraction**, a weak offset dynamo
2. **Mars:** a liquid core of ~ 1650–1675 km below a molten silicate layer (2023 reanalyses), a wet Noachian record, and a dynamo that died between ~ 4.1 and ~ 3.7 Ga
3. **Jeans escape** $\Phi_j \propto (1 + \lambda)e^{-\lambda}$ lets H go but keeps CO₂; **non-thermal loss** (sputtering, ion pickup) removes the heavy species

Next: Lecture 11



Questions?

Next: **Lecture 11. Gas & Ice Giants**

Lecture notes: formingworlds.github.io/IntroductionToPlanetaryScience