

ميباكوفين

بيانات التركيب :

كل قرص يحتوي علي  
كينتوبروفين ٢٥ مجم  
كل قرص طويل المفعول يحتوي علي  
كينتوبروفين ١٥٠ مجم  
كل كبسولة ممتدة المفعول تحتوي علي  
كينتوبروفين ١٠٠ مجم  
كل ٥ ملل من المحلول المعلق تحتوي علي  
كينتوبروفين ١٢.٥ مجم  
كل أنبويه جيل ١٥ جم تحتوي علي  
كينتوبروفين ٢.٥ %

الخواص :

ميباكوفين ( كينتوبروفين ) هو مستحضر مضاد الالتهاب من الفئة الغير ستيرويدية والتي لها تأثيرات مسكنة وخافضة للحرارة ويعمل ميباكوفين كمضاد للإلتهاب ومسكن وخافض للحرارة عن طريق تثبيط بناء البروستاجلاندينز والليوكوترينز  
ميباكوفين له أسرع وأعلي تأثير بالنسبة لباقي المركبات غير السيترويدية كما انه سريع الامتصاص في الجهاز الهضمي

دواعي الاستعمال :

- كعلاج مساعد في حالات التهابات الأنف والأذن والحنجرة المصحوبة بألم
- الالتهابات المصحوبة بألم وأشكال الأمراض الروماتيزمية المختلفة : الروماتويد ، الأم المفاصل ، التهاب الفقرات التيبسي، التهاب مفاصل الفقرات
- الروماتيزم الالامفاصي
- حالات النقرص الحادة
- الحالات الالتهابية المصحوبة بألم في حالات أمراض النساء مثل ألم الطمث والتهاب الحوض
- الألم الناشئ عن مصادر أخرى مثل عرق النسا واللمباجو وآلام الأعصاب والتقلصات

الجرعة و الاستخدام:

للاستخدام عن طريق الفم :-

الالتهاب المفصلي ( روماتويد ) ، التهاب المفاصل المتخللة: يبدأ العلاج بجرعة ٧٥ مجم ثلاث مرات في اليوم أو ٥٠ مجم ٤ مرات يوميا أو ٢٠٠مجم مرة واحدة يوميا.  
الجد الأقصى للجرعة ٣٠٠ مجم يوميا

الأم الطمئي

٢٥ – ٥٠ مجم كل ٦ إلي ٨ ساعات اذا دعت الضرورة

للأطفال

لا ينصح بأستخدام ميباكوفين للأطفال أقل من ١٢ سنة حيث لم تجري دراسات كافية علي هذا السن بعد

الاستخدام الموضعي :-

يستخدم في المناطق المصابة ٢- ٣ مرات يوميا

نواهي الاستعمال

يجب عدم أستخدام ميباكوفين في الأشخاص اللذين لديهم حساسية للمستحضر وايضا للأسبرين ومضادات الالتهاب الغير ستيرويدية الأخرى  
يجب عدم أستخدام ميباكوفين في حالات الاصابه أو سابق الاصابه بقرحة المعدة أو الاثني عشر

الأعراض الجانبية

يتحمل الجسم ميباكوفين بشكل جيد فلا تحدث ايه تغيرات بولوجية غير مستحبة سواءً في الكبد أو الدم أو الكلوي ونادراً ما يحدث اضطرابات هضمية ( ألم بالمعدة ، غثيان )

الحمل والرضاعة

يجب أستخدام ميباكوفين بحذر في حالات الحمل والرضاعة

الاحتياطات الخاصة

- يجب ضبط جرعات ميباكوفين مع جميع الادوية التي لها قابلية اتحاد كبيرة مع بروتينات الدم عند الاستخدام المتزامن
- قد يسبب استخدام ميباكوفى حدوث نزيف بالجهاز الهضمي وقرحة المعدة وبعض اضطرابات الجهاز الهضمي
- في بعض المرضى قد يؤثر علي الكلوي وفي هذه الحالة يجب ضبط الجرعة

العصوة :

أقراص

عليه تحتوي علي ١٠ أقراص

اقراص ممتدة المفعول

عليه تحتوي علي ١٠ أقراص طويلة المفعول

كبسولات ممتدة المفعول

عليه تحتوي علي ١٠ كبسولات ممتدة المفعول

محلول معلق

رجاجه تحتوي علي ١٢٠ مل

حل

أنبويه تحتوي علي ١٥ جم

التخزين :

يحفظ في مكان جاف و في درجة حرارة لا تزيد عن ٣٠ ٥ م .

يحفظ بعيدا عن متناول الأطفال

Mepacofen

Composition:

Each **tablet** contains:  
• Ketoprofen 25 mg  
Each **Long acting tablet** contains:  
• Ketoprofen 150 mg  
Each **SR Capsule** contains:  
• Ketoprofen 100 mg  
Each **5 ml** contains:  
• Ketoprofen 12.5 mg  
Each **15 gm gel** contains:  
• Ketoprofen 2.5 %

Properties:

Mepacofen is a non-steroidal anti-inflammatory drug with analgesic and antipyretic properties.

The anti-inflammatory, analgesic, and antipyretic properties of Mepacofen are due to its inhibitory effects on prostaglandin and leukotriene synthesis, to have antibradykinin activity, as well as to have lysosomal membrane-stabilizing action. Mepacofen has a rapid onset of action and higher peak effect than other non-steroidal drugs; it is rapidly absorbed from gastrointestinal tract.

Indications:

- Mepacofen temporarily relieves cold, sinus and flu symptoms as nasal and sinus congestion, stuffy nose, headache, sore throat, minor body aches pains and fever.
- Painful inflammatory and degenerative forms of osteoarthritis, rheumatoid arthritis, ankylosing spondylitis and spondyloarthritis.
- Non-articular rheumatism.
- Acute attacks of gouty arthritis.
- Painful inflammatory gynecological conditions as dysmenorrhea and pelvic inflammatory diseases.
- Sciatica, lumbago, neuralgia and muscle cramps.

Dosage & Administration:

For Oral dosage forms:

- Osteoarthritis and Rheumatoid arthritis: starting dose is 75 mg three times daily or 50 mg four times daily or 200 mg administered once daily.
- Maximum daily dose is 300 mg/day
- Pain of dysmenorrhea: 25-50 mg every 6 to 8 hours as necessary.
- Pediatric use: not recommended for children under 12 years of age as studies are not yet preformed on this age group.

For Topical dosage form:

- To be applied over the affected area two to three times daily.

CONTRA- INDICATIONS:

- Mepacofen is contraindicated in patients who have shown hypersensitivity to it.
- Mepacofen should not be given to patients in whom aspirin or other nonsteroidal anti-inflammatory drugs induce hypersensitivity reactions.
- Active peptic ulceration or history of recurrent peptic ulceration.

SIDE EFFECTS:

Mepacofen is a highly tolerated drug with no significant change in hemogram or liver and kidney functions tests.

Generally the reported side effects were mild and transient in the form of digestive disorders as gastric pain, nausea and vomiting.

PREGNANCY & LACTATION:

Mepacofen should be used with caution in pregnant and lactating females.

PRECAUTIONS & WARNING:

- Mepacofen is highly protein-bound and might necessitate dosage adjustments if administrated concomitantly with protein bound drugs.
- GIT bleeding and ulceration can occur with or without warning symptoms; dyspepsia and other minor GIT symptoms occur commonly.
- In some patients, renal toxicity occurs, leading to reduction in renal blood flow and renal volume, so dose should be adjusted.

Package:

Tablets:  
Box containing 10 tablets.  
Long acting Tablets:  
Box containing 10 long acting tablets.  
SR Capsules:  
Box containing 10 capsules SR.  
Suspension:  
Bottle containing 120 ml.  
Gel:  
Tube containing 15 gm.

Storage:

Store in a dry place at temperature not exceeding 30°C.

Keep medications out of reach of children

ARE 31210011- Vrs1

Produced By

Arab Co.For Pharmaceuticals & Medicinal Plants

(MEPACO- MEDIFOOD)-[Enshas-Sharkeya]- EGYPT

Mepacofen gel is produced by Alexandria Company for

Pharmaceuticals and chemical industries for Arab Co. for

pharmaceutical & medicinal plants (MEPACO-MEDIFOOD)-

[Enshas-Sharkeya]-Egypt

إنتاج

الشركة العربية للأدوية و النباتات الطبية

(ميباكو - ميدي فود) –[انشاص-شرقية]- مصر

ميباكوفن جل انتاج شركة الأسكندرية للأدوية و الصناعات الكيماوية لصالح

الشركة العربية للأدوية و النباتات الطبية (ميباكو - ميدي فود)-

[انشاص-شرقية]- مصر

ARE 31210011- Vrs1