

Comparing Global Temperature Datasets: Evidence for a Significant Increase in Temperature in the Last 50 Years

Francesco Ragaini

On the last centuries the global temperature has changed and many different surveys have collected data about that.

This report intends to compare data from five different observers and to test if the anomaly in the last fifty years can be considered as a statistical fluctuation.

To achieve this goal, datasets from Nasa, HadCRUT, Japan Meteorological office, Berkeley earth, Ncdc NOAA, these datasets have been compared with a Kalmogorv-Smirnov test, then all of that has been split in data before and after 1975 and compared again with KS test.


The Kalmogorv-Smirnov test on the entire datasets has reported positive results (figure 1), conversely the K.S. test on the split dataset has reported negative results (figure 2).

This result suggests that all the datasets analyzed describe the same effect and the timeseries of the anomalies after the 1975 and before 1975 have different distributions. According to this result, all surveys registered the same effect and in the last fifty years the temperature has increased.

Data analysis

All the surveys analyzed in this study have released their datasets on their respective websites (see bibliography). The datasets are organized as dataserries of annual anomalies: discrepancy of annual global temperature from a certain baseline, considered as zero point. However, each dataset considers a different temperature as the zero point. Therefore, the first step is to adjust the datasets by applying an offset to set the zero point at the same temperature. Only after this correction, Kalmogorv-Smirnov tests can be applied to the datasets.

Data comparison



	HadCRUT	Nasa	japan Met.	Noaa	Berkeley
HadCRUT	1.000	0.124	0.097	0.288	1.000
Nasa	0.124	1.000	0.632	0.955	0.217
japan Met.	0.097	0.632	1.000	0.751	0.051
Noaa	0.288	0.955	0.751	1.000	0.288
Berkeley	1.000	0.217	0.051	0.288	1.000

FIG. 1 All surveys's datasets are consistent with the others by K.S. test. The plot represents normalized histogram with annual anomalies divided by survey. In the table are reported the p values.

As shown in figure 1, all anomaly datasets exhibit a high degree of consistency with each other, as evidenced by the K.S. p values reported in the table. While the Japan Met. dataset appears to be the most different from the

others, its p value respect to all the other datasets is still greater than 0.05. Overall, these findings suggest a high level of agreement among the different surveys regarding the measurement of global temperature anomalies.




FIG. 2 Anomalies before and after 1975 are not consistent. Each plot is an histogram that represent anomalies distributions before and after 1975. In the table are reported the p values by K.S. tests.

Figure 2 shows that all datseries before and after 1975 are not compatible by K.S. tests, as every p-value is under 0.05. The year 1975 has been chosen beacouse it shows a significant change in trend.

Conclusions

According to the first result, (fig 1) is possible to assert that different surveys are misuring similar distributions, so can be concluded that all of that are misuring the same effect. Japan Met. data are the most different to the others, this could appen because Japan Met. zero period was the most different.

In the second claim, (fig 2) is shown that the anomalies in the last fifty years can't be considered as statistical fluctuation. According to that result can be interesting to search for a trend in the last fifty years of data.

Bibliography

Nasa	HadCRUT	Japan Metereological office	Ncdc Noaa	Berkeley
Nasa dataset	HadCRUT dataset	Japan Met. dataset	Noaa dataset	Berkeley dataset
Base: 1951-1980	Base: 1961-1990	Base: 1991-2020	Base: 1901-2000	Base: 1951-1980