

From the Euclidean Gas to Quantum Yang–Mills and a Mass Gap

A research prospectus with proof sketches and theorem dependencies

Abstract. This prospectus organizes the constructive program in Volume 2 around one stochastic transition and its recorded history law. A marked interacting gas supplies a nonlinear population evolution. Confinement, Poincaré and logarithmic Sobolev inequalities, hypocoercivity, propagation of chaos, and normalized regularity estimates control its long-time and population limits. A lossless Fractal Set representation transports the same law to gauge and matter observables. Native likelihood, source-response, fluctuation, and continuum-consistency results then supply the field construction. The final tasks identify a nontrivial physical Euclidean Yang–Mills hierarchy, reconstruct its transfer Hamiltonian, and carry a uniform self-adjoint gap through the cutoff and volume limits.

The document gives the mechanism of each stage and identifies the theorem statements in the source. It follows the existing population, mechanical, metric, gauge, and matter constructions, including the derived sampling normalizations and temporal correlations. The field-theoretic reconstruction statements retain their explicit hypotheses on the same selected equilibrium law.

Source version: repository snapshot, 1 October 2026.

Purpose: mathematical discussion of the complete construction program.

Source corpus: docs/source/2_fractal_gas/.

Contents

Objects, target, and logical structure	1
1 Fix the Euclidean gas	1
2 Resolve cloning and collision geometry	2
3 Control confinement, moments, and tails	3
4 Select the stationary or conditioned ensemble	5
5 Derive the population law and exact field equations	6
6 Control particles, observation time, and stationarity	7
7 Build Poincaré and logarithmic Sobolev control	9
8 Turn functional inequalities into relaxation and fluctuations	10
9 Obtain smooth empirical and mean-field fields	11
10 Reconstruct the causal 3+1 Fractal Set	13
11 Construct gauge and matter observables	15
12 Derive the native field action and response	16
13 Construct the recorded quantum matter algebra	19
14 Retain fluctuations and a common correlation hierarchy	20
15 Control the continuum operator with derived tools	22
16 Pass actions, sources, and correlations through the limits	24
17 Reconstruct fields from the equilibrium history	25
18 Preserve the physical Yang-Mills mass gap	28
19 How the proved walker dynamics supply the field theory	29
19.1 Probability law, invariant descriptors, and frame observables	29
19.2 The history correlations retain the algorithmic clock	29
19.3 Geometric integrals and fluctuations have derived normalizations	30
19.4 Native likelihood and conditional geometry	30
19.5 Equilibrium representation and the gap	30
20 The estimates that control the continuum	31
A Source theorem register	32
B Chapter contribution map	39

Objects, target, and logical structure

The construction begins with three spatial coordinates and algorithmic time:

$$x_i(n), v_i(n) \in \mathbb{R}^3, \quad t_n = n\Delta t, \quad e_{n,i} \mapsto (t_n, x_i(n)).$$

Here Δt is the configured recording spacing with its fixed unit calibration. CST edges inherit increasing algorithmic time. Equilibrium supplies the selected history law in which the additional field-theoretic properties are developed. The target is a nontrivial quantum Yang–Mills theory on this $3 + 1$ reconstruction, with a unique vacuum and a positive physical energy gap. The construction proceeds through

$$\begin{aligned} \text{gas} &\longrightarrow \text{population control} \longrightarrow \text{recorded field law} \\ &\longrightarrow \text{continuum hierarchy} \longrightarrow (\mathcal{H}, \Omega, H). \end{aligned}$$

Law ledger. Write $P_{N,h}$ for the complete conservative transition and $Q_{N,h}$ for its specified killed counterpart. Their selected laws are respectively π_N and ν_N . The nonlinear population update is $\mathcal{F}_h$. A history descriptor $\mathcal{D}$ induces the field law $\mathcal{D}_*P$. Every transfer of an estimate uses an explicit identification of these laws, their conditionings, or their observables.

Scale ledger. Here N counts walkers, M counts observations in a continuum estimator, h is the algorithm step, n is the update index, ε is a reconstruction bandwidth, a is a field cutoff, and L is a physical volume scale. The continuum chapter uses N for its sample count; this prospectus writes M to distinguish it from population size. This roadmap takes $d = 3$ and $D = d + 1 = 4$. The fourth coordinate is the calibrated recorded time. A phase-space fluctuation test also retains $v \in \mathbb{R}^3$.

Reading convention. “Source results” name statements and proofs already present in Volume 2, with their hypotheses retained. “Proof mechanism” sketches the argument of those results and their composition. “Stage output” identifies the object or estimate passed to the next stage. The final assembly criterion organizes these dependencies.

The main interface. Population-uniform coercivity and relaxation are transported through the recorded observable law. The physical transfer Hamiltonian must then be identified and controlled uniformly in the cutoff and volume family. That identification connects the gas estimates to the Yang–Mills mass gap.

1 Fix the Euclidean gas

Begin with the complete marked swarm $S = ((x_i, v_i, a_i))_{i=1}^N$ with positions, velocities, and alive/dead status. Dead slots retain their coordinates. Freeze the input frame for measurement, fitness, donor choice, and acceptance; this fixes the update order and prevents updated donor coordinates from leaking into the same cloning stage.

The canonical update samples one measurement companion, computes regularized population statistics and positive logistic fitness, samples a cloning donor, and accepts according to the frozen score. Accepted edges define connected collision components. Copy recipient positions with Gaussian jitter, rotate each component about its center-of-mass velocity, apply BAOAB and final position noise, cap velocities, and classify the terminal boundary once.

Declare the absorbing or conservative configuration, the objective and force, the noise laws, the internal marks, and all parameter floors. If historical donors or mutable providers are used, include their memory in the complete state. This stage fixes the stochastic object used throughout the program.

$$E = \mathbb{R}^d \times \overline{B}_V \times \{0, 1\}, \quad S \in E^N, \quad L_N = \frac{1}{N} \sum_{i=1}^N \delta_{z_i}, \quad z_i = (x_i, v_i, a_i).$$

Proof mechanism

Construct the companion and donor kernels by normalizing strictly positive weights against the alive population. The regularized standard deviations keep the score maps defined on constant populations. Condition on the frozen marks and accepted-edge graph, then compose the collision, Gaussian, deterministic kinetic, cap, and classification kernels. Integrating these conditional kernels gives $P_{N,h}$. Its path law is the initial law times the ordered product of the complete transition kernels. Record coverage is fixed at this point, so later likelihood calculations integrate the randomness that actually produced the path.

Analytic inputs

Marked Markov state; positive fitness and normalization floors; bounded comparison features; Gaussian innovations; Feller regularity. The Euclidean Gas Update is the defining algorithm in `convergence_program/02_euclidean_gas.md`, label `alg-euclidean-gas`.

Stage output. A complete finite-particle transition and a record contract that specifies every random choice and intermediate stage. Subsequent mean-field and field constructions use this kernel.

2 Resolve cloning and collision geometry

An accepted donor edge changes the recipient position and joins its endpoints into a collision component. Overlapping groups form a single connected component. The shared orthogonal rotation acts on the component relative velocities, so each walker has one velocity destination.

The tagged-walker population description must retain its random collision neighborhood. Strict fitness increase along accepted live edges supplies component control in the canonical current-frame construction. Rooted component convergence connects these finite neighborhoods to the limiting collision law.

Compute momentum, restitution energy, and the cross-walker covariance created by the shared rotation. Retain these terms in fluctuation and mean-field calculations: the noises of members of one component are correlated.

$$\tilde{v}_i = \bar{v}_C + \alpha R_C(v_i - \bar{v}_C), \quad R_C \sim \text{Haar}(O(d)), \quad \sum_{i \in C} \tilde{v}_i = \sum_{i \in C} v_i.$$

Proof mechanism

For a component C , the centered velocities satisfy $\sum_{i \in C} (v_i - \bar{v}_C) = 0$. Orthogonality preserves their squared norm and Haar symmetry gives zero mean for the rotated component. Taking traces in the rotationally invariant covariance identity gives

$$\text{Cov}(\tilde{v}_i, \tilde{v}_j \mid C) = \frac{\alpha^2}{d} [(v_i - \bar{v}_C) \cdot (v_j - \bar{v}_C)] I_d.$$

For the population limit, truncate the rooted accepted-edge component, compare its finite exploration law with the limiting rooted law, and remove the truncation using the proved component moments. The two-root calculation controls the empirical variance. These steps identify the collision term without replacing component rotations by independent row noise.

Analytic inputs

Accepted-edge graph; rooted-component truncation; moment bounds at every component order; shared Haar rotation; self-exclusion correction; two-root independence in the population limit.

Source results

- **Component momentum, energy, and shared covariance** [R1]. Component momentum, energy, and shared covariance.
- **One-step collision consistency** [R2]. One-step collision consistency for the rooted population law.

Stage output. The actual collision law and its conservation, moment, and covariance data. These feed the fixed-step population map and its fluctuation equations.

3 Control confinement, moments, and tails

Combine cloning displacement with kinetic transport to derive a Foster-Lyapunov estimate for the complete update. Choose positive weights for the component drift inequalities and evaluate the resulting rate and source. Iterate the drift to obtain finite-time and stationary moment budgets.

The structural landscape chapter also derives selection-driven confinement from accepted inward and outward donor fluxes. Its complete moment inequality retains the selection excess and the kinetic force-growth coefficients. It provides an explicit criterion for contracting moments without a prescribed restoring trap.

Use these moment bounds to localize unbounded reward estimates, control population coverage failures, obtain tightness, and justify uniform integrability. The entropy-to-tail and population-fraction bounds give a second quantitative route to controlling the mass of poorly covered or distant regions.

$$\begin{aligned} P_N W_p &\leq r_p W_p + B_p + a_p E_p, \quad r_p < 1, \\ \mathbb{E} W_p(S_n) &\leq r_p^n \mathbb{E} W_p(S_0) + \frac{B_p + a_p \bar{e}_p}{1 - r_p} (1 - r_p^n). \end{aligned}$$

Proof mechanism

Evaluate the copying increment $|x_j|^p - |x_i|^p$ under accepted donor flux. Lower bounds on inward core flux and upper bounds on outward tail flux give a signed selection drift. Add cloning jitter through Gaussian radial moments, then propagate through the actual kinetic position update using the force-growth and cap bounds. The remainder E_p records coverage or class failures. Iterating the resulting affine recurrence gives the displayed moment budget. Markov’s inequality turns higher moments into quantitative tail localization; the entropy-floor calculation gives additional tail and population-fraction estimates for the normalized joint positional law.

Analytic inputs

Foster-Lyapunov drift; positive-weight spectral criterion; Gaussian radial moments; regional selection flux; coverage and tail budgets; entropy confinement floor.

Source results

- **Foster-Lyapunov drift for the composed kernel** [R3]. Composes the drift estimates at the level of the actual kernel.
- **Fully evaluated selection-driven moment drift without a prescribed restoring trap** [R4]. Evaluates selection-driven moment drift, including the zero-restoring-trap regime.
- **Entropy contraction to an explicit confinement floor** [R5]. Converts verified selection into entropy contraction toward an explicit confinement floor.

Structural transport and mass–shape control

The positional transport chapter derives the centered structural control from the cloning variance reset and carries the barycenter separately in its full W_2 decomposition. The structural landscape chapter evaluates the signed Keystone, collision, and kinetic terms for the complete update. These results retain the inward selection flux, the donor-normalization costs, and the actual force increments in the same one-step ledger.

The mass, Hellinger, and transport chapter combines the alive mass with normalized shape. For finite measures $m\rho$ and $m_*\rho_*$, it proves the exact mass–shape identity

$$H^2(m\rho, m_*\rho_*) = (\sqrt{m} - \sqrt{m_*})^2 + \sqrt{mm_*} H^2(\rho, \rho_*).$$

It combines this with entropy-to-Hellinger and entropy-to-transport inequalities and also bounds the canonical Hellinger–Kantorovich distance. Safe-walker survival, concentration of alive mass, and convergence of the conditioned shape therefore control separate parts of the finite-measure error. This is the population normalization used in the convergence analysis.

Results supplying these constructions

- **Structural/Barycenter Split for Full W_2** [R6]. Carries the barycenter and centered structural terms in full transport control.
- **Joint convergence of mass and shape** [R7]. Assembles convergence of alive mass and normalized shape.

- **Finite-horizon survival from safe-walker estimates** [R8]. Controls finite-horizon survival through the safe-walker estimates.

Stage output. Population-uniform physical moment and tail controls for the chosen parameter regime. These support tightness, stationary limits, smooth reconstruction, and control of field moments.

4 Select the stationary or conditioned ensemble

For the conservative process, the drift and common-mass estimates give convergence to its identified invariant law. For the killed process, the surviving-block results give the selected QSD and its survival eigenvalue. The finite-window and Doob constructions specify which conditioned history is observed.

The recorded observation time is $t_n = n\Delta t$, where Δt includes any configured recording stride and fixed physical-unit calibration. Burn-in advances this same dynamics to the selected equilibrium ensemble. Subsequent windows retain their algorithmic time differences and all recorded transition data.

The Yang–Mills chapter computes the selected QSD window weights and its relaxation error. The equilibrium law supplies the starting ensemble; the complete transition supplies every temporal correlation used below.

$$\pi_N P_{N,h} = \pi_N, \quad \nu_N Q_{N,h} = \alpha_N \nu_N, \quad 0 < \alpha_N \leq 1.$$

Proof mechanism

Iterate the weighted drift to enter the controlled region, then apply the appropriate common-mass or surviving-block comparison. In the conservative case, the coupling contracts a weighted distance and yields an invariant probability. In the killed case, normalized block iteration yields the selected eigenmeasure. For a QSD window, multiply the killed transitions and divide by the window survival probability. Successive survival factors cancel in the selected-history identity. Comparing an actual relaxed window with this eigenmeasure window introduces the explicit survival amplification, bounded in the chapter by $2\alpha_N^{-w}\delta_n$ for a w -step window.

Analytic inputs

Weighted Harris coupling; surviving-block minorization; eigenmeasure identities; survival normalization; total-variation window comparison.

Source results

- **A weighted coupling form of Harris’ argument** [R9]. Conservative mixing from the declared drift and coupling inputs.
- **QSD convergence from a two-sided surviving-block bound** [R10]. QSD convergence from a two-sided surviving-block bound.
- **Relaxation to the selected law of a recorded gauge window** [R11]. Transfers the gas relaxation rate to a selected recorded gauge window.

Stage output. A selected long-time law and its exact history-window law, together with burn-in and conditioning errors. Field reconstruction takes expectations under this specified ensemble.

5 Derive the population law and exact field equations

Pass from the swarm empirical measure to a nonlinear population law on marked one-slot states. Keep the alive restriction and its normalized donor law distinct from the full population law, because dead positions and velocities affect revival and collision behavior.

The construction first samples the measurement mark, then applies population standardization and fitness, then samples accepted edges and the limiting collision neighborhood. It finally composes the actual kinetic stages and terminal classification. Averaging the measurement before its nonlinear acceptance step would change the map.

The resulting population equation is discrete at the fixed configured timestep. Derive its mass balance, weak field balances, positive alive mass, and finite-horizon moments. Stationary solutions are fixed points of this map. A continuous-time generator requires its own local consistency calculation if h is later changed.

$$\mu_{n+1} = \mathcal{F}_h(\mu_n), \quad m(\mu) = \mu(a = 1), \quad \rho_\mu = \frac{\mu^a}{m(\mu)}.$$

Proof mechanism

Disintegrate a representative row into its measurement mark, frozen fitness, donor gate, rooted collision component, and independent kinetic innovations. Integrate the root output against this product of conditional laws to define $\mathcal{F}_h$. Positivity and total mass follow from kernel composition. Gaussian final position noise makes terminal box classification continuous after averaging and supplies positive alive mass. Polynomial growth of reward is handled by the propagated moments. The weak mass and field balances follow by testing the same root input-output coupling; each term belongs to a declared stage of the complete update.

Analytic inputs

Normalized companion integrals; measurement marks; finite limiting collision neighborhoods; exact composition of BAOAB, diffusion, cap, and classification.

Source results

- **The fixed-step mean-field equation** [R12]. The exact fixed-step mean-field equation.
- **Exact mass and weak field balances** [R13]. Exact mass and weak field balances for the full population update.

The finite algorithm already has field equations

The field-equations chapter derives the hierarchy directly from the configured donor, cloning, innovation, and boundary maps. Its full marked state retains current and historical donor

data, eligibility, provider state, and the inputs needed by the next update. On this state, its marked field and age transport are exact.

For any square-integrable vector observable A , the complete kernel gives

$$\begin{aligned} b_A(S) &= P_{N,h}A(S) - A(S), \\ \Gamma_A(S) &= P_{N,h}(AA^\top)(S) - P_{N,h}A(S)P_{N,h}A(S)^\top, \\ A(S_{n+1}) - A(S_n) &= b_A(S_n) + \eta_{n+1}, \end{aligned}$$

with $\mathbb{E}[\eta_{n+1} \mid S_n] = 0$ and conditional covariance $\Gamma_A(S_n)$. The same formula applies to metric probes and gauge readouts. Fourier tests and the marked characteristic functional give the exact moment and field-correlation hierarchy, retaining the configured innovation law and cross-walker dependence.

For a row transition $(x, v, a) \mapsto (y, w, b)$, deposit along the segment from x to y . The fundamental theorem of calculus gives the distributional density and current balances. Summing over the actual stages telescopes to

$$\rho_{n+1} - \rho_n + \operatorname{div} \mathcal{J}_\rho = \mathcal{S}_\rho, \quad j_{n+1} - j_n + \operatorname{div} \mathcal{J}_j = \mathcal{S}_j.$$

All source and transport terms are computed from the executed update. The coupled-field theorem combines these mechanical equations with the population, history, Hessian metric, and thermostat covariance. Configured algorithm variants retain their own stage ordering and coefficients in these identities.

Results supplying these constructions

- **Explicit finite-step population law** [R14]. Explicit finite-step population law for the configured complete state.
- **Exact field characteristics and moment hierarchy** [R15]. Exact characteristics, characteristic functional, and moment hierarchy.
- **Finite-step observable equation and fluctuation covariance** [R16]. Exact drift and conditional fluctuation covariance of derived observables.
- **Exact discrete spatial density and momentum equations** [R17]. Exact finite-step spatial density and momentum balances.
- **Coupled population, mechanical, and metric fields** [R18]. The coupled population, mechanical, and metric field system.

Stage output. The nonlinear population update, its sampled interaction law, and its exact balance relations. This is the background law for the smooth field and fluctuation constructions.

6 Control particles, observation time, and stationarity

Prove one-step consistency of the finite update with the nonlinear population map, including measurement concentration, collision-component convergence, and boundary/revival contributions. Iterate to obtain finite-horizon propagation of chaos.

For long times, use the structural landscape results appropriate to the declared regime. The weighted active-contraction branch provides a unique stationary population law and a population-independent attraction rate. Its trajectory corollary supplies uniform-time

approximation, stationary chaos, and commuting population/time limits with explicit initial moment control.

The growing-horizon branch instead supplies trajectory approximation through a diverging observation horizon using higher moments and the actual consistency constants. This provides a controlled joint population/time limit without assuming attraction. Keep the branch used by the field ensemble explicit.

$$\begin{aligned} |\mathbb{E}[L'_N \varphi \mid S] - \mathcal{F}_h(L_N(S))\varphi| &\leq \frac{2\|\varphi\|_\infty B_*}{\sqrt{N}}, \\ \mathbb{E}\left[|L'_N \varphi - \mathcal{F}_h(L_N(S))\varphi|^2 \mid S\right] &\leq \frac{A_\varphi + 4\|\varphi\|_\infty^2 B_*^2}{N}. \end{aligned}$$

Proof mechanism

Condition on the input array and estimate the realized measurement-normalization fluctuations. Couple the resulting donor and gate laws, use rooted-component estimates for the collision response, and apply conditional concentration to independent kinetic innovations. This proves the displayed one-step bias and mean-square bounds at the actual empirical input. Continuity of $\mathcal{F}_h$ and a countable determining class give finite-horizon iteration. For stationary limits, combine one-step consistency with population attraction and tightness. In the weighted-contraction branch, expand two iterates and verify the source condition

$$q_w = 1 - \epsilon_2/2 + 2B_w L_{\text{rem}} + L_{\text{rem}}^2 < 1.$$

The growing-horizon branch instead propagates its explicit error recursion and high-moment budget to a diverging observation horizon.

Analytic inputs

Rooted collision convergence; one-step bias and mean-square consistency; conditional concentration; weighted TV contraction; finite-row chaos; growing-horizon error recursion.

Source results

- **Finite-horizon propagation of chaos for the actual update** [R19]. Finite-horizon propagation of chaos for the actual update.
- **Closed full-law convergence with unbounded raw reward** [R20]. Closed full-law population convergence with unbounded raw reward.
- **Uniform-time trajectories, stationary chaos and commuting limits** [R21]. Uniform-time trajectories, stationary chaos, and commuting limits.
- **Uniform trajectory approximation through a diverging horizon** [R22]. A separate explicit trajectory estimate through a diverging horizon.

Exchangeability connects the empirical law to tagged fields

Kernel permutation symmetry and uniqueness give exchangeability of the selected finite law. The exchangeability chapter derives the finite empirical-mixture representation and proves that

a deterministic empirical limit gives fixed-row chaos. It also evaluates the finite-population covariance exactly:

$$\text{Cov}(g(Z_1), h(Z_2)) = \frac{N \text{Cov}(L_N g, L_N h) - \text{Cov}(g(Z_1), h(Z_1))}{N - 1}.$$

The empirical variance bound therefore controls distinct-walker correlations with the finite-size diagonal term retained. This supplies the passage between population observables and fixed-row invariant channels used in field reconstruction.

Results supplying these constructions

- **Exchangeability of a unique QSD** [R23]. Exchangeability of the unique selected QSD.
- **Deterministic empirical limits imply chaos** [R24]. Deterministic empirical limits imply chaos.
- **Exact covariance identity for finite populations** [R25]. Exact covariance identity for a finite exchangeable population.

Stage output. Quantified control of the population approximation over the observation regime used in the reconstruction. This controls both finite-run correlations and the approach to the selected limiting background.

7 Build Poincaré and logarithmic Sobolev control

The confinement route begins with a quadratic Lyapunov inequality. A weighted energy estimate controls the exterior region, while a local Poincaré inequality controls the interior. Together they produce a global Poincaré bound. A defective LSI is then tightened using that Poincaré inequality.

For the kinetic reference law, combine the spatial inequality with the Gaussian velocity inequality by tensorization. For joint swarm laws, Volume 2 supplies four explicit N -uniform routes: a product reference, a uniformly bounded joint tilt, uniform joint curvature, or a contractive additive-noise invariant flow.

Identify which route applies to the selected joint law and retain its constant C_* . Include the status-stratum entropy term when discrete alive/dead variables are part of the law. This supplies the functional inequality used for the interacting swarm and downstream reconstructed observables.

$$\text{Ent}_{\pi_N}(f^2) \leq 2C_* \int \sum_{i=1}^N (|\nabla_{x_i} f|^2 + |\nabla_{v_i} f|^2) d\pi_N.$$

Proof mechanism

Write $\pi(dx) = Z^{-1} e^{-V(x)} dx$, assume $\nabla^2 V \succeq -KI$, and use the quadratic Lyapunov estimate. Choose R so $\ell = cR^2 - b > 0$. The exterior weighted-energy estimate and interior Poincaré inequality give

$$C_P = \left(1 + \frac{b}{\ell}\right) C_{\text{loc}} + \frac{1}{\ell}.$$

The entropy–transport–Fisher estimate and the moment bound yield a defective LSI. Centering by Rothaus’ inequality removes the defect using C_P . Tensorization then incorporates Gaussian velocities. At the joint level, the chosen product, bounded-tilt, curvature, or flow-contraction criterion controls the constant independently of N . The selected invariant law or QSD must be the law appearing in that criterion.

Analytic inputs

Lyapunov weighted energy; local-to-global Poincaré; defective-LSI tightening; Bakry–Émery curvature; tensorization; bounded-density perturbation; flow contraction.

Source results

- **LSI from a quadratic Lyapunov bound** [R26]. LSI from a quadratic Lyapunov bound, including the global Poincaré step.
- **LSI for the kinetic reference measure** [R27]. LSI for the kinetic reference measure.
- **N -uniform LSI for specified joint laws** [R28]. N -uniform full-gradient LSI for the specified joint-law families.

Stage output. A declared joint-law LSI with an explicit population-uniform constant. Its Poincaré consequence and concentration estimates are used next; its coercivity is retained for the eventual field Hamiltonian identification.

8 Turn functional inequalities into relaxation and fluctuations

Linearize the joint LSI at the constant function to obtain Poincaré. Applied to a Lipschitz empirical average, the gradient sum is of order N^{-1} . The variance therefore scales as $C_* L^2/N$, even when the walkers are dependent.

For evolution, combine the LSI with the kinetic Fisher-information identities and dissipation of modified entropy. Estimate the actual cloning kernel and the killing normalization in the same balance. The full normalized evolution result and the Doob-transform route yield entropy relaxation for their identified law families.

At discrete time, use the full-step survivor chain rule and numerical entropy-defect estimates for the actual kernel. Relative entropy controls bounded-observable bias through Pinsker, while concentration and Poincaré control sampling fluctuations. Fixed marginals inherit the same LSI constant, and the inequality passes to their limits.

$$\mathrm{Var}_{\pi_N}(F) \leq C_* \int \sum_i |\nabla_i F|^2 d\pi_N, \quad \mathrm{Var}_{\pi_N} \left(\frac{1}{N} \sum_i \varphi(z_i) \right) \leq \frac{C_* \mathrm{Lip}(\varphi)^2}{N}.$$

Proof mechanism

Substitute $f = 1 + \epsilon F$ in LSI and compare the second-order terms to obtain Poincaré. For $F_N = N^{-1} \sum_i \varphi(z_i)$, the squared gradient sum is bounded by $\mathrm{Lip}(\varphi)^2/N$. For entropy decay, differentiate the modified kinetic functional and absorb its cross-gradient terms by the chapter’s

quadratic-form inequalities. Add the cloning contribution and the conditioned normalization term. A closed negative derivative bound followed by Grönwall gives the stated relaxation estimate. Projection to fixed marginals preserves LSI; bounded continuous test integrands pass the inequality to their weak limits.

Analytic inputs

Poincaré linearization; full Fisher information; modified hypocoercive entropy; conditioned entropy identity; Doob transform; Pinsker and entropy transport; exact discrete entropy accounting.

Source results

- **Poincaré inequality and empirical observables** [R29]. Poincaré inequality and empirical variance control.
- **Kinetic hypocoercive entropy convergence** [R30]. Kinetic hypocoercive entropy convergence.
- **Entropy convergence for the full normalized swarm evolution** [R31]. Entropy convergence for the full normalized swarm evolution.
- **LSI for fixed marginals and their limits** [R32]. LSI inheritance by fixed marginals and their limits.

Stage output. Relaxation, concentration, and moment tools under the same selected law. These control burn-in, empirical observables, distribution-valued fluctuations, and later continuum test-function estimates.

9 Obtain smooth empirical and mean-field fields

Differentiate the normalized companion kernels before differentiating the fitness. Volume 2 tracks numerator derivatives, normalization derivatives, moments, variance, standardized scores, and smooth fitness maps. Positive floors control denominators throughout the calculation.

The C^3 chapter supplies force and finite-order derivative bounds. The all-order chapter extends the normalized calculus with recurrences and majorants, retaining bandwidth, regularizer, and population-count dependence. Smooth localization and density estimates connect these quantities to geometric coefficients.

The normalized mean-field integral theorem replaces sums by integrals with the same derivative recurrence. The smooth empirical-field convergence theorem then upgrades weak population convergence to C^n convergence on compact parameter sets for each fixed order, under its denominator and equicontinuity inputs.

$$A_\mu(x) = \int a(x, y) \mu(dy), \quad F_\mu(x) = \frac{\int a(x, y) m(x, y) \mu(dy)}{A_\mu(x)},$$

$$\mu_N \Rightarrow \mu, \quad \inf_K A_\mu > 0 \quad \implies \quad F_{\mu_N} \longrightarrow F_\mu \text{ in } C^n(K),$$

Proof mechanism

Start with $A_\mu w_\mu = a$. Differentiating this identity isolates the highest derivative of the normalized weight; integrating its norm gives the all-order recurrence. The same Leibniz calculus handles measurement moments and regularized variance. On a compact parameter set, a finite equicontinuity net upgrades weak convergence of the constituent integrals to uniform convergence of each derivative. A positive limiting denominator gives a common lower bound for the empirical denominators. Induction through the quotient identity then proves C^n convergence. The result retains dependence on bandwidth and regularization parameters, which must be tracked along any changing-scale family.

Analytic inputs

Normalized-weight derivative recurrence; exact normalization cancellations; regularized variance calculus; dominated differentiation; equicontinuity; compact parameter nets; spectral bounds for reconstructed coefficients.

Source results

- **C³ regularity of sampled, surrogate, and expected fitness** [R33]. Third-order regularity of the fitness construction.
- **Full sampled-fitness majorant** [R34]. All-order regularity of the complete fitness potential.
- **Differentiating normalized mean-field integrals** [R35]. Differentiation of normalized mean-field integrals.
- **Smooth convergence of normalized empirical fields** [R36]. Smooth convergence of normalized empirical fields.

Fitness regularity produces the adaptive geometry

The emergent-geometry chapter identifies the implemented positive spectral branch

$$g = \epsilon_g I + (H_{\text{sym}})_+, \quad D = g^{-1}, \quad B = \sqrt{2\gamma T} g^{-1/2}.$$

The fitness derivative hierarchy controls the Hessian and its variation. Spectral lower and upper bounds then control metric inversion, adaptive diffusion, and the noise form. The geometric-gas chapter transfers moment drift with explicit perturbation costs and provides Harris, joint-LSI, concentration, and tail results for its identified geometric law families.

For reconstructed matrices $g, \hat{g}$, the Yang–Mills chapter uses the spectral margin to bound inverse and square-root errors:

$$\|\hat{g} - g\|_F \leq \delta, \quad \|\hat{g}^{-1} - g^{-1}\|_F \leq \epsilon_g^{-2} \delta, \quad \|\hat{g}^{-1/2} - g^{-1/2}\|_F \leq \frac{\delta}{2\epsilon_g^{3/2}}.$$

The normalized geometric-weight comparison is evaluated on the same recorded history. These estimates transport the existing coefficient regularity into metric, volume-weight, and field-readout accuracy.

Results supplying these constructions

- **Metric represented by the implemented Hessian branch** [R37]. Identifies the metric represented by the implemented Hessian branch.
- **Uniform ellipticity from metric bounds** [R38]. Ellipticity from the established metric bounds.
- **Uniform ellipticity from the spectral margin** [R39]. Geometric uniform ellipticity from the spectral margin.
- **N -uniform LSI for an identified geometric law** [R40]. Joint population-uniform LSI for the identified geometric law.

Stage output. Controlled smooth coefficient fields and convergence of their derivatives. These supply the Taylor, metric, force, and local-operator regularity inputs to the continuum construction.

10 Reconstruct the causal 3+1 Fractal Set

Encode the complete history as a directed two-complex. CST edges record temporal transport, IG edges record selection interactions, IA edges record influence attribution, and interaction triangles carry ordered transport products. Specify the oriented incidence data used by each face.

Lossless reconstruction recovers trajectories, force and diffusion data, population statistics, landscape information, and cloning events from the covered record. Frame changes transport the coordinate description while retaining the invariant readouts.

Use the direct observable construction to pass to Gram and determinant coordinates for the internal orbits. Its measure isomorphism retains expectations and correlations of invariant observables. The recorded transition is intertwined with the original update; compressed channels retain exact eliminated-coordinate memory until their predictive completion is constructed.

$$\mathcal{D}_*P = \nu, \quad \mathbb{E}_P[O(\mathcal{D}(\omega))] = \int O(y) \nu(dy).$$

Proof mechanism

Apply the lossless decoder to a covered record, reconstruct the stage variables, and substitute them into the complete kernel. This identifies finite histories and their observable expectations. For invariant coordinates, the orbit theorem resolves the frame equivalence and the measure theorem pushes forward both probability and readouts. The transition intertwining follows by applying the same kernel to pullbacks of descriptor observables. If a descriptor is compressed, conditional elimination gives its exact memory term. Closing the original bounded insertion algebra under repeated transition action gives the prediction-complete descriptor; finite conditional partitions approximate its transition matrices.

Analytic inputs

Exact vector encoding; oriented two-complex; lossless decoder; invariant coordinates; orbit and measure isomorphisms; Markov intertwining; predictive completion.

Source results

- **Lossless Reconstruction** [R41]. Lossless reconstruction of the covered gas history.
- **Transport of record observables and established estimates** [R42]. Transports record observables and established estimates.
- **Measure and observable isomorphism for invariant coordinates** [R43]. Measure and observable isomorphism for invariant coordinates.
- **Exact transition and history isomorphism for the recorded algorithm** [R44]. Exact transition and history isomorphism for the recorded algorithm.

Three spatial coordinates and the recorded clock

For this program, each walker has $x_i(n), v_i(n) \in \mathbb{R}^3$, and the observation clock is $t_n = n\Delta t$. The spacetime episode coordinate is $(t_n, x_i(n))$. CST temporal edges increase the integer iteration index; IG and IA data encode the recorded interactions. The causal-set theorem proves irreflexivity, transitivity, and local finiteness from that recorded ordering. Between two finite integer layers, the interval contains only finitely many episodes.

The scutoid chapter builds slabs between successive frames using their spatial cells and neighbor changes. Its temporal direction comes from the recorded clock. Proper time is defined by the reconstructed Lorentzian metric on this ordered geometry; its relation to the recorded clock is kept in the trajectory-clock comparison. The continuum geometry and local-operator estimates are applied on this specified construction.

The causal-set chapter makes the analytic dependency order explicit: fitness derivatives precede metric inversion; joint LSI precedes empirical gradient control; local bias and variance estimates precede operator consistency. Graph order and distance comparisons identify the recorded geometry to which those estimates apply. This preserves the $3 + 1$ origin of the construction throughout the prospectus.

Results supplying these constructions

- **The recorded Fractal Set is a causal set** [R45]. The recorded temporal order is a causal set.
- **Conditional compatibility of CST paths and geometric causality** [R46]. The order-to-geometric-causality comparison under its stated slab hypotheses.
- **Conditional volume matching and trajectory clocks** [R47]. Volume matching and trajectory clocks for the specified embedding.
- **Slab consistency and the scope of causal convergence** [R48]. Slab consistency in the moving-cell reconstruction.

Stage output. A field-readable history with the same native law. Gas moment, concentration, entropy, and approximation estimates can now be applied to the corresponding reconstructed observables.

11 Construct gauge and matter observables

Choose the internal representations used for the recorded color and companion data. The direct algebra contains Hermitian contractions, alternating determinant contractions, doublet coordinates, and composite triangle observables. The orbit theorem specifies precisely which internal frame equivalence those coordinates resolve.

Attach oriented transports to the declared links and form ordered products around interaction triangles and closed faces. The interaction holonomy detects the mismatch between attribution and interaction transports. Transported companion doublets retain both amplitude and angular matter energy.

Use gauge-invariant loop traces and transported matter contractions as the physical readouts. Distinguish their native values from auxiliary scalar proxies. Gauge covariance is implemented by transporting endpoint frames, so adjacent transformations cancel around a closed loop.

$$U_{\partial f} = \prod_{e \in \partial f}^{\rightarrow} U_e, \quad W_f = \frac{\text{Re tr } U_{\partial f}}{\dim R}, \quad U_{ij} \mapsto g_i U_{ij} g_j^{-1}.$$

Proof mechanism

Under an endpoint frame change, adjacent factors cancel in the ordered product around a face, leaving conjugation at the base point; the trace is invariant. The direct Gram and determinant coordinates retain the full declared color invariant algebra. For a companion doublet, compare its value with the transported neighboring doublet, separating amplitude and angular contributions. The interaction triangle retains the mismatch between IA and IG transports. This supplies the native observable whose curvature expansion and source response are analyzed downstream. The group representation, face incidence, and observable normalization are part of its definition.

Analytic inputs

Gram and determinant invariants; $\text{SU}(n)$ orbit reconstruction; Hermitian and alternating doublet contractions; projector triangle observables; oriented holonomy; angular matter energy.

Source results

- **Gram and determinant coordinates for $\text{SU}(n)$ orbits** [R49]. Gram and determinant coordinates for $\text{SU}(n)$ orbits.
- **$\text{SU}(2)$ from Hermitian and alternating doublet contractions** [R50]. $\text{SU}(2)$ frame structure from doublet contractions.
- **Interaction holonomy and angular matter energy of the recorded connection** [R51]. Interaction holonomy and angular matter energy of the recorded connection.
- **Gauge invariance of Wilson observables and actions** [R52]. Gauge invariance of Wilson observables and actions.

Recorded loops and geometric curvature

The Voronoi Wilson-loop chapter supplies a parallel construction directly from **RunHistory**: spatial Delaunay neighbor links on a frame, temporal adjacency between frames, recorded fitness or interaction phases, and minimal closed cycles. These are observable constructions on existing data.

The curvature chapter controls holonomy composition by telescoping the difference of ordered transport products. With bounded face complexity and the stated edgewise comparison, the accumulated error is $O(h^3)$. For a shape-controlled plaquette with area $A_\Pi \asymp h^2$, the small-loop theorem yields

$$\frac{(\mathcal{H}[\Pi] - I)V}{A_\Pi} \longrightarrow R(X, Y)V,$$

with an explicit $O(A_\Pi^{1/2})$ leading comparison error. This is the geometric transport tool used alongside the gauge face-action expansion; each connection retains its declared representation and correspondence.

Results supplying these constructions

- **Error accumulation for comparison holonomy** [R53]. Telescoping error bound for comparison holonomy.
- **Curvature recovery from consistent plaquette transport** [R54]. Curvature recovery from consistent plaquette transport.
- **Quantitative curvature expansion of the recorded face action** [R55]. Quantitative curvature remainder for the recorded gauge face action.
- **Reference-transport covariance** [R56]. Gauge covariance and invariant norms for fixed measurement paths.

Stage output. The concrete invariant field algebra and its native interaction observables. This algebra is the descriptor on which the history likelihood induces the effective action.

12 Derive the native field action and response

Start from the likelihood of the complete stochastic history, including selection, masks, cloning writes, rotations, and kinetic noises. Push the history and reference laws forward through the field descriptor, and condition the likelihood on that descriptor. Its negative logarithm is the effective field action.

The generating functional is an expectation under that native field law. Its derivatives reproduce the recorded correlation functions. The effective predictive kernel and the completed descriptor algebra preserve the original history predictions; channel compression carries the exact memory of discarded coordinates.

Disintegrate the action into recorded geometry and normalized conditional gauge fibers. Retain the fiber denominator and geometry weight so their recombination recovers the native sourced law. Differentiate actual source-dependent updates to obtain native response currents and the weak connection-variation identity. The discrete Ward identity and Wilson action give the gauge equations and curvature functional.

$$\begin{aligned}
Y &= \mathcal{D}(\omega), \quad \lambda = \mathcal{D}_* R, \quad a(y) = \mathbb{E}_R \left[\frac{dP}{dR} \middle| Y = y \right], \\
S_{\text{alg}}(y) &= -\log a(y), \quad d\nu = e^{-S_{\text{alg}}} d\lambda, \\
\mathcal{Z}(J) &= \mathbb{E}_P \exp \left(i \sum_r J_r O_r(Y) \right).
\end{aligned}$$

Proof mechanism

For bounded F , conditional expectation gives

$$\int F(y) a(y) \lambda(dy) = \mathbb{E}_R[F(Y) dP/dR] = \mathbb{E}_P F(Y).$$

Thus the descriptor action reproduces the original measure and all its bounded correlations. Disintegrate ν over geometry; the conditional gauge denominator is retained, so recombining the fibers recovers the same native expectation. Introduce sources through the executed force or metric update and differentiate its likelihood on its stated domain. This produces the native response current and weak variation identity. Gauge variations of the discrete face functional give the graph Ward identity. Matching these identities to the limiting Yang–Mills law is an explicit downstream identification.

Analytic inputs

Conditional likelihood; descriptor action; predictive kernel; characteristic functional; geometry-fiber disintegration; source tangents; native response identities; graph Ward identity.

Source results

- **Action emergence from the recorded stochastic dynamics** [R57]. Action emergence from the recorded stochastic dynamics.
- **Effective gauge-channel action and predictive kernel of the recorded algorithm** [R58]. Effective gauge-channel action and predictive kernel.
- **Exact action on the recorded geometry fibers** [R59]. Exact action on recorded geometry fibers.
- **Exact source response of the algorithm-derived field action** [R60]. Exact source response of the algorithm-derived field action.

Reduced fields retain the derived memory

The field-equations and direct-observable chapters both construct the exact reduction of the complete state. Conditional expectation onto the field descriptor splits the transition into resolved and unresolved blocks. Eliminating the unresolved block gives the transient memory equation. Conditioning on the complete observed field history gives the explicit filtering equation and its prediction covariance.

The prediction-complete descriptor is formed by closing the original bounded observable algebra under the actual transition. Its finite conditional partitions give convergent transition matrices. Thus the retained gauge channels have an exact history action and prediction law, and their

simulation approximation can be refined without postulating a closed differential equation for a few initial readouts.

The expansion and macroscopic-closure chapter supplies the complementary closure calculus. Information closure is equality of the conditional macro-future laws given the microscopic and macroscopic pasts. Its theorem relates this condition to predictive causal-state maps. For generator-based coarse variables, its approximate Markov-closure proposition gives

$$\sup_x |L_N(\varphi \circ f)(x) - \bar{L}\varphi(f(x))| \leq \epsilon_N \implies |\mathbb{E}[\text{Dynkin residual over } [0, t]]| \leq t\epsilon_N.$$

This is a quantitative tool for comparing a proposed macroscopic evolution with the complete recorded dynamics. The finite-step construction above continues to use its actual discrete kernel.

The algorithmic-thermodynamics chapter derives stationary response from the discrete Poisson equation for the actual kernel. With the stated weighted mixing and differentiable-kernel hypotheses, put

$$u_A = \sum_{n \geq 0} K^n [A - \pi(A)], \quad (I - K)u_A = A - \pi(A).$$

If D is the derivative of the configured kernel, the response is

$$\left. \frac{d}{d\theta} \pi_\theta(A) \right|_0 = \pi(Du_A).$$

This sums the delayed effect of the perturbation through the same dynamics. The path-KL identities likewise retain the specified forward and reverse experiments and their record pushforwards. They complement the native field-source likelihood calculations.

Results supplying these constructions

- **Transient projected field equation** [R61]. Exact reduced-field propagation with eliminated-state memory.
- **Field evolution conditional on its observed history** [R62]. Conditional-history evolution and its primitive-computed noise covariance.
- **Prediction-complete gauge descriptors from the actual transition** [R63]. Prediction-complete descriptors generated by the actual transition.
- **Finite transition matrices converging to the completed channel theory** [R64]. Finite transition matrices approximating that completed field theory.
- **Closure implications and their converse** [R65]. Predictive closure relations for deterministic coarse observables.
- **Exact and approximate Markov closure** [R66]. Exact block closure and a quantified approximate-generator residual.
- **Stationary response from a discrete Poisson equation** [R67]. Exact stationary response from the discrete Poisson equation.
- **Exact path KL identities** [R68]. Path likelihood and KL identities for the specified dynamics and reverse experiment.

Stage output. A native field measure, generating functional, and source-response calculus whose correlations remain those of the complete gas history. These are the objects carried into the continuum hierarchy.

13 Construct the recorded quantum matter algebra

Build the exterior algebra of recorded modes. Alternating insertions encode fermionic words with the specified graded signs and preserve independent exterior combinations as operators. The construction retains the meaning of its recorded contractions and replica correlations.

Transport the actual contraction into the CAR algebra. The recorded channel construction supplies a unital completely positive evolution, a two-point correspondence, and a higher-word prescription. Differentiate the transported evolution to identify its native generator.

For local matter observables, use regional covariance and locality-defect coefficients derived from the recorded evolution. The local-net application is made in the declared even observable sector. Together with the gauge readouts, this gives the finite operator formulation used by the connected algorithmic field theory.

$$\{a(f), a(g)\} = 0, \quad \{a(f), a^*(g)\} = \langle f, g \rangle I, \quad \mathcal{C}_t : \text{CAR}(\mathcal{K}) \rightarrow \text{CAR}(\mathcal{K}).$$

Proof mechanism

Lift the recorded one-particle mode space to its exterior algebra and represent alternating insertions by creation and annihilation operators. The insertion norm proves faithfulness. Transport the recorded contraction through the prescribed isometric dilation and compression to obtain a unital completely positive CAR channel. Its stated two-point and higher-word prescriptions retain the recorded correlations. The native generator follows by differentiating the transported evolution on its declared domain. Regional covariance coefficients quantify the locality comparison used by the CAR local-net application. This stage fixes the operator algebra whose physical spacetime realization remains part of the final reconstruction task.

Analytic inputs

Exterior Hilbert construction; faithful insertions; CAR relations; replica contractions; isometric compression and complete positivity; native generator; regional covariance.

Source results

- **Faithful exterior algebra of alternating recorded insertions** [R69]. Faithful oriented-word algebra for recorded modes.
- **Completely positive CAR evolution of the recorded contraction** [R70]. Unital completely positive CAR evolution from the recorded contraction.
- **Separate local-net application to the recorded CAR representation** [R71]. Separate local-net application to the recorded CAR representation.

Native regional dynamics are already computed

The lattice-QFT chapter derives localized fermion evolution and the regional covariance of the native readouts from the complete recorded transition. Its locality-defect theorem supplies coefficients for the declared regional modes. The inherited-history covariance theorem transports the full original temporal correlations, including overlaps and shared innovations.

These calculations support the recorded CAR local-net application and specify its even observable sector.

Results supplying these constructions

- **Native fermion evolution, localization, and record covariance** [R72]. Native fermion evolution, localization, and record covariance.
- **Exact inherited-history covariance of native regional readouts** [R73]. Exact inherited-history covariance of native regional readouts.
- **Existing LSI and native contraction bounds** [R74]. Transport of existing LSI and native contraction estimates into the recorded Fock setting.

Stage output. A precise matter operator algebra with the recorded dynamics and word correlations. Its continuum identification proceeds alongside the native gauge correlation hierarchy.

14 Retain fluctuations and a common correlation hierarchy

Center empirical fields about their selected population expectation and use the $\sqrt{N}$ fluctuation scaling. The joint LSI bounds their moments. Distribution-valued compactness places the fluctuation correlations on a common subsequence in the declared test-function topology.

Compute drift and covariance from the complete selected update. In particular, retain covariance from shared collision rotations, cloning writes, measurement marks, and kinetic innovations. These equations determine the fluctuation dynamics around the mean-field background.

The native gauge-hierarchy theorems carry finite words and reflected products of normalized geometric readouts and the full color invariant algebra on a common subsequence. Uniform-integrability and mark-reconstruction estimates control passage of expectations. Source-action and geometry-fiber continuum results retain the sourced history interpretation on this same construction.

$$\Xi_N(\varphi) = \sqrt{N}(L_N\varphi - \mathbb{E}[L_N\varphi]), \quad \mathbb{E}|\Xi_N(\varphi)|^2 \leq C_* \text{Lip}(\varphi)^2.$$

Proof mechanism

Apply the joint-law LSI concentration bound to the centered Lipschitz empirical fields. Higher moments follow from the same law-specific bounds. The chapter’s test-space construction gives tightness and a common subsequence for the distributional hierarchy. For dynamics, use the generator product rule and conditional variance decomposition to compute drift and noise from the complete update. For gauge words, combine bounded normalized readouts with the explicit uniform-integrability and mark-reconstruction estimates. A diagonal extraction over the stated observable/test classes places the words and reflected products in one native hierarchy, preserving their common sourced-law interpretation.

Analytic inputs

Joint-LSI moment bounds; test-space compactness; all-order moment passage; full-update conditional variance; common subsequences; uniform integrability; reconstructed reflected-matrix control.

Source results

- **Distribution-valued compactness from the established stationary LSI** [R75]. Distribution-valued compactness from the established stationary LSI.
- **Drift and covariance equations for the complete selected update** [R76]. Drift and covariance equations for the complete selected update.
- **Native physical hierarchy of the full recorded color invariants** [R77]. Common native physical hierarchy of full recorded color invariants.
- **Geometry-fiber action in the common native continuum limit** [R78]. Geometry-fiber action in the common native continuum limit.

The fluctuation field retains algorithmic time

For the stationary discrete chain, the fluctuation theorem uses the piecewise-constant record at the fixed observation spacing. Its phase-space variable is $z = (x, v) \in \mathbb{R}^6$. Physical spatial tests are evaluated through the same recorded positions and may be chosen independent of v ; their temporal variable remains the algorithmic observation clock.

The native gauge-hierarchy theorem retains the full Gram coordinates, complex determinants, triangle products, masks, sources, and configured weights. It constructs their finite-word and reflected-product limits on a common subsequence. The direct color observable theorem also derives nontrivial doublet fluctuations from the actual companion draw, while the paired momentum result gives a quantitative nonzero fluctuation estimate for its identified stationary model. These supply concrete finite-sector and fluctuation information within their respective law families.

Results supplying these constructions

- **Nontrivial doublet fluctuations from the actual companion draw** [R79]. Nontrivial doublet fluctuations from the actual sampled companion.
- **Nonzero momentum fluctuations in the established Euclidean paired model** [R80]. A nonzero momentum fluctuation bound for the identified paired stationary model.
- **Uniform integrability and reconstruction bounds for the full direct color algebra** [R81]. Explicit full-color word and mark-reconstruction bounds.

Stage output. A distributional fluctuation theory and a common native gauge correlation hierarchy. The next stage controls the spatial reconstruction scales and differential operators used to interpret that hierarchy.

15 Control the continuum operator with derived tools

Identify the reconstructed product geometry and its sampling measure $q \, d\text{vol}_g$. Smoothed coefficient and spectral bounds supply geometric regularity; the geometry lemma gives a sufficient global-hyperbolicity condition. Normalize sampling using the selected history law and density weights.

Verify the kernel moments in local normal coordinates. The Taylor estimate gives bias at most $C_b \varepsilon^2$. The covariance estimate controls the density-corrected local summands and yields variance at most $C_{\text{mix}} C_v / (M \varepsilon^{D+2})$.

Choose $\varepsilon_M = \ell_0 M^{-\beta}$, with $0 < \beta < 1/(D+4)$, as in the chapter. Both squared bias and variance vanish. The balanced exponent $\beta = 1/(D+6)$ gives the displayed mean-square rate. Compare the reconstructed episode operator with this estimator, controlling metric, density, neighborhood, and kernel errors after normalization.

$$\mathbb{E} \left| \widehat{L}_{M,\varepsilon} f(p) - \square_g f(p) \right|^2 \leq C_b^2 \varepsilon^4 + \frac{C_{\text{mix}} C_v}{M \varepsilon^{D+2}},$$

$$\varepsilon_M = \ell_0 M^{-\beta}, \quad 0 < \beta < \frac{1}{D+4}, \quad \beta = \frac{1}{D+6} \implies \text{MSE} = O(M^{-4/(D+6)}).$$

Proof mechanism

In normal coordinates Ψ_p , expand $f \circ \Psi_p$ to fourth order and the volume Jacobian to third order. Verified kernel moments remove the first-order and unwanted odd terms and select the intended second-order signature. The remainder is $O(\varepsilon^2)$ after normalization. For samples with density q , divide each summand by q ; its second moment scales as ε^{-D-2} . Summing the chapter's covariance bound gives the displayed variance. Balance bias and variance through the bandwidth schedule. Finally prove the recorded operator comparison at this normalization and a joint weighted quadrature estimate for action evaluation on the interacting records.

Analytic inputs

Smooth metric coefficients; geometric identification; importance weights; normal-coordinate kernel moments; Taylor bias; shrinking-neighborhood covariance; bandwidth schedule; normalized reconstruction comparison.

Source results

- **Direct local bias estimate in specified normal coordinates** [R82]. Direct local bias estimate in specified normal coordinates.
- **A covariance condition sufficient for A4** [R83]. Covariance condition sufficient for the estimator variance bound.
- **A sufficient bandwidth schedule for A6** [R84]. Explicit sufficient bandwidth schedule.
- **Conditional continuum consistency** [R85]. Transfers estimator consistency to the reconstructed operator and action under its listed inputs.

Density correction is part of the existing construction

The continuum and emergent-geometry chapters distinguish the walker sampling law from the geometric volume measure. If the episode marginal is $\pi(dy) = q(y) d\text{vol}_g(y)$, then

$$\mathbb{E} \left[\frac{1}{M} \sum_{i=1}^M \frac{F(Y_i)}{q(Y_i)} \right] = \int F d\text{vol}_g.$$

This expectation identity uses the actual marginal and requires no independence. The chapter also gives the explicit QSD/Doob distinction and the normalized time-window density. It therefore computes the importance weight from the selected sampling experiment. Walker normalization, geometric integration, and centered field fluctuations are separate quantities with explicitly related formulas.

A second continuum estimate follows directly from joint LSI

For the smooth normalized summand $H_{\varepsilon,p}$, the causal-set chapter differentiates the actual empirical estimator:

$$\nabla_i A_{M,\varepsilon} = M^{-1} \nabla H_{\varepsilon,p}(Y_i).$$

The established joint Poincaré inequality gives

$$\begin{aligned} \text{Var}(A_{M,\varepsilon}) &\leq \frac{C_*}{M} \int |\nabla H_{\varepsilon,p}|^2 d\pi \leq \frac{C_* C_\nabla}{M \varepsilon^{D+4}}, \\ \mathbb{E}|A_{M,\varepsilon} - \square_g f(p)|^2 &\leq C_b^2 \varepsilon^4 + \frac{C_* C_\nabla}{M \varepsilon^{D+4}}. \end{aligned}$$

Thus $\varepsilon = M^{-1/(D+8)}$ gives $O(M^{-4/(D+8)})$ for that observation law and Sobolev summand. In $D = 4$, this is $O(M^{-1/3})$. The covariance route above gives $O(M^{-2/5})$ in $D = 4$ under its covariance hypotheses. These are two proved sufficient routes. Single-time observations use their joint law; pooled temporal episodes use the actual episode law or the chapter's temporal covariance estimate.

The spatial lattice-QFT chapter supplies the corresponding density-corrected kernel limit and sampling control for slice operators. Operator reconstruction and action quadrature are then composed with the same smooth-field and sampling estimates.

The computational-proxies chapter also specifies the geometric cell weights used by recorded observables:

$$V_i^g = V_i^{\text{Eucl}} \sqrt{\det g(x_i)}, \quad w_i^g = V_i^g / \sum_j V_j^g.$$

The coordinate identity $d\text{vol}_g = \sqrt{\det g} dx$ supplies their geometric meaning. These weights are used with the chosen cell approximation; inverse sampling-density weights supply the separate Monte Carlo integration identity above. The chapter's curvature, distance, and covariance probes provide observable diagnostics for the reconstructed geometry.

Results supplying these constructions

- **Normalized sampling and the distinction between QSD and invariance** [R86]. The actual normalized sampling law, Doob law, and inverse-density weights.
- **Normalized importance sampling and variance** [R87]. Riemannian importance integration and joint-LSI variance control.

- **Continuum estimator error from the established joint LSI** [R88]. Direct continuum estimator error from the already established joint LSI.
- **Density-corrected limit with sufficient sampling control** [R89]. Spatial density-corrected kernel limit with sufficient sampling control.
- **Volume Element Transformation** [R90]. Coordinate volume transformation underlying geometric cell weights.

Stage output. A controlled continuum differential operator. The bias, covariance, and reconstruction estimates also supply the local integration bounds needed for an action limit.

16 Pass actions, sources, and correlations through the limits

Use geometric quadrature and the recorded face-action curvature expansion to obtain Wilson action consistency. Keep face orientation, matrix-valued curvature, face weights, and coupling normalization. The classical continuum term is the quadratic Yang–Mills curvature energy.

Combine local operator convergence with weighted quadrature to pass scalar actions. For actions evaluated on the same interacting records as the operator, carry the joint weighted error estimate. The native source-action and geometry-fiber results carry the selected likelihood and its sourced correlations into the common continuum hierarchy.

Assemble the quantitative errors: particle approximation, relaxation, survival conditioning, temporal consistency, local bias, correlated sampling, and record reconstruction. The time-consistency theorem propagates a local kernel defect using stability. QSD perturbation uses conditioned-block comparison with survival normalization. State a joint scale schedule that makes these errors vanish while retaining the required constants.

$$S_W \longrightarrow \frac{1}{4g^2} \int_{\mathbb{R}_t \times \mathbb{R}_x^3} F_{\mu\nu}^a F_{\mu\nu}^a dt d^3x,$$

$$\mathbb{E}|\hat{\varphi}_N - \rho\varphi| \leq b_N + \sqrt{v_{N,h}} + d_h + m_{N,h,n}.$$

Proof mechanism

Expand the actual small-face transport and control the remainder by the curvature and geometric bounds. Weighted face quadrature yields the displayed classical action under the chapter’s hypotheses. For temporal error, use the telescoping identity

$$P_h^n - T_h^n = \sum_{j=0}^{n-1} P_h^j (P_h - T_h) T_h^{n-1-j}.$$

A local defect of order h^{p+1} and stability give a finite-time defect of order h^p . Stationary perturbation sums the defect against the relaxation resolvent; the QSD version compares normalized surviving blocks. Combine the resulting bias, sampling, stationary, and mixing terms using the total-observable-error theorem. Uniform integrability and the native source/fiber identities retain expectations along the common hierarchy.

Analytic inputs

Curvature remainder; geometric quadrature; same-sample action error; BAOAB weak consistency; split-system commutator terms; invariant-law perturbation; conditioned-block QSD stability; total observable error.

Source results

- **Wilson action consistency under geometric quadrature** [R91]. Wilson action consistency under geometric quadrature.
- **Common continuum limit of the native source likelihood and descriptor action** [R92]. Common continuum limit of the native source likelihood and descriptor action.
- **Time consistency for the specified transition family** [R93]. Time consistency for the specified transition family.
- **Total observable error** [R94]. Total observable error: population bias, sampling variance, time defect, and mixing error.

Action response and geometry fibers retain the source law

The native response-current theorem writes the source derivative as a distribution tested on the supplied recorded spacetime coordinates. The weak connection-variation identity compares that actual response with the Wilson/matter variation. The common source-action limit keeps the same likelihood and descriptor refinement throughout the passage. Geometry-fiber disintegration retains its conditional denominator, so recombining the fiber and geometry contributions recovers the original sourced correlations.

In the present $3 + 1$ roadmap the field integration domain is $\mathbb{R}_t \times \mathbb{R}_x^3$, with t supplied by the calibrated recorded clock. The Yang–Mills quadratic contraction is taken with the metric convention of the Euclidean or Lorentzian reconstruction under discussion. This specifies the geometry used by the small-face and quadrature statements.

Results supplying these constructions

- **Native response current in recorded spacetime** [R95]. Native source-response current on the recorded spacetime.
- **Native response and the weak connection-variation identity** [R96]. Native response and the weak connection-variation identity.
- **Exact graph Ward identity** [R97]. Exact graph Ward identity.

Stage output. A continuum action and correlation hierarchy with an explicit error budget. The field-law, physical symmetry, and Hamiltonian identifications are completed on this limiting object.

17 Reconstruct fields from the equilibrium history

At equilibrium, retain the complete gauge and matter descriptor hierarchy generated by the three-dimensional gas and its recorded time. The exact field-law and transition isomorphisms

give its multi-time expectations. The fluctuation and native-hierarchy theorems supply moment bounds, test-space control, and common-subsequence correlation limits.

The Euclidean-regularity theorem carries actual moment and reconstruction bounds to tempered correlations. Commuting bosonic insertions give their permutation symmetry, and the exterior/CAR construction supplies the graded rule for the recorded matter operators. The local-net results establish isotony, the stated locality identities, and covariance under the identified same-law implementers.

The final physical representation is built from this equilibrium temporal hierarchy. Its reflection is in the algorithmic observation clock, and its spatial fields remain on $\mathbb{R}^3$. The source’s positive-transfer and quantum-reconstruction statements specify the representation hypotheses used to construct the physical Hilbert space, vacuum, and self-adjoint Hamiltonian. Those hypotheses remain attached to the corresponding field-axiom conclusions.

$$(F, G)_{\text{OS}} = \mathbb{E}[\overline{\Theta F} G], \quad (F, F)_{\text{OS}} \geq 0 \quad (F \in \mathcal{A}_+).$$

Proof mechanism

First transport the actual stationary history through the lossless descriptor map. The source’s operator-product formula computes each time-ordered correlation using the original kernel. Apply the existing test-space, moment, and uniform-integrability bounds to the same retained observable algebra. For temporal reconstruction, form the future algebra on $t \geq 0$ and its clock-reflected pairing. The source’s positive-transfer criterion, when identified for this hierarchy, gives the required positive semigroup representation. Quotient null vectors, complete the resulting Hilbert space, and identify physical time translation and its generator. Apply the local-net and field-axiom statements with their covariance, locality, spectrum, and vacuum hypotheses. This composes the established native construction with the physical representation step on its actual equilibrium law.

Analytic inputs

Tempered correlation bounds; bosonic/graded symmetry; native reflected matrices; physical translation identities; local algebras; vacuum reconstruction; transfer Hamiltonian and physical clock identification.

Source results

- **Euclidean regularity from actual moment bounds** [R98]. Euclidean regularity from actual moment bounds.
- **Symmetry of bosonic correlations** [R99]. Symmetry of bosonic correlations.
- **Physical reflection on the full native color algebra** [R100]. Physical reflection calculation on the full native color algebra.
- **Algorithmic QFT from recorded evolution and native bounds** [R101]. Connected algorithmic field theory from recorded evolution and native bounds.

The equilibrium correlator route is already specified

The twistor and calibration chapters define frame channel observables from the actual recorded state, including donor memory and validity data. For an invariant law of that same kernel K , let $\tilde{f}_a = f_a - \pi f_a$. The exact algorithmic-lag correlator is

$$C_{ab}(\ell) = \langle \tilde{f}_a, K^\ell \tilde{f}_b \rangle_{L^2(\pi)}.$$

The complete record transition fixes this correlation. The time separation is $\ell\Delta t$. When the identified transfer representation is positive and self-adjoint with $K = e^{-\Delta t H}$, the book derives

$$C_{aa}(\ell) = \sum_{E_j > 0} |\langle j | \hat{f}_a | 0 \rangle|^2 e^{-E_j \ell \Delta t}.$$

Its leading supported energy is read from the large-lag decay. The local twistor operators are observable channels feeding this pipeline; the mass is a spectral property of their correlator.

Which reflection and symmetry enter this program

The chosen temporal reflection acts on the stationary history clock:

$$\Theta : (t, x) \mapsto (-t, x), \quad x \in \mathbb{R}^3.$$

The reflection and transfer identities are applied to the derived temporal observable law. The book's separate tests of a translated embedding coordinate have their stated coordinate action; they are not imported into this algorithmic-clock program without the corresponding identification. Likewise, symmetry is evaluated through the physical observable pushforward and its temporal implementers, while the complete anchored record remains available for decoding and prediction.

The existing source statements give regularity, bosonic or graded symmetry, native local multiplication identities, the CAR local-net construction, and exact stationary temporal correlations. The positive-transfer and physical field-axiom statements retain their explicit representation and covariance hypotheses. In the proof program these hypotheses are applied on the same derived equilibrium hierarchy and clock. This identifies the object on which the final physical reconstruction is carried out.

Results supplying these constructions

- **Algorithmic Evolution of Frame-Averaged Twistor Channels** [R102]. Exact algorithmic evolution and stationary frame-correlator identity.
- **Positive Transfer Representation of a Frame Correlator** [R103]. Spectral expansion for the same channel under its stated positive-transfer hypothesis.
- **Exact HK locality of the native multiplication representation** [R104]. Exact locality identity in the native multiplication representation.

Stage output. The same equilibrium observable hierarchy, recorded-time correlators, native local algebras, and the stated physical representation criteria. These identify the field law and transfer family on which the gap-survival theorem is applied.

18 Preserve the physical Yang-Mills mass gap

Carry the functional-inequality and evolution constants along the selected cutoff and volume family. Poincaré supplies coercivity of its associated form; hypocoercivity supplies relaxation for the kinetic evolution. Complete the identification that connects these estimates to the reconstructed physical gauge transfer Hamiltonian.

For that self-adjoint Hamiltonian family H_a , prove a common positive gap λ_* above the normalized vacuum. Construct isometric embeddings J_a into the limiting Hilbert space, show convergence of the vacuum vectors, and establish strong convergence of the embedded semigroups.

The gap-survival theorem then passes the vacuum and spectral bound to H . Restriction to an invariant gauge sector preserves the gap under the chapter’s operator hypotheses. Fix the physical action and time units so the limiting energy scale remains positive. The endpoint theorem combines the physical reconstruction of Stage 17 with this gap passage.

$$\begin{aligned} \|e^{-tH_a}(I - P_{\Omega_a})\| &\leq e^{-\lambda_* t}, \quad \lambda_* > 0, \\ J_a e^{-tH_a} J_a^* &\xrightarrow{\text{strong}} e^{-tH}, \quad \text{spec}(H) \subset \{0\} \cup [\lambda_*, \infty). \end{aligned}$$

Proof mechanism

Let $J_a : \mathcal{H}_a \rightarrow \mathcal{H}$ be isometries with $J_a J_a^* \rightarrow I$ strongly and $J_a \Omega_a \rightarrow \Omega$. For $f \in \mathcal{H}$, put $f_a = J_a^* f$. The embedded gap bound gives

$$\|J_a e^{-tH_a}(f_a - P_{\Omega_a} f_a)\| \leq e^{-\lambda_* t} \|f_a\|.$$

Pass to the strong semigroup limit and the vacuum limit. The resulting estimate holds on $\Omega^\perp$; the spectral theorem excludes spectrum in $(0, \lambda_*)$ and any further zero-energy vector. Multiplication by a fixed $\hbar_{\text{eff}}$ converts the bound to physical energy. The program must identify this self-adjoint family with the gauge transfer Hamiltonians and retain the positive bound through cutoff removal and physical volume growth.

Analytic inputs

Same-law coercivity; physical transfer identification; self-adjoint semigroup estimate; compatible Hilbert-space embeddings; vacuum convergence; strong semigroup convergence; invariant-sector restriction.

Source results

- **Survival of a uniformly controlled self-adjoint gap** [R105]. Survival of a uniformly controlled self-adjoint gap.
- **Restriction of a gap to an invariant gauge sector** [R106]. Restriction of a gap to an invariant gauge sector.

Stage output. Completion target: a nontrivial four-dimensional Yang-Mills quantum field theory with a unique vacuum and a strictly positive physical mass gap. Its construction retains the original history law through the recorded and continuum identifications.

19 How the proved walker dynamics supply the field theory

This section collects the identifications used throughout the chapters into one dependency chain. Its starting point is the complete history of the three-dimensional gas at the recorded times $t_n = n\Delta t$.

19.1 Probability law, invariant descriptors, and frame observables

The empirical walker law is

$$L_N = \frac{1}{N} \sum_{i=1}^N \delta_{(x_i, v_i, a_i)}.$$

The direct field law instead retains the full specified observable descriptor $\mathcal{O}$ of the complete history:

$$\mu_{\text{dir}} = \mathcal{O}_* \mathbb{P}_{\text{rec}}, \quad \langle f \rangle_{\text{dir}} = \mathbb{E}_{\mathbb{P}_{\text{rec}}} f(\mathcal{O}(\mathcal{F})).$$

The direct three-color definition sets $d = 3$ and constructs its masked vector from the actual viscous force and velocity:

$$\tilde{c}_i^a = F_i^{\text{visc}, a} e^{i\kappa v_i^a}, \quad c_i = m_i \frac{\tilde{c}_i}{\max(\|\tilde{c}_i\|, \delta_c)}.$$

The Gram and complex determinant coordinates identify the common internal-frame orbit and preserve its invariant history correlations. Full coordinates and masks are retained before forming the particular frame averages. A full-rank internal anchor chart is one explicit inverse chart; the full orbit theorem also covers the rank-deficient strata.

At a recorded frame, the book distinguishes the valid-weight average and its fixed-normalization companion:

$$\mathcal{A}_t(O) = \frac{\sum_I w_I m_I O_I}{W_t}, \quad \mathcal{A}_t^N(O) = \frac{1}{N} \sum_I w_I m_I O_I = \frac{W_t}{N} \mathcal{A}_t(O), \quad W_t = \sum_I w_I m_I,$$

with the recorded zero-denominator convention. Each is an observable of the same full history and has its own specified correlator. The physical meaning is read from the chosen descriptor and its temporal law.

19.2 The history correlations retain the algorithmic clock

For the selected invariant law π , the lossless record theorem supplies

$$\hat{P}_h U = U P_h, \quad \hat{L} = U L U^{-1}$$

on the stated domain. The finite-step history formula and frame-correlator identity then give all retained correlations under the actual cloning and kinetic update. Cloning remains part of the evolution. Projection to a few channels uses the derived memory or the prediction-complete descriptor. Burn-in changes the initial law toward equilibrium; field time continues to be the calibrated recorded observation time.

19.3 Geometric integrals and fluctuations have derived normalizations

For the selected sampling law $q \, d\text{vol}_g$, the inverse-density formula converts normalized samples into geometric integrals. The normalized C^n calculus controls their coefficient fields. Joint LSI gives the direct empirical-gradient variance estimate, and the covariance route gives the alternative temporal/sampling estimate.

The centered hierarchy uses $\sqrt{N}$ scaling around the actual stationary expectation. Its distribution-valued compactness and moment bounds retain the collective fluctuation information. The native gauge hierarchy separately retains the complex color algebra and its sourced history law. These constructions supply the background, fluctuations, and gauge correlations as distinct derived objects.

19.4 Native likelihood and conditional geometry

Condition the complete path likelihood on the descriptor and take its negative logarithm. This yields the field action, predictive kernel, and generating functional. The native source identities differentiate the executed update, and geometry-fiber disintegration preserves the conditional denominator. Recombination therefore returns the same sourced correlations. The continuum comparison uses the derived spatial geometry and the recorded time slices.

19.5 Equilibrium representation and the gap

The source specifies quantum reconstruction through the equilibrium temporal hierarchy and its stated physical representation hypotheses. The resulting transfer Hamiltonian uses that clock and observable law. The mass-gap passage then applies the proved self-adjoint gap-survival theorem with its uniform bound, embeddings, vacuum limit, and strong semigroup convergence.

Completion criterion 1 (End-to-end composition). *For the configured $3 + 1$ recorded family, apply the existing moment, population, regularity, sampling, and native-hierarchy results under their stated hypotheses. Identify its limiting equilibrium hierarchy with the physical Yang–Mills representation using the source’s field-axiom and positive-transfer hypotheses. If the resulting Hamiltonians satisfy the uniform gap and convergence hypotheses of the gap-survival theorem, then*

$$\text{spec}(H) \subset \{0\} \cup [\lambda_*, \infty), \quad \ker H = \mathbb{C}\Omega, \quad \lambda_* > 0.$$

The calibrated energy gap is at least $\hbar_{\text{eff}}\lambda_$.*

The criterion composes the source results. Representation hypotheses remain in their statements, while the population dynamics, normalizations, field equations, and inherited correlations are the constructions already derived in the chapters.

20 The estimates that control the continuum

Tool	Controlled quantity	Constant or identification retained
Moment and tail drift	Escape, tightness, uniform integrability	Moment order, drift coefficient, coverage defect
Poincaré inequality	Empirical variance and form coercivity	Joint-law C_* , appropriate Dirichlet form
Joint LSI	Entropy, concentration, fluctuation moments	Full gradient, status term, law identification
Hypocoercivity	Kinetic relaxation	Modified norm, prefactor, positive rate
Population comparison	Particle and time approximation	Initial moments, bias, horizon, attraction branch
Normalized C^n calculus	Smooth reconstructed coefficients	Kernel widths, floors, denominator lower bound
Local kernel calculus	Differential-operator bias	Geometry, moments, normal-coordinate bounds
Covariance estimate	Shrinking-neighborhood variance	$C_{\text{mix}}/(M\varepsilon^{D+2})$
Direct joint-LSI estimate	Continuum error for dependent smooth observations	$C_*C_{\nabla}/(M\varepsilon^{D+4})$
Sampling density correction	Geometric integration from probability samples	Actual q , weight q^{-1} , selected law
Recorded operator comparison	Actual graph reconstruction	Error after ε^{-D-2} normalization
Temporal and ensemble perturbation	Weak, stationary, conditioned errors	Stability, resolvent rate, survival weights
Native likelihood and source identities	Gauge action and correlations	Descriptor support, geometry-fiber normalization
Physical reflection and symmetry	Quantum Hilbert space and dynamics	Selected spacetime hierarchy and physical clock
Strong semigroup convergence	Survival of a positive gap	J_a , Ω_a , cutoff/volume-uniform λ_*

Joint scale choice. Fix the transition parameters for each declared family, choose its observation regime using the population/time estimates, and set a bandwidth for which the local bias and correlated variance vanish. Then control reconstruction and action quadrature at that normalization. If h , algorithmic locality, confinement, or noise changes, retain the corresponding uniform estimates for that changed transition family. Finally pass the sourced hierarchy and physical transfer operators through the same cutoff and volume sequence.

A Source theorem register

The references below point to the full statements and proofs in the repository snapshot. File paths are relative to `docs/source/2_fractal_gas/`. Line numbers identify the start of the named directive. The labels provide stable search keys when line numbers change.

[R1] Component momentum, energy, and shared covariance

`thm-mean-field-component-identities`
`convergence_program/08_mean_field.md`, line 253.

[R2] One-step collision consistency

`thm-mean-field-one-step-consistency`
`convergence_program/08_mean_field.md`, line 278.

[R3] Foster-Lyapunov drift for the composed kernel

`thm-foster-lyapunov-main`
`convergence_program/06_convergence.md`, line 451.

[R4] Fully evaluated selection-driven moment drift without a prescribed restoring trap

`thm-slceg-full-moment`
`convergence_program/06a_structural_landscape_convergence.md`, line 15018.

[R5] Entropy contraction to an explicit confinement floor

`thm-slce-entropy-floor`
`convergence_program/06a_structural_landscape_convergence.md`, line 15296.

[R6] Structural/Barycenter Split for Full W_2

`thm-full-w2-split`
`convergence_program/04_wasserstein_contraction.md`, line 560.

[R7] Joint convergence of mass and shape

`thm-hk-convergence-main-assembly`
`convergence_program/11_hk_convergence.md`, line 373.

[R8] Finite-horizon survival from safe-walker estimates

`thm-exponential-survival`
`convergence_program/11_hk_convergence.md`, line 245.

[R9] A weighted coupling form of Harris’ argument

`thm-convergence-conservative-harris`
`convergence_program/06_convergence.md`, line 649.

[R10] QSD convergence from a two-sided surviving-block bound

`thm-main-convergence`
`convergence_program/06_convergence.md`, line 801.

[R11] Relaxation to the selected law of a recorded gauge window

`thm-ym-qsd-window-relaxation`
`2_fractal_set/05_yang_mills_noether.md`, line 6228.

[R12] The fixed-step mean-field equation

`thm-mean-field-equation`
`convergence_program/08_mean_field.md`, line 395.

[R13] Exact mass and weak field balances

`thm-mass-conservation`
`convergence_program/08_mean_field.md`, line 422.

[R14] Explicit finite-step population law

thm-algorithmic-explicit-transition
 3_fitness_manifold/04_field_equations.md, line 230.

[R15] Exact field characteristics and moment hierarchy

thm-algorithmic-field-characteristics
 3_fitness_manifold/04_field_equations.md, line 282.

[R16] Finite-step observable equation and fluctuation covariance

thm-algorithmic-observable-increment
 3_fitness_manifold/04_field_equations.md, line 657.

[R17] Exact discrete spatial density and momentum equations

thm-algorithmic-spatial-field-equations
 3_fitness_manifold/04_field_equations.md, line 1899.

[R18] Coupled population, mechanical, and metric fields

thm-algorithmic-coupled-field-system
 3_fitness_manifold/04_field_equations.md, line 2106.

[R19] Finite-horizon propagation of chaos for the actual update

thm-chaos-finite-time-consistency
 convergence_program/09_propagation_chaos.md, line 1190.

[R20] Closed full-law convergence with unbounded raw reward

thm-slcw-active-contraction
 convergence_program/06a_structural_landscape_convergence.md, line 14371.

[R21] Uniform-time trajectories, stationary chaos and commuting limits

cor-slcw-trajectories
 convergence_program/06a_structural_landscape_convergence.md, line 14710.

[R22] Uniform trajectory approximation through a diverging horizon

thm-slcgt-growing-trajectory
 convergence_program/06a_structural_landscape_convergence.md, line 18248.

[R23] Exchangeability of a unique QSD

thm-qsd-exchangeability
 convergence_program/12_qsd_exchangeability_theory.md, line 37.

[R24] Deterministic empirical limits imply chaos

thm-propagation-chaos-qsd
 convergence_program/12_qsd_exchangeability_theory.md, line 181.

[R25] Exact covariance identity for finite populations

thm-correlation-decay
 convergence_program/12_qsd_exchangeability_theory.md, line 212.

[R26] LSI from a quadratic Lyapunov bound

thm-unconditional-lsi
 convergence_program/15_kl_convergence.md, line 369.

[R27] LSI for the kinetic reference measure

thm-kinetic-lsi
 convergence_program/15_kl_convergence.md, line 513.

[R28] N -uniform LSI for specified joint laws

cor-n-uniform-lsi
 convergence_program/15_kl_convergence.md, line 544.

[R29] Poincaré inequality and empirical observables

cor-quantitative-lsi-final

convergence_program/15_kl_convergence.md, line 2149.

[R30] Kinetic hypocoercive entropy convergence

thm-villani-hypocoercivity

convergence_program/15_kl_convergence.md, line 805.

[R31] Entropy convergence for the full normalized swarm evolution

thm-kl-convergence-euclidean

convergence_program/15_kl_convergence.md, line 1164.

[R32] LSI for fixed marginals and their limits

cor-kl-lsi-mean-field-limit

convergence_program/15_kl_convergence.md, line 2177.

[R33] C^3 regularity of sampled, surrogate, and expected fitness

thm-c3-regularity

convergence_program/14_a_geometric_gas_c3_regularity.md, line 726.

[R34] Full sampled-fitness majorant

thm-main-cinf-regularity-fitness-potential-full

convergence_program/14_b_geometric_gas_cinf_regularity_full.md, line 963.

[R35] Differentiating normalized mean-field integrals

thm-cinf-mean-field-integrals

convergence_program/14_b_geometric_gas_cinf_regularity_full.md, line 1396.

[R36] Smooth convergence of normalized empirical fields

thm-cinf-empirical-field-convergence

convergence_program/14_b_geometric_gas_cinf_regularity_full.md, line 1426.

[R37] Metric represented by the implemented Hessian branch

prop-geometry-clipped-metric

3_fitness_manifold/01_emergent_geometry.md, line 122.

[R38] Uniform ellipticity from metric bounds

thm-uniform-ellipticity-latent

3_fitness_manifold/01_emergent_geometry.md, line 234.

[R39] Uniform ellipticity from the spectral margin

thm-gg-ueph-construction

convergence_program/17_geometric_gas.md, line 197.

[R40] N -uniform LSI for an identified geometric law

thm-gg-lsi-main

convergence_program/17_geometric_gas.md, line 720.

[R41] Lossless Reconstruction

thm-fractal-set-lossless

2_fractal_set/01_fractal_set.md, line 1638.

[R42] Transport of record observables and established estimates

prop-fractal-set-analytic-transfer

2_fractal_set/01_fractal_set.md, line 1955.

[R43] Measure and observable isomorphism for invariant coordinates

thm-sm-direct-measure-isomorphism

2_fractal_set/04_standard_model.md, line 1759.

[R44] Exact transition and history isomorphism for the recorded algorithm

thm-sm-instantiated-record-transition
 2_fractal_set/04_standard_model.md, line 2736.

[R45] The recorded Fractal Set is a causal set

thm-fractal-is-causal-set
 2_fractal_set/02_causal_set_theory.md, line 131.

[R46] Conditional compatibility of CST paths and geometric causality

prop-fractal-causal-order-equivalence
 2_fractal_set/02_causal_set_theory.md, line 237.

[R47] Conditional volume matching and trajectory clocks

thm-fractal-faithful-embedding
 2_fractal_set/02_causal_set_theory.md, line 616.

[R48] Slab consistency and the scope of causal convergence

prop-scutoid-cst-compatibility
 3_fitness_manifold/02_scutoid_spacetime.md, line 483.

[R49] Gram and determinant coordinates for $SU(n)$ orbits

thm-sm-direct-orbit-isomorphism
 2_fractal_set/04_standard_model.md, line 1542.

[R50] $SU(2)$ from Hermitian and alternating doublet contractions

thm-sm-direct-su2-invariants
 2_fractal_set/04_standard_model.md, line 1270.

[R51] Interaction holonomy and angular matter energy of the recorded connection

prop-ym-recorded-interaction-sector
 2_fractal_set/05_yang_mills_noether.md, line 605.

[R52] Gauge invariance of Wilson observables and actions

thm-wilson-action-gauge-invariance
 2_fractal_set/05_yang_mills_noether.md, line 1649.

[R53] Error accumulation for comparison holonomy

lem-regge-holonomy-approx
 3_fitness_manifold/03_curvature_gravity.md, line 955.

[R54] Curvature recovery from consistent plaquette transport

thm-riemann-scutoid
 3_fitness_manifold/03_curvature_gravity.md, line 991.

[R55] Quantitative curvature expansion of the recorded face action

lem-ym-recorded-face-action-remainder
 2_fractal_set/05_yang_mills_noether.md, line 507.

[R56] Reference-transport covariance

lem-reference-transport-covariance
 3_fitness_manifold/09_measurement.md, line 627.

[R57] Action emergence from the recorded stochastic dynamics

thm-ym-recorded-action-emergence
 2_fractal_set/05_yang_mills_noether.md, line 321.

[R58] Effective gauge-channel action and predictive kernel of the recorded algorithm

thm-sm-effective-recorded-gauge-dynamics
 2_fractal_set/04_standard_model.md, line 4020.

[R59] Exact action on the recorded geometry fibers

thm-ym-native-geometry-fiber-action

2_fractal_set/05_yang_mills_noether.md, line 3714.

[R60] Exact source response of the algorithm-derived field action

thm-ym-native-source-response

2_fractal_set/05_yang_mills_noether.md, line 1686.

[R61] Transient projected field equation

thm-algorithmic-transient-field-memory

3_fitness_manifold/04_field_equations.md, line 1641.

[R62] Field evolution conditional on its observed history

thm-algorithmic-field-filter

3_fitness_manifold/04_field_equations.md, line 1768.

[R63] Prediction-complete gauge descriptors from the actual transition

thm-sm-prediction-complete-descriptors

2_fractal_set/04_standard_model.md, line 2083.

[R64] Finite transition matrices converging to the completed channel theory

thm-sm-predictive-partition-convergence

2_fractal_set/04_standard_model.md, line 2201.

[R65] Closure implications and their converse

thm-closure-equivalence-cosmo

3_fitness_manifold/06_cosmology.md, line 351.

[R66] Exact and approximate Markov closure

prop-cosmo-generator-closure

3_fitness_manifold/06_cosmology.md, line 391.

[R67] Stationary response from a discrete Poisson equation

thm-algorithmic-stationary-response

3_fitness_manifold/05_holography.md, line 873.

[R68] Exact path KL identities

prop-algorithmic-path-kl-identities

3_fitness_manifold/05_holography.md, line 797.

[R69] Faithful exterior algebra of alternating recorded insertions

thm-lqft-oriented-word-algebra

2_fractal_set/03_lattice_qft.md, line 681.

[R70] Completely positive CAR evolution of the recorded contraction

thm-lqft-record-car-channel

2_fractal_set/03_lattice_qft.md, line 1018.

[R71] Separate local-net application to the recorded CAR representation

thm-ym-hk-record-instantiation

2_fractal_set/05_yang_mills_noether.md, line 9275.

[R72] Native fermion evolution, localization, and record covariance

thm-lqft-native-local-fermion-evolution

2_fractal_set/03_lattice_qft.md, line 1585.

[R73] Exact inherited-history covariance of native regional readouts

thm-lqft-inherited-history-covariance

2_fractal_set/03_lattice_qft.md, line 1332.

[R74] Existing LSI and native contraction bounds

cor-lqft-fock-lsi-transfer

2_fractal_set/03_lattice_qft.md, line 1186.

[R75] Distribution-valued compactness from the established stationary LSI

thm-ym-spacetime-fluctuation-compactness

2_fractal_set/05_yang_mills_noether.md, line 5512.

[R76] Drift and covariance equations for the complete selected update

prop-ym-complete-fluctuation-equations

2_fractal_set/05_yang_mills_noether.md, line 5750.

[R77] Native physical hierarchy of the full recorded color invariants

thm-ym-native-physical-gauge-hierarchy

2_fractal_set/05_yang_mills_noether.md, line 7629.

[R78] Geometry-fiber action in the common native continuum limit

thm-ym-native-fiber-continuum

2_fractal_set/05_yang_mills_noether.md, line 3914.

[R79] Nontrivial doublet fluctuations from the actual companion draw

thm-sm-native-doublet-fluctuations

2_fractal_set/04_standard_model.md, line 1087.

[R80] Nonzero momentum fluctuations in the established Euclidean paired model

cor-ym-paired-momentum-fluctuations

2_fractal_set/05_yang_mills_noether.md, line 6405.

[R81] Uniform integrability and reconstruction bounds for the full direct color algebra

lem-ym-physical-gauge-word-uniform-integrability

2_fractal_set/05_yang_mills_noether.md, line 7546.

[R82] Direct local bias estimate in specified normal coordinates

lem-continuum-local-bias

convergence_program/16_continuum_discharge.md, line 509.

[R83] A covariance condition sufficient for A4

lem-continuum-a4-mixing

convergence_program/16_continuum_discharge.md, line 597.

[R84] A sufficient bandwidth schedule for A6

lem-continuum-a6-scaling

convergence_program/16_continuum_discharge.md, line 687.

[R85] Conditional continuum consistency

cor-continuum-consistency-conditional

convergence_program/16_continuum_discharge.md, line 735.

[R86] Normalized sampling and the distinction between QSD and invariance

lem-continuum-a3-qsd-sampling

convergence_program/16_continuum_discharge.md, line 327.

[R87] Normalized importance sampling and variance

prop-monte-carlo-riemannian-latent

3_fitness_manifold/01_emergent_geometry.md, line 1116.

[R88] Continuum estimator error from the established joint LSI

cor-cst-inherited-lsi-consistency

2_fractal_set/02_causal_set_theory.md, line 966.

[R89] Density-corrected limit with sufficient sampling control

prop-density-corrected-limit
2_fractal_set/03_lattice_qft.md, line 2185.

[R90] Volume Element Transformation

thm-volume-element-transformation
3_fitness_manifold/07_computational_proxies.md, line 1155.

[R91] Wilson action consistency under geometric quadrature

thm-continuum-limit-ym
2_fractal_set/05_yang_mills_noether.md, line 4858.

[R92] Common continuum limit of the native source likelihood and descriptor action

thm-ym-continuum-source-action
2_fractal_set/05_yang_mills_noether.md, line 2714.

[R93] Time consistency for the specified transition family

thm-correct-continuum-limit
2_fractal_set/05_yang_mills_noether.md, line 5180.

[R94] Total observable error

thm-total-error-bound
convergence_program/13_quantitative_error_bounds.md, line 719.

[R95] Native response current in recorded spacetime

thm-ym-native-response-current
2_fractal_set/05_yang_mills_noether.md, line 2376.

[R96] Native response and the weak connection-variation identity

prop-ym-native-connection-variation-identity
2_fractal_set/05_yang_mills_noether.md, line 4158.

[R97] Exact graph Ward identity

lem-ym-discrete-ward
2_fractal_set/05_yang_mills_noether.md, line 4584.

[R98] Euclidean regularity from actual moment bounds

thm-os-os0-fg
2_fractal_set/05_yang_mills_noether.md, line 7063.

[R99] Symmetry of bosonic correlations

thm-os-os4-fg
2_fractal_set/05_yang_mills_noether.md, line 7085.

[R100] Physical reflection on the full native color algebra

prop-ym-native-color-reflected-matrices
2_fractal_set/05_yang_mills_noether.md, line 8050.

[R101] Algorithmic QFT from recorded evolution and native bounds

thm-ym-algorithmic-qft-synthesis
2_fractal_set/05_yang_mills_noether.md, line 9530.

[R102] Algorithmic Evolution of Frame-Averaged Twistor Channels

thm-effective-twistor-spectral-meaning
2_fractal_set/08_twistor_formulation.md, line 854.

[R103] Positive Transfer Representation of a Frame Correlator

cor-effective-twistor-positive-transfer
2_fractal_set/08_twistor_formulation.md, line 914.

[R104] Exact HK locality of the native multiplication representation

thm-ym-native-multiplication-locality
 2_fractal_set/05_yang_mills_noether.md, line 8890.

[R105] Survival of a uniformly controlled self-adjoint gap

thm-mass-gap-rg-fixed-point
 2_fractal_set/05_yang_mills_noether.md, line 5225.

[R106] Restriction of a gap to an invariant gauge sector

thm-gauge-sector-mass-gap
 2_fractal_set/05_yang_mills_noether.md, line 7017.

B Chapter contribution map

This map records the contributions used in the proof program. Chapter-local configurations, laws, and theorem hypotheses are retained when their results are composed. The experiment and calibration chapters supply observable definitions and diagnostics alongside the analytic results.

Microscopic definition and configured variants

Complete state, update order, parameter floors, metric/noise choices, and each variant’s own transition.

1_the_algorithm/01_algorithm_intuition.md
 02_fractal_gas_latent.md
 03_parameter_constraints.md
 04_gas_variants.md
 convergence_program/01_fragile_gas_framework.md
 02_euclidean_gas.md

Cloning, transport, and recurrence

Keystone pressure, reset and signed drift, barycenter/shape decomposition, kinetic estimates, moment/tail budgets, Harris and QSD convergence, quantified population/time limits.

convergence_program/03_cloning.md
 04_single_particle.md
 04_wasserstein_contraction.md
 05_kinetic_contraction.md
 06_convergence.md
 06a_structural_landscape_convergence.md
 07_discrete_qsd.md

Population law and exchangeability

Fixed-step rooted population map, exact balances, particle consistency, survival normalization, mass–shape convergence, empirical mixtures, fixed-row chaos.

convergence_program/08_mean_field.md
 09_propagation_chaos.md
 11_hk_convergence.md
 12_qsd_exchangeability_theory.md

Entropy, quantitative errors, and regularity

Poincaré and LSI, full-update entropy identities, hypocoercivity, observable error, normalized derivative calculus, smooth empirical fields, identified adaptive-law bounds.

convergence_program/10_kl_hypocoercive.md
 13_quantitative_error_bounds.md
 14_a_geometric_gas_c3_regularity.md
 14_b_geometric_gas_cinf_regularity_full.md
 15_kl_convergence.md
 17_geometric_gas.md

Causal and geometric continuum

Recorded causal order, slabs and trajectory clocks, regular metric coefficients, sampling correction, two estimator-error routes, operator reconstruction, holonomy and curvature comparison.

convergence_program/16_continuum_discharge.md
 2_fractal_set/02_causal_set_theory.md
 3_fitness_manifold/01_emergent_geometry.md
 02_scutoid_spacetime.md
 03_curvature_gravity.md

Exact derived finite fields

Marked characteristic hierarchy, exact density/current balances, conditional metric law, drift/covariance, field memory and filtering, coupled population/mechanical/metric system.

3_fitness_manifold/04_field_equations.md

Recorded gauge and matter theory

Lossless history, invariant coordinates, CAR dynamics, native action and sources, Ward identities, gauge hierarchies, Wilson consistency, spectral-gap passage.

2_fractal_set/01_fractal_set.md
 03_lattice_qft.md
 04_standard_model.md
 05_yang_mills_noether.md
 3_fitness_manifold/08_voronoi_wilson_loops.md

Spectral channels and empirical checks

Implemented channel and correlator definitions, algorithmic-time lag, source-grounded calibration, positive-transfer spectral criterion, observable diagnostics.

2_fractal_set/06_empirical_validation.md
 07_qft_calibration_report.md
 08_twistor_formulation.md
 09_qft_calibration.md
 partvi_experiments.md

Response, measurement, and geometric extensions

Actual-kernel path KL and stationary Poisson response, graph-cut and boundary observables, equilibrium centering and coarse closure, computational geometric probes, reference-transport covariance for measurement. These chapters develop additional observables and extensions of the same recorded system.

3_fitness_manifold/05_holography.md
 06_cosmology.md
 07_computational_proxies.md
 09_measurement.md