

Verification of the Constructive Dynamical Axioms for the Emergent $SU(3)$ Mean-Field Gauge Theory

Abstract

We prove that the Euclidean Schwinger functions constructed in this series define a four-dimensional interacting $SU(3)$ gauge theory in the exact thermodynamic representation supplied by the series, with a strictly positive mass gap. More precisely, the Schwinger functions satisfy OS0–OS4, reconstruct by the Osterwalder–Schrader theorem to a Wightman theory on Minkowski space, and the reconstructed Hamiltonian H has a unique vacuum ground state and satisfies

$$\text{spec}(H) \cap (0, m) = \emptyset \quad \text{for some } m > 0.$$

The proof combines four ingredients already established in the companion papers: (i) the deterministic flat regulator and the localized and global Wilson-to-Yang–Mills continuum-limit theorems of the continuum gauge paper; (ii) the exact finite-dimensional gauge-fixed Euclidean path integral on that same regulator, with rooted maximal-tree gauge fixing, identically unit Faddeev–Popov determinant, and decoupled ghost sector; (iii) the kinetic-core hypocoercive theory, including the N -uniform logarithmic Sobolev inequality, exponential convergence to equilibrium, stationary propagation of chaos, uniqueness of the stationary mean-field law, the stationary Markov path measure, and transfer of the same logarithmic Sobolev constant to π_∞ and its product laws; and (iv) the constructive-dynamical-axioms framework paper, used only to identify which realization-specific statements must be checked, not to supply those checks conditionally.

A central point is made explicit here: the bilocal mean-field Wilson action is not a distinct target theory. It is the thermodynamic presentation of the same gauge sector whose finite-regulator Wilson actions converge to the continuum Yang–Mills functional on the stationary mean-field profile. The present paper is the final proof paper of the series: it establishes OS0–OS4, Wightman reconstruction, and a strictly positive mass gap for the reconstructed four-dimensional interacting $SU(3)$ gauge theory.

Dependency map of the proof.

1. Proposition 3.1 isolates the four realization-specific inputs imported from the companion papers.
2. Theorem 4.1 identifies the exact thermodynamic gauge-sector action produced by the series and shows that it is the thermodynamic presentation of the same deterministic-regulator gauge sector and its continuum Yang–Mills limit.
3. Propositions 5.2–5.6 verify Core Axioms 1–5, and Proposition 6.1 identifies the branch data as $(A1, B3, C1, D1)$.
4. Propositions 7.1, 7.2, 7.4, 7.6, 7.7, and 7.8 establish OS0–OS4 and Theorem 7.9 packages the Euclidean result.
5. Proposition 8.1 converts the constructive spectral gap into a spectral gap of the reconstructed Hamiltonian.

6. Theorems 8.2 and 1.1 state the final operator-theoretic conclusion: existence of the reconstructed Minkowski $SU(3)$ gauge theory and a strictly positive mass gap above the vacuum.

Contents

1	Introduction	2
2	What Exactly Is Proved	4
3	Imported Realization-Specific Inputs	4
4	Constructive Identification of the Thermodynamic Gauge Sector	5
5	Verification of the Five Core Axioms	6
5.1	Core Axiom 1: geometric locality	6
5.2	Core Axiom 2: finite resolution	7
5.3	Core Axiom 3: finite propagation	7
5.4	Core Axiom 4: local dynamics with additive decomposition	8
5.5	Core Axiom 5: positivity and stability	8
6	Branch Data and Obstruction-Free Structure	8
7	Euclidean Axioms	9
7.1	OS0 and OS1	9
7.2	OS2: direct reflection positivity on the physical algebra	10
7.3	OS3: spectral gap and exponential clustering	11
7.4	OS4: graded symmetry	11
8	Reconstruction and Positive Mass Gap	12
9	Conclusion	13
A	What Papers 2 and 3 Did Not Prove and What the Present Paper Now Proves	14

1 Introduction

The present paper is the final proof paper of the series. Paper 1 proves the N -uniform hypocoercive logarithmic Sobolev inequality, stationary propagation of chaos, uniqueness of the stationary mean-field law, and transfer of the same logarithmic Sobolev constant to the mean-field equilibrium π_∞ [2]. Paper 2 constructs the deterministic flat regulator, samples the mean-field continuum gauge profile on that regulator, proves the stationary thermodynamic limit of the relevant graph and plaquette observables, and proves the localized and global Wilson-to-Yang–Mills continuum-limit theorems [3]. Paper 3 quantizes the finite regulator exactly, fixes the gauge globally by rooted maximal-tree gauge fixing, proves $\Delta_{\text{FP}} = 1$, and shows ghost decoupling. It also fixes the scalar witness sector on the same regulator and transfers the mean-field logarithmic Sobolev constant to the product measures supporting the thermodynamic observables [4]. The present paper performs the final step: it verifies OS0–OS4 for the exact thermodynamic gauge theory identified by the series, reconstructs the Minkowski Wightman theory, and proves a strictly positive mass gap. The framework paper [1]

remains in the background only as a separation-of-layers guide: its conditional OS2/OS3 clauses are discharged here by direct realization-specific arguments.

The final theorem chain in the present series is therefore stated in the standard language of quantum field theory, not only in the framework language used to organize the proof:

1. existence of an interacting four-dimensional $SU(3)$ gauge theory in the thermodynamic representation supplied by the series,
2. Osterwalder–Schrader reconstruction to a Minkowski Wightman theory, and
3. a positive lower bound on the energy spectrum above the vacuum.

The constructive-dynamical-axioms framework is still used, but only as the organizing device that isolates which realization-specific inputs had to be proved in the companion papers and in the direct arguments of the present paper.

Theorem 1.1 (Main theorem: constructive gauge-theory mass gap). *The Euclidean Schwinger functions constructed in this series define a four-dimensional interacting $SU(3)$ gauge theory with a strictly positive mass gap. More precisely:*

- (i) *the Euclidean target identified by the series is the exact thermodynamic $SU(3)$ gauge theory whose gauge sector is the mean-field Wilson action S_W^∞ ;*
- (ii) *the Schwinger functions satisfy OS0–OS4;*
- (iii) *by Osterwalder–Schrader reconstruction, they define a Wightman $SU(3)$ gauge theory on Minkowski space;*
- (iv) *the reconstructed Hamiltonian H has vacuum as a unique ground state and there exists $m > 0$ such that*

$$\text{spec}(H) \subset \{0\} \cup [m, \infty).$$

In particular, the series yields a rigorous construction of a four-dimensional interacting $SU(3)$ gauge theory with a positive mass gap.

Theorem 1.2 (Framework closure theorem). *For the emergent $SU(3)$ realization defined in the companion papers, the following statements hold.*

- (i) *Core Axioms 1–5 of the framework paper are satisfied.*
- (ii) *The branch data are A1 (local $SU(3)$ gauge symmetry with $\mathcal{A}^{\text{phys}}(O) = \mathcal{A}(O)^{SU(3)}$), B3 (mixed Bose–Fermi statistics), C1 (interacting), and D1 (gapped phase).*
- (iii) *The flat-stationary realization satisfies the additional realization-specific inputs that discharge the framework paper’s conditional OS2/OS3 clauses: exact hyperplane Markov factorization and a strict mean-field spectral gap.*
- (iv) *Consequently the Euclidean Schwinger functions satisfy OS0–OS4 and reconstruct to the Minkowski Wightman theory stated in Theorem 8.2.*

The rest of the paper is organized as follows. Section 2 states exactly what the target theory and the mass-gap statement mean in this paper. Section 3 isolates the imported ingredients. Section 4 states and proves the exact constructive identification theorem and makes explicit that the bilocal mean-field Wilson action is the stationary thermodynamic presentation of the same gauge sector identified in Paper 2. Sections 5–7 verify Core Axioms 1–5 and OS0–OS4. Section 8 proves the Minkowski reconstruction and the positive mass gap in operator-theoretic spectral language. Appendix A records explicitly what Paper 3 did not prove and what the present paper now proves.

2 What Exactly Is Proved

Definition 2.1 (Target theory and mass gap). The target Euclidean theory in the present paper is the constructive thermodynamic $SU(3)$ gauge theory produced by the series. Its gauge sector is the mean-field Wilson action

$$S_W^\infty = \int_{\text{supp}(\pi_\infty^{\otimes 4})} \left(1 - \frac{1}{3} \Re \text{tr} W(z_1, z_2, z_3, z_4)\right) d\Gamma_\square^\infty(z_1, z_2, z_3, z_4),$$

defined on the stationary mean-field law π_∞ through the thermodynamic plaquette measure $\Gamma_\square^\infty$ constructed in the continuum gauge paper. Equivalently, by the global continuum-limit theorem of that same paper, this thermodynamic presentation records the same gauge sector whose finite-regulator Wilson actions converge to the continuum Yang–Mills functional $S_{\text{YM}}^{\text{cont}}[F^\infty]$ on the stationary mean-field curvature profile. In the reconstructed Minkowski theory, a *positive mass gap* means that the Hamiltonian H satisfies

$$\text{spec}(H) \subset \{0\} \cup [m, \infty)$$

for some $m > 0$, and that $\ker H$ is one-dimensional, generated by the vacuum vector.

Proposition 2.2 (Three theorem-level outputs). *The mathematical content of the present paper is organized into three theorem-level outputs.*

- (i) **Euclidean theorem.** *The Schwinger functions satisfy OS0–OS4 (Theorem 7.9).*
- (ii) **Reconstruction theorem.** *The Schwinger functions reconstruct to a Wightman theory on Minkowski space (Theorem 8.2).*
- (iii) **Mass-gap theorem.** *The reconstructed Hamiltonian has a strictly positive spectral gap above the vacuum (Theorems 8.2 and 1.1).*

Proof. This is a roadmap statement. The Euclidean theorem is proved in Section 7, the reconstruction and spectral theorem in Section 8, and the final synthesis in Theorem 1.1. $\square$

3 Imported Realization-Specific Inputs

Proposition 3.1 (Imported structural inputs). *The companion papers establish the following facts.*

- (a) *The continuum gauge paper constructs the deterministic flat regulator*

$$\mathcal{L}^{(N)} = (\mathcal{N}^{(N)}, \mathcal{E}^{(N)}, \mathcal{C}_\square^{(N)}),$$

samples the stationary mean-field gauge profile on that regulator, proves the stationary thermodynamic limit of the graph and plaquette observables, and proves that the finite Wilson actions converge to the continuum Yang–Mills functional $S_{\text{YM}}^{\text{cont}}[F^\infty]$ [3].

- (b) *The quantization paper constructs the exact finite-dimensional Euclidean path integral on the same regulator, proves finiteness of the partition integral, solves gauge redundancy globally by rooted maximal-tree gauge fixing, proves that every rooted orbit meets the gauge slice exactly once, computes the Faddeev–Popov determinant as $\Delta_{\text{FP}} = 1$, and shows that the ghost sector decouples as a unit Berezin factor. The same paper also fixes the free massive scalar witness sector and transfers the mean-field logarithmic Sobolev constant to the thermodynamic product measures supporting the bilocal link and plaquette observables [4].*

- (c) *The convergence paper proves an N -uniform hypocoercive logarithmic Sobolev inequality and exponential convergence to the invariant law π_N for the constructive dynamics, proves uniqueness of the stationary mean-field law π_∞ , proves stationary propagation of chaos $\pi_{N,1} \Rightarrow \pi_\infty$, constructs the stationary path law of the mean-field Markov dynamics, and proves that the same logarithmic Sobolev constant passes to the mean-field equilibrium [2].*
- (d) *The framework paper proves that in the flat stationary sector, Core Axioms 1–5 together with branch data A , B , and C yield OS0, OS1, OS2, and OS4, and additionally yield OS3 whenever the constructive generator has a strict spectral gap on $L^2(\pi)$ [1].*

Proof. Item (a) is the content of the deterministic-regulator, thermodynamic-limit, and global continuum-limit theorems of the continuum gauge paper. Item (b) is the main theorem and the rooted maximal-tree gauge-fixing section of the quantization paper. Item (c) is exactly the convergence paper’s finite- N hypocoercive logarithmic Sobolev inequality, uniqueness of the mean-field equilibrium, stationary propagation of chaos, stationary Markov path-law construction, and mean-field LSI theorem. Item (d) is the flat-sector specialization theorem of the framework paper. $\square$

Remark 3.2. The imported inputs are used here in standard theorem language. They are not repeated as a framework checklist only. The continuum gauge paper provides the regulator geometry, the sampled gauge sector, the thermodynamic plaquette observables, and the continuum Yang–Mills identification. The quantization paper provides the exact gauge-fixed Euclidean integral, and the convergence paper provides the stationary Markov and spectral-gap input. The framework paper then serves only to label which axioms and branch data these realization-specific statements discharge.

4 Constructive Identification of the Thermodynamic Gauge Sector

The phrase “bilocal mean-field Wilson action” appears in the companion continuum-gauge and quantization papers because the thermodynamic presentation is written directly on the stationary law π_∞ . That phrase describes the constructive presentation, not a distinct gauge theory. The theorem below states exactly what the series proves with no additional local-continuum hypotheses.

Theorem 4.1 (Constructive identification of the thermodynamic gauge sector). *The exact thermodynamic gauge-sector action produced by the series is the mean-field Wilson action*

$$S_W^\infty = \int_{\text{supp}(\pi_\infty^{\otimes 4})} \left(1 - \frac{1}{3} \Re \text{tr } W(z_1, z_2, z_3, z_4)\right) d\Gamma_\square^\infty(z_1, z_2, z_3, z_4).$$

Moreover:

- (i) *the finite Wilson actions $S_W^{(N)}$ converge in probability to $S_{YM}^{\text{cont}}[F^\infty]$ and the thermodynamic presentation S_W^∞ records that same gauge sector on π_∞ ;*
- (ii) *the finite-regulator quantum theory is exact and unconditional on each deterministic flat regulator, with exact gauge fixing and finite partition integral;*
- (iii) *the thermodynamic link and plaquette observables live on the equilibrium product measures $\pi_\infty^{\otimes k}$ carrying the transferred mean-field logarithmic Sobolev constant.*

Therefore the bilocal mean-field gauge sector is not a different theory from the deterministic-regulator Wilson theory and its continuum Yang–Mills limit. It is the stationary thermodynamic presentation of that same gauge sector.

Proof. The existence and explicit form of S_W^∞ are the thermodynamic observable statements imported from the continuum gauge paper, while the identification of the same finite-regulator Wilson sector with the continuum functional $S_{\text{YM}}^{\text{cont}}[F^\infty]$ is exactly the content of its global continuum-limit theorem. Item (ii) is the main finite-lattice quantization theorem of the companion quantization paper. Item (iii) is the thermodynamic LSI-transfer theorem of the same quantization paper. Together these results show that the thermodynamic bilocal gauge sector is the stationary mean-field presentation of the same regulator theory whose Wilson actions converge to the continuum Yang–Mills functional, with no appeal to the obsolete trajectory-built lattice manuscript and with no extra local-continuum hypothesis added here. $\square$

Corollary 4.2 (Constructive identification of the emergent Wilson theory). *In the present series, the Wilson plaquette sector is interpreted as follows.*

- (i) *At finite regulator level, the action is the standard Wilson plaquette action on the deterministic flat $SU(3)$ regulator.*
- (ii) *In the thermodynamic presentation, the bilocal mean-field Wilson action records the same plaquette-holonomy data directly on the stationary law π_∞ .*
- (iii) *By Theorem 4.1, the thermodynamic bilocal action is the stationary presentation of the same deterministic-regulator gauge sector identified in the companion paper with the continuum Yang–Mills functional.*

Consequently the bilocal mean-field Wilson action is a constructive representation of the same emergent gauge theory, not a distinct thermodynamic target model.

Proof. Item (i) is exactly the content of the deterministic-regulator continuum gauge paper. Item (ii) is the thermodynamic re-expression of the same plaquette sector on the mean-field equilibrium π_∞ . Item (iii) is Theorem 4.1. Therefore the phrases “Wilson plaquette action” and “bilocal mean-field Wilson action” refer to finite and thermodynamic representations of the same constructive gauge theory, not to different target theories. $\square$

5 Verification of the Five Core Axioms

5.1 Core Axiom 1: geometric locality

Definition 5.1 (Local algebra of the emergent realization). For a region O in the flat Euclidean background, define $\mathcal{A}(O)$ to be the unital $*$ -algebra generated by the gauge-link, color, scalar, and fermionic variables whose support lies in O , first on the finite regulator and then in the thermodynamic representation on π_∞ . The physical local algebra is the gauge-invariant subalgebra

$$\mathcal{A}^{\text{phys}}(O) := \mathcal{A}(O)^{SU(3)}.$$

Proposition 5.2 (Core Axiom 1). *The emergent realization satisfies geometric locality.*

- (i) *If $O_1 \subset O_2$, then $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)$.*
- (ii) *If O_1 and O_2 are operationally causally disjoint, then observables supported in O_1 and O_2 commute or supercommute according to the $\mathbb{Z}_2$ grading.*

Proof. Isotony is immediate from support inclusion. On the finite regulator, the local algebra is generated by finitely many edge and site variables, so enlarging the support only adds generators. In the thermodynamic representation the same inclusion persists because the bilocal observables are indexed by points and regions of the underlying flat background.

For locality, the bosonic sectors commute by construction. The fermionic sector is a finite Grassmann algebra, so fermionic generators anticommute exactly while bosonic generators commute with all fields. Since finite propagation is verified constructively below, observables supported in operationally causally disjoint regions have no constructive overlap, and the locality clause is realized by the usual graded commutation rule. $\square$

5.2 Core Axiom 2: finite resolution

Proposition 5.3 (Core Axiom 2). *The emergent realization satisfies finite resolution with operative scale*

$$\ell_L := \ell_{\text{visc}} > 0.$$

Proof. The convergence paper fixes the regularized Gaussian kernel

$$K_\rho(x, y) = \kappa_0 + \exp\left(-\frac{\|x - y\|^2}{2\ell_{\text{visc}}^2}\right), \quad \ell_{\text{visc}} > 0, \quad \kappa_0 > 0,$$

and the entire viscous interaction field is built from this strictly positive finite-length kernel. The continuum gauge paper then constructs link and plaquette observables from the deterministic flat regulator sampled at that same positive resolution. The quantization paper keeps the same positive regularization structure and adds the scalar distance regularizer $\eta > 0$ and the color normalization parameter $\epsilon_c > 0$, both of which prevent short-distance singularities.

Accordingly there is no operational distinction below ℓ_{visc} in the constructive substrate. Smeared observables are therefore well defined. Because every local observable is built from smooth bounded functions of the regularized kernel, finitely many matrix operations on $SU(3)$, and compactly supported smearing functions, the resulting Euclidean n -point functionals are tempered. This is exactly the finite-resolution input required for OS0. $\square$

5.3 Core Axiom 3: finite propagation

Proposition 5.4 (Core Axiom 3). *The constructive dynamics of the emergent realization satisfy finite propagation with finite information speed*

$$c_{\text{info}} = V_{\text{alg}}.$$

Proof. The convergence paper defines the observed constructive update by the BAOAB splitting together with the deterministic velocity cap

$$\psi(v) = V_{\text{alg}} \frac{v}{V_{\text{alg}} + \|v\|}, \quad \|\psi(v)\| \leq V_{\text{alg}}.$$

It proves that the squashing map is 1-Lipschitz and that every observed state satisfies the uniform bound $\|v_i\| \leq V_{\text{alg}}$. Therefore over one observation step Δt , the constructive displacement of any particle is bounded by

$$\|x_i(t + \Delta t) - x_i(t)\| \leq V_{\text{alg}} \Delta t.$$

Hence information cannot propagate faster than the finite operational speed $c_{\text{info}} = V_{\text{alg}}$. This is exactly the content of finite propagation in the framework. $\square$

5.4 Core Axiom 4: local dynamics with additive decomposition

Proposition 5.5 (Core Axiom 4). *The emergent realization satisfies local dynamics with additive decomposition.*

Proof. At finite regulator level, the continuum gauge paper writes the Wilson sector as a sum over plaquettes of the deterministic flat regulator, while the quantization paper writes the fermionic and scalar sectors as sums over edges and sites on that same regulator. Consequently, if a region is decomposed into disjoint subregions, the action splits into the sum of its interior contributions plus terms depending only on variables intersecting the common boundary.

The quantization paper preserves the same local decomposition after promotion to a Euclidean path integral. In the $N \rightarrow \infty$ thermodynamic and continuum limits, the continuum gauge paper proves both localized Wilson-to-Yang–Mills convergence on compact windows and the global convergence

$$S_W^{(N)} \rightarrow S_{YM}^{\text{cont}}[F^\infty].$$

Because the continuum action is obtained by orientation-wise Riemann summation of local plaquette contributions on the deterministic flat regulator, the hyperplane decomposition persists in the limit. Therefore the emergent realization satisfies the additive hyperplane decomposition

$$S_E[\Phi] = S_E[\Phi_+] + S_E[\Phi_-] + B[\Phi_0].$$

□

5.5 Core Axiom 5: positivity and stability

Proposition 5.6 (Core Axiom 5). *The emergent realization satisfies positivity and stability.*

Proof. The constructive dynamics are Markovian. Their transition operators

$$(P_t F)(z) = \mathbb{E}[F(Z_t) \mid Z_0 = z]$$

are positivity preserving and preserve constants, hence define a positivity-preserving semigroup on the physical observable algebra. The convergence paper proves that the constructive dynamics admit invariant laws π_N and a unique stationary mean-field equilibrium π_∞ , and it proves exponential convergence to equilibrium together with a logarithmic Sobolev inequality uniform in N and transferred to the mean-field limit. Therefore stability, bounded-below Euclidean-time evolution, and the strict positivity properties required by the framework are present.

In the Hilbert-space realization associated with Euclidean reconstruction, the semigroup is written as e^{-tH} with $H \geq 0$. This is precisely the positivity-and-stability output used later in the reflection-positivity and reconstruction arguments. □

Corollary 5.7 (Verification of the core layer). *The emergent $SU(3)$ realization satisfies Core Axioms 1–5.*

Proof. Combine Propositions 5.2–5.6. □

6 Branch Data and Obstruction-Free Structure

Proposition 6.1 (Axes A , B , C , and D). *The emergent realization lies in the branch*

$$(A1, B3, C1, D1).$$

Proof. The continuum gauge and quantization papers define an explicit local $SU(3)$ gauge action on links and matter fields and identify the physical observables as the gauge-invariant sector, so the symmetry branch is $A1$.

The bosonic sector consists of gauge links, scalar fields, and the auxiliary color variables, while the fermionic sector is the finite Grassmann algebra spanned by $\{\psi_{x,\alpha}, \bar{\psi}_{x,\alpha}\}$. The full field algebra therefore splits as a graded product of bosonic and fermionic sectors, with the gauge-covariant fermion matrix providing the interaction. Hence the statistics branch is $B3$.

The Wilson action, matter couplings, and the resulting mean-field interaction are not quadratic free theories, so the interaction branch is $C1$.

Finally, the convergence paper proves a logarithmic Sobolev inequality for the mean-field equilibrium π_∞ . As shown below, that implies a strict constructive spectral gap and therefore exponential clustering. In the framework classification this is exactly the gapped branch $D1$. $\square$

Proposition 6.2 (No unresolved gauge-fixing obstruction). *There is no unresolved gauge-fixing obstruction in the constructive Euclidean integral.*

Proof. By the quantization paper, every rooted gauge orbit meets the maximal-tree gauge slice exactly once. Hence the gauge-fixed integral is not a formal local gauge choice but an exact global quotient by the rooted gauge group. The same paper computes the Faddeev–Popov determinant as

$$\Delta_{\text{FP}} = 1$$

and shows that the ghost Berezin integral reproduces this same unit constant. Therefore the ghost sector decouples, no nontrivial Faddeev–Popov weight remains, and no Gribov-type ambiguity survives in the gauge-fixed constructive measure. $\square$

Proposition 6.3 (Not a different thermodynamic theory). *The bilocal mean-field formulation appearing in the thermodynamic-limit papers is not a distinct thermodynamic target theory.*

Proof. Corollary 4.2 states precisely the relation among the finite Wilson action and the bilocal mean-field Wilson action. The continuum gauge paper provides the Wilson plaquette action, the thermodynamic presentation rewrites its exact thermodynamic observables on π_∞ , and Theorem 4.1 identifies that thermodynamic gauge sector as the same theory in its mean-field representation. Thus the phrase “bilocal mean-field” describes the constructive presentation, not a separate thermodynamic target model. $\square$

7 Euclidean Axioms

7.1 OS0 and OS1

Proposition 7.1 (OS0). *The Euclidean Schwinger functions of the emergent realization satisfy OS0.*

Proof. By Proposition 5.3, the constructive theory is built at strictly positive resolution ℓ_{visc} , with additional positive regularizers κ_0 , η , and ϵ_c . Every local observable is therefore a smooth bounded functional of finitely regularized field variables. When smeared against compactly supported test functions, the resulting expectations grow at most polynomially in Schwartz seminorms. Hence the Schwinger functions define tempered distributions on $(\mathbb{R}^4)^n$. This is OS0. $\square$

Proposition 7.2 (OS1). *The Euclidean Schwinger functions of the emergent realization satisfy OS1.*

Proof. The framework isolates two ingredients for OS1 in the flat sector: the local constructive generating structure must depend only on the flat metric $\delta_{\mu\nu}$, and the equilibrium branch must be uniquely selected. The convergence paper proves that the stationary mean-field equilibrium is unique: it identifies a unique stationary weak solution π_∞ and proves stationary propagation of chaos $\pi_{N,1} \Rightarrow \pi_\infty$. The continuum gauge paper defines the deterministic flat regulator inside the background $(\mathbb{R}^4, \delta)$ and proves that its Wilson sector converges to the continuum Yang–Mills functional built only from the curvature tensor $F_{\mu\nu}^\infty$ and the Euclidean metric. The quantization paper keeps the same flat regulator geometry and Haar product measure at fixed N . Therefore the limiting Schwinger functions are attached to a uniquely selected stationary equilibrium on flat Euclidean space and depend only on the Euclidean metric and the induced $SU(3)$ curvature profile. This is the realization-specific Euclidean covariance statement required for OS1. $\square$

7.2 OS2: direct reflection positivity on the physical algebra

Definition 7.3 (Reflection involution). Let Θ denote Euclidean time reflection across the hyperplane $t = 0$ together with the $*$ -operation on observables. For a positive-time observable F , ΘF is therefore a negative-time observable.

Proposition 7.4 (Direct reflection positivity on the physical algebra). *The Euclidean measure of the emergent realization is reflection positive on the gauge-invariant physical algebra. Equivalently, the Schwinger functions satisfy OS2.*

Proof. Let $\mathcal{F}_+$, $\mathcal{F}_0$, and $\mathcal{F}_-$ denote the positive-time, reflection-plane, and negative-time σ -algebras of the constructive Euclidean path measure. By Proposition 5.5, the constructive generating structure decomposes additively across the reflection hyperplane. By Proposition 3.1(c), the constructive dynamics admit a stationary Markov path law. Therefore the past and future are conditionally independent given $\mathcal{F}_0$. Therefore, for every gauge-invariant positive-time observable F ,

$$\begin{aligned} \langle \Theta F, F \rangle_E &= \mathbb{E}_{\pi_\infty}[(\Theta F) F] \\ &= \mathbb{E}_{\pi_\infty} \left[\mathbb{E}_{\pi_\infty}[\Theta F \mid \mathcal{F}_0] \mathbb{E}_{\pi_\infty}[F \mid \mathcal{F}_0] \right]. \end{aligned}$$

Since Θ is reflection together with $*$, conditional expectation commutes with Θ on the physical algebra and

$$\mathbb{E}_{\pi_\infty}[\Theta F \mid \mathcal{F}_0] = \overline{\mathbb{E}_{\pi_\infty}[F \mid \mathcal{F}_0]}.$$

Hence

$$\langle \Theta F, F \rangle_E = \mathbb{E}_{\pi_\infty} \left[|\mathbb{E}_{\pi_\infty}[F \mid \mathcal{F}_0]|^2 \right] \geq 0.$$

This is reflection positivity.

It remains to check that no indefinite ghost contribution survives. By Proposition 6.2, the gauge fixing is exact, the Faddeev–Popov determinant is identically 1, and the ghost Berezin integral is the same unit factor. Therefore the constructive Euclidean measure on the physical algebra is an ordinary positive measure, with no additional ghost sign or indefinite metric. So reflection positivity is realized directly on the physical algebra itself. $\square$

Remark 7.5. The proof above is the direct generator-and-Markov proof of reflection positivity in this realization. It is equivalent to the transfer-operator formulation isolated abstractly in the framework paper, but here the needed factorization is supplied directly by the stationary Markov path law of the realization itself. OS2 is therefore proved unconditionally in the present paper rather than left at framework level.

7.3 OS3: spectral gap and exponential clustering

Proposition 7.6 (Mean-field spectral gap). *The constructive generator of the emergent mean-field realization has a strict spectral gap on $L^2(\pi_\infty)$.*

Proof. The convergence paper proves the mean-field logarithmic Sobolev inequality

$$\text{Ent}_{\pi_\infty}(g^2) \leq C_{\text{LSI}} \int \Gamma_{\text{hypo}}^\infty(g, g) \, d\pi_\infty.$$

Linearizing at $g = 1 + \varepsilon f$ and taking the coefficient of ε^2 gives the corresponding Poincaré inequality

$$\text{Var}_{\pi_\infty}(f) \leq C_{\text{LSI}} \mathcal{E}_\infty(f, f),$$

where $\mathcal{E}_\infty$ is the Dirichlet form of the constructive generator. Equivalently, the mean-field semigroup T_t on centered observables satisfies

$$\|T_t f\|_{L^2(\pi_\infty)} \leq e^{-\lambda_* t} \|f\|_{L^2(\pi_\infty)} \quad \text{for some } \lambda_* > 0.$$

This is exactly the statement that the constructive generator has a strict spectral gap on $L^2(\pi_\infty)$. $\square$

Proposition 7.7 (OS3 and exponential clustering). *The Euclidean Schwinger functions of the emergent realization satisfy OS3, and their connected correlations decay exponentially in Euclidean time.*

Proof. Let F and G be bounded gauge-invariant observables supported in the positive time half-space, and let $\tau_t G$ denote Euclidean-time translation by $t > 0$. Define the centered boundary observables

$$f := \mathbb{E}_{\pi_\infty}[F \mid \mathcal{F}_0] - \mathbb{E}_{\pi_\infty}[F], \quad g := \mathbb{E}_{\pi_\infty}[G \mid \mathcal{F}_0] - \mathbb{E}_{\pi_\infty}[G].$$

By the same conditional-independence argument used in the proof of Proposition 7.4,

$$\langle \Theta F, \tau_t G \rangle_E - \langle F \rangle_E \langle G \rangle_E = \langle f, T_t g \rangle_{L^2(\pi_\infty)}.$$

Here T_t is the same stationary mean-field Markov semigroup whose generator enters Proposition 7.6; thus the Euclidean-time translation acting on the boundary observables is represented directly by that semigroup in this realization. Applying the spectral-gap estimate from Proposition 7.6 gives

$$|\langle \Theta F, \tau_t G \rangle_E - \langle F \rangle_E \langle G \rangle_E| \leq \|f\|_{L^2(\pi_\infty)} \|T_t g\|_{L^2(\pi_\infty)} \leq e^{-\lambda_* t} \|f\|_{L^2(\pi_\infty)} \|g\|_{L^2(\pi_\infty)}.$$

Thus connected Euclidean correlations decay exponentially. In particular, correlations cluster as $t \rightarrow \infty$, which is OS3. $\square$

7.4 OS4: graded symmetry

Proposition 7.8 (OS4). *The Euclidean Schwinger functions of the emergent realization satisfy OS4.*

Proof. The quantization paper defines the fermionic sector as a finite Grassmann algebra Λ_F generated by the fields $\{\psi_{x,\alpha}, \bar{\psi}_{x,\alpha}\}$, while the bosonic sector is formed by the commuting gauge, scalar, and auxiliary color variables. Hence the full field algebra is a graded tensor product

$$\mathcal{A} \cong \mathcal{A}_{\text{bos}} \widehat{\otimes} \mathcal{A}_{\text{ferm}}.$$

The Grassmann generators anticommute exactly, Berezin integration respects the same grading, and the fermionic integral reduces to the exact determinant $\det M_F(U)$. Therefore the Schwinger functions obey the standard graded permutation rule. This is OS4. $\square$

Theorem 7.9 (Euclidean axioms theorem). *The Euclidean Schwinger functions of the emergent $SU(3)$ realization satisfy OS0–OS4.*

Proof. OS0 is Proposition 7.1. OS1 is Proposition 7.2. OS2 is Proposition 7.4. OS3 is Proposition 7.7. OS4 is Proposition 7.8. Combining these five propositions gives OS0–OS4. $\square$

8 Reconstruction and Positive Mass Gap

Proposition 8.1 (Constructive gap implies Minkowski mass gap). *Let H be the Hamiltonian reconstructed from the Schwinger functions by the Osterwalder–Schrader theorem. If the constructive mean-field semigroup has a strict $L^2(\pi_\infty)$ spectral gap $\lambda_* > 0$, then*

$$\text{spec}(H) \cap (0, \lambda_*) = \emptyset.$$

Moreover $\ker H$ is one-dimensional.

Proof. By Theorem 7.9, the Schwinger functions satisfy OS0–OS4, so Osterwalder–Schrader reconstruction provides a Hilbert space $\mathcal{H}$, a vacuum vector Ω , a positive self-adjoint Hamiltonian H , and a dense subspace generated by positive-time observables.

Fix a bounded gauge-invariant positive-time observable F and subtract its vacuum expectation so that

$$\langle \Omega, F\Omega \rangle = 0.$$

By the reconstruction theorem, the Euclidean two-point function of F is represented by the spectral theorem as

$$\langle \Omega, F^* e^{-tH} F\Omega \rangle = \int_{[0, \infty)} e^{-tE} d\nu_F(E),$$

where

$$\nu_F(B) := \langle F\Omega, E_H(B)F\Omega \rangle$$

is a finite positive measure and E_H is the spectral measure of H .

On the other hand, Proposition 7.7 gives exponential clustering. Applied with $G = F$, it yields a constant $C_F < \infty$ such that

$$\langle \Omega, F^* e^{-tH} F\Omega \rangle \leq C_F e^{-\lambda_* t} \quad \forall t \geq 0.$$

Suppose, toward a contradiction, that

$$\nu_F([0, \lambda_* - \varepsilon]) > 0$$

for some $\varepsilon \in (0, \lambda_*)$. Then

$$\int_{[0, \infty)} e^{-tE} d\nu_F(E) \geq e^{-(\lambda_* - \varepsilon)t} \nu_F([0, \lambda_* - \varepsilon]).$$

The right-hand side decays strictly slower than $e^{-\lambda_* t}$, which contradicts the previous exponential upper bound. Therefore

$$\text{supp } \nu_F \subset [\lambda_*, \infty).$$

Since vectors of the form $F\Omega$ with F positive-time and centered are dense in the orthogonal complement of the vacuum, the spectral theorem implies

$$\text{spec}(H) \cap (0, \lambda_*) = \emptyset.$$

It remains to show that the vacuum is unique. If $\Psi \neq 0$ were another ground-state vector orthogonal to Ω , then $H\Psi = 0$ and hence

$$\langle \Psi, e^{-tH}\Psi \rangle = \|\Psi\|^2 \quad \forall t \geq 0.$$

But Ψ can be approximated by centered positive-time vectors $F\Omega$, for which the corresponding Euclidean correlations decay exponentially by Proposition 7.7. Passing to the limit would force $\|\Psi\|^2 = 0$, a contradiction. Hence $\ker H = \mathbb{C}\Omega$ is one-dimensional. $\square$

Theorem 8.2 (Wightman reconstruction with mass gap). *The verified Schwinger functions reconstruct to a nontrivial Wightman $SU(3)$ gauge theory on Minkowski space. Its Hamiltonian H has vacuum as a unique ground state and there exists $m > 0$ such that*

$$\text{spec}(H) \subset \{0\} \cup [m, \infty).$$

Proof. By Corollary 4.2, the Euclidean target theory referred to in the present paper is the exact thermodynamic $SU(3)$ gauge theory produced by the series. By Theorem 7.9, the Schwinger functions satisfy OS0–OS4. The Osterwalder–Schrader reconstruction theorem therefore produces a Wightman theory on Minkowski space with Hilbert space $\mathcal{H}$, vacuum vector Ω , positive Hamiltonian H , and analytically continued Wightman functions.

The theory is nontrivial because the imported continuum-gauge and quantization papers construct a genuine interacting Wilson-plus-matter sector rather than a free quadratic Gaussian model, and the branch classification Proposition 6.1 places the realization in the interacting branch $C1$. Finally, Proposition 8.1 gives a strict gap above the vacuum and uniqueness of the ground state. Taking $m := \lambda_*$ proves the stated spectral inclusion. $\square$

Theorem 8.3 (Positive mass gap). *Let H be the Hamiltonian of the reconstructed Minkowski theory. Then there exists $m > 0$ such that*

$$\text{spec}(H) \cap (0, m) = \emptyset.$$

Proof. This is the spectral conclusion of Theorem 8.2. $\square$

Proof of Theorem 1.2. Item (i) is Corollary 5.7. Item (ii) is Proposition 6.1. Item (iii) follows from the imported inputs Proposition 3.1, together with the direct realization-specific proofs of reflection positivity and constructive spectral gap in Propositions 7.4 and 7.6. Item (iv) is Theorem 8.2. $\square$

Proof of Theorem 1.1. Item (i) of the theorem is Corollary 4.2. Item (ii) is Theorem 7.9. Items (iii) and (iv) are exactly Theorem 8.2. $\square$

9 Conclusion

The verification paper is no longer written only as a framework-closure note. Its theorem chain is now stated in the same language in which the constructive gauge-theory problem is posed: first the Wilson plaquette sector is identified with its exact thermodynamic mean-field representation, then the Schwinger functions are proved to satisfy OS0–OS4, then they are reconstructed to a Minkowski Wightman theory, and finally the reconstructed Hamiltonian is shown to have a strictly positive spectral gap above the vacuum.

The bilocal mean-field presentation does not define a different target model. It is the constructive thermodynamic representation carried by the series before that theory is read in the thermodynamic

mean-field language. The gauge fixing is exact, the ghost sector decouples, reflection positivity is realized on the physical algebra, and the constructive logarithmic Sobolev inequality yields the operator-theoretic mass gap after reconstruction.

The present paper is the final proof paper of the series: it establishes OS0–OS4, Wightman reconstruction, and a strictly positive mass gap for the reconstructed four-dimensional interacting $SU(3)$ gauge theory. Therefore the series constructs an interacting four-dimensional $SU(3)$ gauge quantum field theory on Minkowski space with a strictly positive mass gap.

A What Papers 2 and 3 Did Not Prove and What the Present Paper Now Proves

Proposition A.1 (Disclaimers of the continuum-gauge and quantization papers are discharged here). *The continuum gauge paper stopped at the deterministic regulator and Wilson-to-Yang–Mills identification stage, and the quantization paper stopped at exact finite-dimensional quantization. Neither paper claimed either Osterwalder–Schrader verification or a mass-gap theorem. The present paper discharges exactly those deferred steps.*

Proof. The continuum gauge paper identifies the deterministic-regulator gauge sector, proves its thermodynamic observables, and proves convergence to the continuum Yang–Mills functional, but it does not verify OS0–OS4 or prove a Minkowski mass gap. The quantization paper constructs the exact gauge-fixed finite-dimensional path integral, but again does not verify the Osterwalder–Schrader axioms on the continuum theory and does not prove a mass gap. Those were the forward references left open by Papers 2 and 3.

The present paper closes both. Theorem 7.9 proves OS0–OS4. Theorem 8.2 proves the Minkowski reconstruction together with the positive mass gap. Accordingly the earlier disclaimers remain correct for the continuum gauge and quantization papers taken in isolation, but they no longer apply to the completed series. $\square$

References

- [1] G. Duran Ballester, *Constructive Dynamical Axioms for Quantum Field Theory: Version VII: Core Axioms, Specialization Axes, Spectral Type, Scope Boundaries, and Detailed Framework Correspondences*, 2026.
- [2] G. Duran Ballester, *Convergence, Hypocoercivity, and Mean-Field Limit for the Kinetic Core of the Geometric Gas*, 2026.
- [3] G. Duran Ballester, *Local Continuum Yang–Mills Limit from Particle Trajectories*, 2026.
- [4] G. Duran Ballester, *From the Deterministic Regulator Data to a Gauge-Fixed Path Integral for the Finite $SU(3)$ Yang–Mills Lattice Theory*, 2026.
- [5] K. Osterwalder and R. Schrader, *Axioms for Euclidean Green’s functions*, Commun. Math. Phys. 31 (1973), 83–112; and *Axioms for Euclidean Green’s functions. II*, Commun. Math. Phys. 42 (1975), 281–305.