

Homework 1

Professor: Dr. Jaimes

Francisco Javier Vázquez Tavares A00827546

Contents

1	Problem 4.2	1
2	Problem 4.3	2
3	Problem 4.13	2
4	Problem 4.14	3
5	Problem 4.23	3

1 Problem 4.2

Use separation of variable in *cartesian* coordinates to solve the infinite *cubical* well (or particle in a box):

$$V(x, y, z) = \begin{cases} 0, & \forall x, y, z \in [0, a] \\ \infty, & \forall x, y, z \notin [0, a] \end{cases}$$

1. Find the stationary states, and the corresponding energies.
2. Call the distinct energies E_1, E_2, \dots in order of increasing energy. Find E_1, E_2, E_3, E_4, E_5 and E_6 . Determine their degeneracies (that is, the number of different states that share the same energy).
3. What is the degeneracy of E_{14} , and why is this case interesting?

Solution 1: Stationary states

To find the stationary states of the infinite cubical well, we are going to solve the time independent Schrödinger equation,

$$-\frac{\hbar^2}{2m} \nabla^2 \psi = E\psi, \quad \forall x, y, z \in [0, a],$$

with the following boundary conditions $\psi(0, 0, 0) = \psi(a, a, a) = 0$. To solve the equation we are going to use the method of separation of variables, that is, that we assume that the solution of the differential equation has the following form $\psi(x, y, z) = X(x)Y(y)Z(z)$. Substituting this solution to the differential equation, we can perform

some algebraic manipulation,

$$\begin{aligned}
 -\frac{\hbar^2}{2m}\nabla^2\psi &= E\psi \\
 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right)X(x)Y(y)Z(z) &= -\frac{2m}{\hbar^2}EX(x)Y(y)Z(z) \\
 Y(y)Z(z)\frac{\partial^2}{\partial x^2}X(x) + X(x)Z(z)\frac{\partial^2}{\partial y^2}Y(y) + X(x)Y(y)\frac{\partial^2}{\partial z^2}Z(z) &= -\frac{2m}{\hbar^2}EX(x)Y(y)Z(z) \\
 \frac{1}{X(x)}\frac{\partial^2}{\partial x^2}X(x) + \frac{1}{Y(y)}\frac{\partial^2}{\partial y^2}Y(y) + \frac{1}{Z(z)}\frac{\partial^2}{\partial z^2}Z(z) &= -\frac{2m}{\hbar^2}E
 \end{aligned}$$

2 Problem 4.3

Use

$$\begin{aligned}
 P_l^m(x) &\equiv (1-x^2)^{|m|/2} \left(\frac{d}{dx}\right)^{|m|} P_l(x) \\
 P_l(x) &\equiv \frac{1}{2^l l!} \left(\frac{d}{dx}\right)^l (x^2-1)^l \\
 Y_l^m(\theta, \phi) &= \epsilon \sqrt{\frac{(2l+1)}{4\pi} \frac{(l-|m|)!}{(l+|m|)!}} e^{im\phi} P_l^m(\cos[\theta])
 \end{aligned}$$

to construct Y_0^0 and Y_2^l . Check that they are normalized and orthogonal.

3 Problem 4.13

- Find $\langle r \rangle$ and $\langle r^2 \rangle$ for an electron in the ground state of hydrogen. Express your answers in terms of the Bohr radius.
- Find $\langle x \rangle$ and $\langle x^2 \rangle$ for an electron in the ground state of hydrogen. *Hint:* this requires no new integration-note that $r^2 = x^2 + y^2 + z^2$, and exploit the symmetry of the ground state.
- Find $\langle x^2 \rangle$ in the state $n=2, l=1, m=1$. *Warning:* This state is not symmetrical in x, y, z . Use $x = r \sin \theta \cos \phi$.

4 Problem 4.14

What is the *most probable* value of r , in the ground state of hydrogen? (The answer is not zero!) *Hint:* First you must figure out the probability that the electron would be found between r and $r + dr$.

5 Problem 4.23

In problem 4.3 you showed that

$$Y_2^l(\theta, \phi) = -\sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{i\phi}.$$

Apply the raising operator to find $Y_2^2(\theta, \phi)$. Use equation $A_l^m = \hbar\sqrt{l(l+1) - m(m \pm 1)} = \hbar\sqrt{(l \mp m)(l \pm m + 1)}$ to get the normalization.

[heading=bibintoc,title=References]