

Is Dementia predictable?

F. Di Filippo, E. Manfrin, E. Musiari, E. Palli 17 december 2021

Dataset 1/5

Dataset Dementia and Alzheimer longitudinal

| Subject.ID * | MRI.ID [‡] | Group | Visit [‡] | MR.Delay [‡] | M.F [‡] | Hand [‡] | Age ÷ | EDUC [‡] | SES ÷ | MMSE [‡] | CDR ÷ | eTIV ÷ | nWBV [‡] | ASF [‡] |
|--------------|---------------------|-------------|--------------------|-----------------------|------------------|-------------------|-------|-------------------|-------|-------------------|-------|--------|-------------------|------------------|
| OAS2_0001 | OAS2_0001_MR1 | Nondemented | 1 | 0 | М | R | 87 | 14 | 2 | 27 | 0.0 | 1987 | 0.696 | 0.883 |
| OAS2_0001 | OAS2_0001_MR2 | Nondemented | 2 | 457 | М | R | 88 | 14 | 2 | 30 | 0.0 | 2004 | 0.681 | 0.876 |
| OAS2_0002 | OAS2_0002_MR1 | Demented | 1 | 0 | М | R | 75 | 12 | NA | 23 | 0.5 | 1678 | 0.736 | 1.046 |
| OAS2_0002 | OAS2_0002_MR2 | Demented | 2 | 560 | М | R | 76 | 12 | NA | 28 | 0.5 | 1738 | 0.713 | 1.010 |
| OAS2_0002 | OAS2_0002_MR3 | Demented | 3 | 1895 | М | R | 80 | 12 | NA | 22 | 0.5 | 1698 | 0.701 | 1.034 |
| OAS2_0004 | OAS2_0004_MR1 | Nondemented | 1 | 0 | F | R | 88 | 18 | 3 | 28 | 0.0 | 1215 | 0.710 | 1.444 |
| OAS2_0004 | OAS2_0004_MR2 | Nondemented | 2 | 538 | F | R | 90 | 18 | 3 | 27 | 0.0 | 1200 | 0.718 | 1.462 |
| OAS2_0005 | OAS2_0005_MR1 | Nondemented | 1 | 0 | М | R | 80 | 12 | 4 | 28 | 0.0 | 1689 | 0.712 | 1.039 |
| OAS2_0005 | OAS2_0005_MR2 | Nondemented | 2 | 1010 | M | R | 83 | 12 | 4 | 29 | 0.5 | 1701 | 0.711 | 1.032 |

where SES is Socioeconomic Status, MMSE is Mini Mental State Examination, CDR is Clinical Dementia Rating, eTIV is Estimated Total Intracranial Volume, nWBV is Normalize Whole Brain Volume and ASF is Atlas Scaling Factor.

Source: Kaggle