



POLITECNICO
MILANO 1863


Coupled Markov chains with applications to Approximate Bayesian Computation for model based clustering

E. Bertoni, M. Caldarini, F. Di Filippo, G. Gabrielli, E. Musiani
10 January 2022


Introduction

A complex problem

1/30



Unbiased Markov chain
Monte Carlo methods with
couplings



likelihood




Approximate Bayesian
Computation

The complete method: MCMC + Couplings + ABC

The complete method: MCMC + Couplings + ABC Implementation

2/30



Metropolis Hastings with couplings and ABC

- ① Compute $s_{obs} = S(y_{obs})$;
- ② generate $\mu_x^{(0)} \sim \pi(\mu)$ and $\mu_y^{(0)} \sim \pi(\mu)$ from prior density;
- ③ generate $\sigma_x^{2(0)} \sim \pi(\sigma^2)$ and $\mu_y^{2(0)} \sim \pi(\sigma^2)$ from prior density;
- ④ generate with a maximal coupling two samples of N observations such that $y_{1i} \sim \mathcal{N}(\mu_x^{(0)}, \sigma_x^{2(0)})$ and $y_{2i} \sim \mathcal{N}(\mu_y^{(0)}, \sigma_y^{2(0)})$;
- ⑤ compute $s_x^{(0)} = S(y_1)$ and $s_y^{(0)} = S(y_2)$;
- ⑥ until $K_h(\|s_x^{(0)} - s_{obs}\|) > 0$:
 - ▶ generate $\mu_x^{(0)} \sim \pi(\mu)$ and $\sigma_x^{2(0)} \sim \pi(\sigma^2)$ from prior densities;
 - ▶ generate a sample of N observations such that $y_{1i} \sim \mathcal{N}(\mu_x^{(0)}, \sigma_x^{2(0)})$;
 - ▶ compute $s_x^{(0)} = S(y_1)$;
- ⑦ until $K_h(\|s_y^{(0)} - s_{obs}\|) > 0$:
 - ▶ generate $\mu_y^{(0)} \sim \pi(\mu)$ and $\sigma_y^{2(0)} \sim \pi(\sigma^2)$ from prior densities;
 - ▶ generate a sample of N observations such that $y_{1i} \sim \mathcal{N}(\mu_y^{(0)}, \sigma_y^{2(0)})$;
 - ▶ compute $s_x^{(0)} = S(y_1)$;

⑧ for $i = 1, \dots, N$: given

$$\theta^{(i-1)} = (\mu^{(i-1)}, \sigma^{2(i-1)})$$

- ▶ generate $[\theta_x^{(i)}, \theta_y^{(i)}]$ from a maximal coupling given $[\theta_x^{(i-1)}, \theta_y^{(i-1)}]$;
- ▶ generate from a maximal coupling two samples of N observations $y_1 \sim p(y|\theta_x^{(i)})$ and $y_2 \sim p(y|\theta_y^{(i)})$;
- ▶ compute $s_x^{(i)} = S(y_1)$ and $s_y^{(i)} = S(y_2)$;
- ▶ accept $\theta_x^{(i)}$ with probability

$$\frac{K_h(\|s_x^{(i)} - s_{obs}\|)\pi(\theta_x^{(i)})}{K_h(\|s_x^{(i-1)} - s_{obs}\|)\pi(\theta_x^{(i-1)})}$$

and accept $\theta_y^{(i)}$ with probability

$$\frac{K_h(\|s_y^{(i)} - s_{obs}\|)\pi(\theta_y^{(i)})}{K_h(\|s_y^{(i-1)} - s_{obs}\|)\pi(\theta_y^{(i-1)})}.$$

As output we get two sets of parameter vectors:

$$\mu_x^{(1)}, \dots, \mu_x^{(N)} \sim \pi_{ABC}(\mu | y_{obs});$$

$$\mu_y^{(1)}, \dots, \mu_y^{(N)} \sim \pi_{ABC}(\mu | y_{obs}).$$

$$\sigma_x^{2(1)}, \dots, \sigma_x^{2(N)} \sim \pi_{ABC}(\sigma^2 | y_{obs});$$

$$\sigma_y^{2(1)}, \dots, \sigma_y^{2(N)} \sim \pi_{ABC}(\sigma^2 | y_{obs}).$$

Set $p = \mathcal{N}(X_{t-1}, 0.1^2 * \mathbb{I})$ and $q = \mathcal{N}(Y_{t-1}, 0.1^2 * \mathbb{I})$, then:

- ① sample $X_t \sim p$;
- ② sample $W|X_t \sim \mathcal{U}\{[0, p(X_t)]\}$;
- ③ if $W \leq q(X_t)$ then output (X_t, X_t) , otherwise:
 - ① sample $Y_t \sim q$;
 - ② sample $W^*|Y_t \sim \mathcal{U}\{[0, q(Y_t)]\}$ until $W^* > p(Y_t)$ and output (X_t, Y_t) .

Summary statistic:

Sample mean, Sample Variance

Distance:

L^2 – norm of the difference of $S(y)$ and s_{obs}

Kernel:

$$K(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}, \quad K_h(u) = \frac{K(\frac{u}{h})}{h}$$

Model

$$Y_i | \mu, \sigma^2 \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\sigma^2 \sim \text{InvGa}(a, b)$$

$$\pi(\mu, \sigma) = \pi(\mu) * \pi(\sigma)$$

$$\mu_0 = 34, \quad \sigma_0^2 = 3$$

$$a = b = 1$$

Dataset


100 samples generated from a Gaussian distribution:

$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$


$$\mu_{obs} = 30, \sigma_{obs}^2 = 2$$

$$E[\mu|y] = 30, 19$$

Coupled chains




Sampling histogram with real distribution




$$E[\sigma^2|y] = 2,06$$

Coupled chains



Sampling histogram with real distribution



QUANTILE FUNCTION:

$$r \in (0, 1) \mapsto a + b(1 + 0.8 \left(\frac{1 - \exp(-g \cdot z(r))}{1 + \exp(-g \cdot z(r))} \right) (1 + z(r)^2)^k \cdot z(r)$$

where

$$z(r)$$

is the r -th quantile of the standard Normal distribution.

Model

Prior distribution: $a \sim \mathcal{U}([0, 10])$

$b \sim \mathcal{U}([0, 10])$

$g \sim \mathcal{U}([0, 10])$

$k \sim \mathcal{U}([0, 10])$

Summary statistic:

10 quantile + minimum

Distance:

L^2 – norm of the difference of $S(y)$ and s_{obs}

Kernel:

$$K(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}, \quad K_h(u) = \frac{K(\frac{u}{h})}{h}$$

Dataset

100 samples generated from the quantile distribution:

$$a_{obs} = 3, \quad b_{obs} = 1, \quad g_{obs} = 2, \quad k_{obs} = 0.5, \quad y_{obs}$$

$$y_{obs} = a_{obs} + b_{obs} \left(1 + 0.8 \left(\frac{1 - \exp(-g_{obs} \cdot z(r))}{1 + \exp(-g_{obs} \cdot z(r))} \right) (1 + z(r)^2)^k \cdot z(r) \right)$$

and

$$z(r) \stackrel{iid}{\sim} \mathcal{N}(0, 1)$$

ABC ALGORITHM

INITIALIZATION:

$$\theta^{(0)} = (a^{(0)}, b^{(0)}, g^{(0)}, k^{(0)})$$

$$\theta^{(0)} \sim \mathcal{U}([0, 10]^4)$$

for j in $(1, \dots, n)$:

- $z^{(0,j)}(r) \sim \mathcal{N}(0, 1)$

- $y^{(0,j)} \sim \text{quantilefunction}(z^{(0,j)}(r))$

compute the summaries statistics: $s^{(0)} = S(y^{(0)})$ Untill $Kh(\|s^{(0)} - s_{obs}\|) > 0$

- Generate $\theta^{(0)} \sim \mathcal{U}([0, 10]^4)$ from prior density.

- Generate $z^{(0)}(r) \sim \mathcal{N}(0, 1)$

- Generate a sample of 1000 observations such that
 $y \sim \text{quantilefunction}(z^{(0)}(r), \theta^{(0)})$

- Compute $s^{(0)} = S(y)$

for i in $(1, \dots, M)$:

① given $\theta^{(i-1)} = (a^{(i-1)}, b^{(i-1)}, g^{(i-1)}, k^{(i-1)})$

$\theta^{(i)} \sim \mathcal{N}(\theta^{(i-1)}, I)$

for j in $(1, \dots, n)$:

$z^{(i,j)}(r) \sim \mathcal{N}(0, 1)$

$y^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta^{(i-1)})$

② Compute the summaries $s^{(i)} = S(y^{(i)})$

③ Accept $\theta^{(i)}$ with probability $\frac{Kh(\|s^{(i)} - s_{obs}\|)\pi(\theta^{(i)})}{Kh(\|s^{(i-1)} - s_{obs}\|)\pi(\theta^{(i-1)})}$

parameters b, g, k are fixed equal to the observation parameters

$$a_1^{(0)} \sim \mathcal{U}([0, 10])$$

$$a_2^{(0)} \sim \mathcal{U}([0, 10])$$

for j in $(1, \dots, n_{obs})$:

- $z^{(0,j)}(r) \sim \mathcal{N}(0, 1)$

- $y_1^{(0,j)} \sim \text{quantilefunction}(z^{(0,j)}(r))$

- compute $s_1^{(0)} = S(y_1^{(0)})$

for j in $(1, \dots, n_{obs})$:

- $z^{(0,j)}(r) \sim \mathcal{N}(0, 1)$

- $y_2^{(0,j)} \sim \text{quantilefunction}(z^{(0,j)}(r))$

- compute the summaries statistics: $s_2^{(0)} = S(y_2^{(0)})$

Untill $Kh(\|s_1^{(0)} - s_{obs}\|) > 0$:

- 1 Generate $a_1^{(0)} \sim \mathcal{U}([0, 10])$ from prior density.
- 2 Generate $z_1^{(0)}(r) \sim \mathcal{N}(0, 1)$
- 3 Generate a sample of 1000 observations such that $y_1 \sim \text{quantilefunction}(z_1^{(0)}(r), a_1^{(0)})$
- 4 Compute $s_1^{(0)} = S(y_1)$

Untill $Kh(\|s_2^{(0)} - s_{obs}\|) > 0$:

- 1 Generate $a_2^{(0)} \sim \mathcal{U}([0, 10])$ from prior density.
- 2 Generate $z_2^{(0)}(r) \sim \mathcal{N}(0, 1)$
- 3 Generate a sample of 1000 observations such that $y_2 \sim \text{quantilefunction}(z_2^{(0)}(r), a_2^{(0)})$
- 4 Compute $s_2^{(0)} = S(y_2)$

ABC + COUPLED MARKOV CHAINS for parameter $a_{17/30}$

for i in $(1, \dots, M)$:

① generate $a_1^{(i)}, a_2^{(i)}$ from $\text{maximalcoupling}(a_1^{(i-1)}, a_2^{(i-1)})$

for j in $(1, \dots, n)$:

▶ $z^{(i,j)}(r) \sim \mathcal{N}(0, 1)$

▶ generate: $y_1^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta^{(i-1)})$
 $y_2^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta^{(i-1)})$

② Compute the summaries $s_1^{(i)} = S(y_1^{(i)})$ and $s_2^{(i)} = S(y_2^{(i)})$

③ acceptance:

▶ Accept $a_1^{(i)}$ with probability $\frac{Kh(\|s_1^{(i)} - s_{obs}\|)\pi(a_1^{(i)})}{Kh(\|s_1^{(i-1)} - s_{obs}\|)\pi(a_1^{(i-1)})}$ otherwise

$$a_1^{(i)} = a_1^{(i-1)}$$

▶ Accept $a_2^{(i)}$ with probability $\frac{Kh(\|s_2^{(i)} - s_{obs}\|)\pi(a_2^{(i)})}{Kh(\|s_2^{(i-1)} - s_{obs}\|)\pi(a_2^{(i-1)})}$ otherwise

$$a_2^{(i)} = a_2^{(i-1)}$$

given $(a_1^{(i-1)}, a_2^{(i-1)})$

set $p(\cdot | a_1^{(i-1)}) = \mathcal{N}(a_1^{(i-1)}, 1)$ and $q(\cdot | a_2^{(i-1)}) = \mathcal{N}(a_2^{(i-1)}, 1)$

① $a'_1 \sim p(\cdot | a_1^{(i-1)})$

② if $w_1 \sim \mathcal{U}((0, p(a' | a_1^{(i-1)}))) < q(a' | a_2^{(i-1)}) : a_1^{(i)} = a'_1$ and $a_2^{(i)} = a'_1$


otherwise until $w_2 \sim \mathcal{U}((0, q(a' | a_2^{(i-1)}))) > p(a' | a_1^{(i-1)})$:

① generate $a'_2 \sim q(\cdot | a_2^{(i-1)})$


② sample $I \sim \mathcal{U}((0, 1))$

and then set: $a_1^{(i)} = a'_1$ and $a_2^{(i)} = a'_2$


Coupled chains




Sampling histogram with real distribution




Coupled chains




Sampling histogram




Coupled chains




Sampling histogram



Coupled chains



Sampling histogram



given $\theta = (a, b, g, k)$

$$\theta_1^{(0)} \sim \mathcal{U}([0, 10]^4)$$

$$\theta_2^{(0)} \sim \mathcal{U}([0, 10]^4)$$

for j in $(1, \dots, n_{obs})$:

- $z^{(0,j)}(r) \sim \mathcal{N}(0, 1)$

- $y_1^{(0,j)} \sim \text{quantilefunction}(z^{(0,j)}(r), \theta_1^{(0)})$

- compute $s_1^{(0)} = S(y_1^{(0)})$

for j in $(1, \dots, n_{obs})$:

- $z^{(0,j)}(r) \sim \mathcal{N}(0, 1)$

- $y_2^{(0,j)} \sim \text{quantilefunction}(z^{(0,j)}(r), \theta_2^{(0)})$

- compute the summaries statistics: $s_2^{(0)} = S(y_2^{(0)})$

Until $Kh(\|s_1^{(0)} - s_{obs}\|) > 0$:

- 1 Generate $\theta_1^{(0)} \sim \mathcal{U}([0, 10]^4)$ from prior density.
- 2 Generate $z_1^{(0)}(r) \sim \mathcal{N}(0, 1)$
- 3 Generate a sample of 1000 observations such that $y_1 \sim \text{quantilefunction}(z_1^{(0)}(r), \theta_1^{(0)})$
- 4 Compute $s_1^{(0)} = S(y_1)$

Until $Kh(\|s_2^{(0)} - s_{obs}\|) > 0$:

- 1 Generate $\theta_2^{(0)} \sim \mathcal{U}([0, 10]^4)$ from prior density.
- 2 Generate $z_2^{(0)}(r) \sim \mathcal{N}(0, 1)$
- 3 Generate a sample of 1000 observations such that $y_2 \sim \text{quantilefunction}(z_2^{(0)}(r), \theta_2^{(0)})$
- 4 Compute $s_2^{(0)} = S(y_2)$

for i in $(1, \dots, M)$:

- ① generate $\theta_1^{(i)}, \theta_2^{(i)}$ from $\text{maximalcoupling}(\theta_1^{(i-1)}, \theta_2^{(i-1)})$

for j in $(1, \dots, n)$:

- ▶ $z^{(i,j)}(r) \sim \mathcal{N}(0, 1)$
- ▶ generate: $y_1^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta_1^{(i-1)})$
 $y_2^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta_2^{(i-1)})$

- ② Compute the summaries $s_1^{(i)} = S(y_1^{(i)})$ and $s_2^{(i)} = S(y_2^{(i)})$

- ③ acceptance:

- ▶ Accept $a_1^{(i)}$ with probability $\frac{Kh(\|s_1^{(i)} - s_{obs}\|)\pi(\theta_1^{(i)})}{Kh(\|s_1^{(i-1)} - s_{obs}\|)\pi(\theta_1^{(i-1)})}$ otherwise
 $a_1^{(i)} = \theta_1^{(i-1)}$
- ▶ Accept $a_2^{(i)}$ with probability $\frac{Kh(\|s_2^{(i)} - s_{obs}\|)\pi(\theta_2^{(i)})}{Kh(\|s_2^{(i-1)} - s_{obs}\|)\pi(\theta_2^{(i-1)})}$ otherwise
 $\theta_2^{(i)} = \theta_2^{(i-1)}$

given $(\theta_1^{(i-1)}, \theta_2^{(i-1)})$

set $p(\cdot|\theta_1^{(i-1)}) = \mathcal{N}(\theta_1^{(i-1)}, 0.1^2\mathbb{I})$ and $q(\cdot|\theta_2^{(i-1)}) = \mathcal{N}(\theta_2^{(i-1)}, 0.1^2\mathbb{I})$

① $\theta'_1 \sim p(\cdot|\theta_1^{(i-1)})$

② if $w_1 \sim \mathcal{U}((0, p(\theta'|\theta_1^{(i-1)}))) < q(\theta'|\theta_2^{(i-1)}) : \theta_1^{(i)} = \theta'_1$ and $\theta_2^{(i)} = \theta'_1$

otherwise until $w_2 \sim \mathcal{U}((0, q(\theta'|\theta_2^{(i-1)}))) > p(\theta'|\theta_1^{(i-1)})$:

■ generate $\theta'_2 \sim q(\cdot|\theta_2^{(i-1)})$

and then set: $\theta_1^{(i)} = \theta'_1$ and $\theta_2^{(i)} = \theta'_2$

