




POLITECNICO
MILANO 1863

Coupled Markov chains with applications to Approximate Bayesian Computation for model based clustering

E. Bertoni, M. Caldarini, F. Di Filippo, G. Gabrielli, E. Musiari
15 February 2022



likelihood


intractable



**Unbiased Markov chain
Monte Carlo methods with
couplings**



**Approximate Bayesian
Computation**




The complete method: MCMC + Couplings + ABC

The complete method: MCMC + Couplings + ABC

Implementation

2/14



Maximal coupling

Maximal coupling between two distributions p and q

- sample $X \sim p$;
- sample $W|X \sim \mathcal{U}\{[0, p(X)]\}$;
- if $W \leq q(X)$ then output (X, X) , otherwise:
 - ① sample $Y \sim q$;
 - ② sample $W^*|Y \sim \mathcal{U}\{[0, q(Y)]\}$ until $W^* > p(Y)$
 - ③ output (X, Y) .

Output

Distribution of a pair of random variables X, Y that maximizes $\mathbb{P}(X = Y)$ subject to the marginal constraints $X \sim p$ and $Y \sim q$

Metropolis Hastings with couplings and ABC

Model

$$Y_i | \mu, \sigma^2 \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\sigma^2 \sim \text{InvGa}(a, b)$$

$$\mu_0 = 8, \quad \sigma_0^2 = 4$$

$$a = b = 1$$

Metropolis Hastings with couplings and ABC

Initialization

set $\theta = (\mu, \sigma^2)$

for $k=1,2$:

until $K_h(||s_k^{(0)} - s_{obs}||) > 0$:

- sample $\theta_k^{(0)} \sim \pi(\theta)$
- simulate n_{obs} observations $y_{ki} \sim \mathcal{N}(\mu^{(0)}, \sigma^{2(0)})$ from maximal coupling
- compute $s_k^{(0)} = S(y_k)$;

Iterations

for $i = 1, \dots, M$:

given $\theta^{(i-1)} = (\mu^{(i-1)}, \sigma^{2(i-1)})$

- generate $[\theta_1^{(i)}, \theta_2^{(i)}]$ from a maximal coupling given $[\theta_1^{(i-1)}, \theta_2^{(i-1)}]$;
- for $k=1, 2$:
 - ① simulate n_{obs} observations $y_k^{(i)} \sim \mathcal{N}(\mu^{(i)}, \sigma^{2(i)})$ from maximal coupling;
 - ② compute $s_k^{(i)} = S(y_k)$;
 - ③ accept $\theta_k^{(i)}$ with probability

$$\frac{K_h(\|s_k^{(i)} - s_{obs}\|) \pi(\theta_k^{(i)})}{K_h(\|s_k^{(i-1)} - s_{obs}\|) \pi(\theta_k^{(i-1)})}$$

Output

for $k = 1, 2$

$$\mu_k^{(1)}, \dots, \mu_k^{(M)} \sim \pi_{ABC}(\mu | y_{obs});$$

$$\sigma_k^{2(1)}, \dots, \sigma_k^{2(M)} \sim \pi_{ABC}(\sigma | y_{obs}).$$

Summary statistics: Sample mean, Sample Variance

Distance: $L^2 - norm$

Kernel: $K(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}$, $K_h(u) = \frac{K(\frac{u}{h})}{h}$

Proposal distribution: $\begin{bmatrix} \mu' \\ \log(\sigma^{2'}) \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu^{(i)} \\ \log(\sigma^{2(i)}) \end{bmatrix}, 0.1^2 \cdot \mathcal{I} \right)$

Summary statistics: Sample mean, Sample Variance

Distance: $L^2 - norm$

Kernel: $K(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}$, $K_h(u) = \frac{K(\frac{u}{h})}{h}$

Proposal distribution: $\begin{bmatrix} \mu' \\ \log(\sigma'^2) \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu^{(i)} \\ \log(\sigma^{2(i)}) \end{bmatrix}, 0.1^2 \cdot \mathcal{I} \right)$


Dataset

100 samples generated from a Gaussian distribution:


$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$

$$\mu_{obs} = 10, \sigma_{obs}^2 = 3$$

Coupled chains from one processor



Sampling histogram with estimated distribution




Samplings from all accepted chains




Time Averaged Estimators mean obtained: $\mathbb{E}[H_{k:m}(X_\mu, Y_\mu)] \simeq 9.8$

Coupled chains from one processor



Sampling histogram with estimated distribution



Samplings from all accepted chains



Time Averaged Estimators mean obtained: $\mathbb{E}[H_{k:m}(X_{\sigma^2}, Y_{\sigma^2})] \simeq 2.56$

Numerical experiment: g-and-k distribution

Quantile function

$$r \in (0, 1) \mapsto a + b \left(1 + 0.8 \left(\frac{1 - \exp(-g \cdot z(r))}{1 + \exp(-g \cdot z(r))} \right) \right) (1 + z(r)^2)^k \cdot z(r)$$

where $z(r)$ is the r -th quantile of the standard Normal distribution.

Prior distributions

$$a \sim \mathcal{U}([0, 10])$$

$$b \sim \mathcal{U}([0, 10])$$

$$g \sim \mathcal{U}([0, 10])$$

$$k \sim \mathcal{U}([0, 10])$$

Summary statistics: 10 quantiles

Distance: L^2 – norm

Kernel:

$$K(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}, \quad K_h(u) = \frac{K(\frac{u}{h})}{h}$$

Dataset


100 samples generated as follows:

$$a_{obs} = 3, b_{obs} = 1, g_{obs} = 2, k_{obs} = 0.5$$

$$y_{obs} \sim \text{quantilefunction}(z(r), \theta_{obs})$$

$$z(r) \sim \mathcal{N}(0, 1)$$

Quantile distribution: y_{obs}



Initialization

given $\theta = (a, b, g, k)$

for $k=1,2$:

Untill $K_h(\|s_k^{(0)} - s_{obs}\|) > 0$:

- Generate $\theta_k^{(0)} \sim \mathcal{U}([0, 10]^4)$ from prior density.
- Generate $z(r) \sim \mathcal{N}(0, 1)$
- Generate a sample of n_{obs} observations such that $y_k \sim \text{quantilefunction}(z(r), \theta_k^0)$
- Compute $s_k^{(0)} = S(y_k)$

Iterations

for i in $(1, \dots, M)$:

- generate $\theta_1^{(i)}, \theta_2^{(i)}$ from $\text{maximalcoupling}(\theta_1^{(i-1)}, \theta_2^{(i-1)})$
- simulate n_{obs} observations $y_1^{(i)}, y_2^{(i)}$ from maximal coupling
- Compute the summaries $s_k^{(i)} = S(y_k^{(i)})$
- Accept $\theta_k^{(i)}$ with probability

$$\frac{Kh(\|s_k^{(i)} - s_{obs}\|)\pi(\theta_k^{(i)})}{Kh(\|s_k^{(i-1)} - s_{obs}\|)\pi(\theta_k^{(i-1)})}$$

otherwise $\theta_k^{(i)} = \theta_k^{(i-1)}$;

Iterations

for i in $(1, \dots, M)$

Couplings

for j in $(1, \dots, n_{obs})$:

- ① $z^{(i,j)}(r) \sim \mathcal{N}(0, 1)$

- ② generate:


$$y_1^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta_1^{(i-1)})$$

$$y_2^{(ij)} \sim \text{quantilefunction}(z^{(i,j)}(r), \theta_2^{(i-1)})$$


$$\kappa n(\|S_k^{(i)} - S_{obs}\|) \pi(\theta_k^{(i)})$$

otherwise $\theta_k^{(i)} = \theta_k^{(i-1)}$;

Coupled chains from one core



Sampling histogram with estimated distribution



Samplings from all accepted chains

