ETH Zürich
Department of Physics

# The Berezinskii-Kosterlitz-Thouless Transition in the 2D XY Model

**Lara Turgut** **Francesco Conoscenti** **Ben Bullinger**

Computational Statistical Physics (402-0812-00L), Spring 2025

# Outline

1. Introduction
2. Theoretical Background
3. Monte Carlo Methods
4. Implementation Details
5. Results and Analysis
6. Conclusion

# Outline

1. Introduction
2. Theoretical Background
3. Monte Carlo Methods
4. Implementation Details
5. Results and Analysis
6. Conclusion

# The 2D XY Model

- XY model: classical spins on a 2D lattice, constrained to a plane with $O(2)$ symmetry.

$$H = -J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j = -J \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) \tag{1}$$

  [1]

- Each spin is a unit vector $\mathbf{S}_i = (\cos\theta_i, \sin\theta_i)$. $J > 0$ favors alignment.
- Control parameter: Temperature $T$. We study behavior as $T$ varies.

# The 2D XY Model

- XY model: classical spins on a 2D lattice, constrained to a plane with $O(2)$ symmetry.

$$H = -J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j = -J \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) \tag{1}$$

  [1]

- Each spin is a unit vector $\mathbf{S}_i = (\cos\theta_i, \sin\theta_i)$. $J > 0$ favors alignment.
- Control parameter: Temperature $T$. We study behavior as $T$ varies.
- Unlike conventional phase transitions with symmetry breaking:
  - **Mermin-Wagner theorem:** No spontaneous breaking of continuous symmetries (hence no true long-range order) at $T > 0$ in $d \leq 2$.
  - Yet, the 2D XY model *does* exhibit a phase transition!

# The 2D XY Model

- XY model: classical spins on a 2D lattice, constrained to a plane with $O(2)$ symmetry.

$$H = -J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j = -J \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) \tag{1}$$

  [1]

- Each spin is a unit vector $\mathbf{S}_i = (\cos\theta_i, \sin\theta_i)$. $J > 0$ favors alignment.
- Control parameter: Temperature $T$. We study behavior as $T$ varies.
- Unlike conventional phase transitions with symmetry breaking:
  - **Mermin-Wagner theorem:** No spontaneous breaking of continuous symmetries (hence no true long-range order) at $T > 0$ in $d \leq 2$.
  - Yet, the 2D XY model *does* exhibit a phase transition!
- The Berezinskii-Kosterlitz-Thouless (BKT) transition:
  - Topological phase transition with no conventional order parameter.
  - Transition from quasi-long-range order (QLRO) to exponential disorder.
  - Driven by the unbinding of vortex-antivortex pairs.

# Physical Significance

- The BKT transition represents a unique phase transition mechanism
  - **Infinite-order transition:** All derivatives of free energy are continuous
  - **Topological in nature:** Driven by topological defects (vortices)

# Physical Significance

- The BKT transition represents a unique phase transition mechanism
  - **Infinite-order transition:** All derivatives of free energy are continuous
  - **Topological in nature:** Driven by topological defects (vortices)
- Found in diverse physical systems:
  - Thin films of superfluid helium-4
  - 2D superconductors
  - 2D melting (KTHNY theory)
  - Josephson junction arrays
  - Cold atom systems
  - Quantum spin systems

# Physical Significance

- The BKT transition represents a unique phase transition mechanism
  - **Infinite-order transition:** All derivatives of free energy are continuous
  - **Topological in nature:** Driven by topological defects (vortices)
- Found in diverse physical systems:
  - Thin films of superfluid helium-4
  - 2D superconductors
  - 2D melting (KTHNY theory)
  - Josephson junction arrays
  - Cold atom systems
  - Quantum spin systems
- Universal mechanism: A beautiful example of universality in critical phenomena

# Outline

1. Introduction

2. Theoretical Background

3. Monte Carlo Methods

4. Implementation Details

5. Results and Analysis

6. Conclusion

# Algebraic Order in the 2D XY Model

- Spin-wave approximation (valid at low temperature):

$$H \approx E_0 + \frac{J}{2} \int d^2 r (\nabla \theta(\mathbf{r}))^2 \tag{2}$$

# Algebraic Order in the 2D XY Model

- Spin-wave approximation (valid at low temperature):

$$H \approx E_0 + \frac{J}{2} \int d^2 r (\nabla \theta(\mathbf{r}))^2 \tag{2}$$

- Correlation function analysis within this approximation yields:

$$g(r) = \langle e^{i(\theta(\mathbf{r}) - \theta(0))} \rangle \approx \left(\frac{r}{a}\right)^{-\eta(T)} \quad \text{with} \quad \eta(T) = \frac{k_B T}{2\pi J} \tag{3}$$

- Key property: Algebraic decay of correlations at low temperature.
  - **Low-T phase (QLRO):** $g(r) \sim r^{-\eta(T)}$ with temperature-dependent exponent.
  - **High-T phase (Disordered):** $g(r) \sim e^{-r/\xi(T)}$ (exponential decay).

# Algebraic Order in the 2D XY Model

- Spin-wave approximation (valid at low temperature):

$$H \approx E_0 + \frac{J}{2} \int d^2 r (\nabla \theta(\mathbf{r}))^2 \tag{2}$$

- Correlation function analysis within this approximation yields:

$$g(r) = \langle e^{i(\theta(\mathbf{r}) - \theta(0))} \rangle \approx \left(\frac{r}{a}\right)^{-\eta(T)} \quad \text{with} \quad \eta(T) = \frac{k_B T}{2\pi J} \tag{3}$$

- Key property: Algebraic decay of correlations at low temperature.
  - **Low-T phase (QLRO):** $g(r) \sim r^{-\eta(T)}$ with temperature-dependent exponent.
  - **High-T phase (Disordered):** $g(r) \sim e^{-r/\xi(T)}$ (exponential decay).
- This algebraic decay = **Quasi-Long-Range Order (QLRO)**.
  - Infinite correlation length $\xi = \infty$ in the low-T phase.
  - System is critical at all temperatures below $T_{KT}$.

# The Role of Vortices

- Vortices: topological defects with winding of $2\pi n$ in phase field

$$\oint \nabla\theta \cdot d\mathbf{l} = 2\pi n \tag{4}$$

- Energy of a single vortex (with $n = 1$):

$$\Delta E = \pi J \ln \frac{L}{a} \tag{5}$$

  where $L$ is system size and $a$ is lattice spacing

# The Role of Vortices

- Vortices: topological defects with winding of $2\pi n$ in phase field

$$\oint \nabla\theta \cdot d\mathbf{l} = 2\pi n \tag{4}$$

- Energy of a single vortex (with $n = 1$):

$$\Delta E = \pi J \ln \frac{L}{a} \tag{5}$$

  where $L$ is system size and $a$ is lattice spacing
- Entropy of a single vortex:

$$\Delta S = k_B \ln \frac{L^2}{a^2} = 2k_B \ln \frac{L}{a} \tag{6}$$

# The Role of Vortices

- Vortices: topological defects with winding of $2\pi n$ in phase field

$$\oint \nabla\theta \cdot d\mathbf{l} = 2\pi n \tag{4}$$

- Energy of a single vortex (with $n = 1$):

$$\Delta E = \pi J \ln \frac{L}{a} \tag{5}$$

  where $L$ is system size and $a$ is lattice spacing
- Entropy of a single vortex:

$$\Delta S = k_B \ln \frac{L^2}{a^2} = 2k_B \ln \frac{L}{a} \tag{6}$$

- Free energy:

$$\Delta F = \Delta E - T\Delta S = (\pi J - 2k_B T) \ln \frac{L}{a} \tag{7}$$

# The Role of Vortices

- Vortices: topological defects with winding of $2\pi n$ in phase field

$$\oint \nabla\theta \cdot d\mathbf{l} = 2\pi n \tag{4}$$

- Energy of a single vortex (with $n = 1$):

$$\Delta E = \pi J \ln \frac{L}{a} \tag{5}$$

  where $L$ is system size and $a$ is lattice spacing
- Entropy of a single vortex:

$$\Delta S = k_B \ln \frac{L^2}{a^2} = 2k_B \ln \frac{L}{a} \tag{6}$$

- Free energy:

$$\Delta F = \Delta E - T\Delta S = (\pi J - 2k_B T) \ln \frac{L}{a} \tag{7}$$

- Crucial insight: Vortices are bound in pairs below critical temperature, and unbound above it
  - For $T < \frac{\pi J}{2k_B}$: $\Delta F > 0$ (bound vortex pairs)
  - For $T > \frac{\pi J}{2k_B}$: $\Delta F < 0$ (free vortices proliferate)

# Visualizing Spin Configurations and Vortices




$T = 0.85 \lesssim T_{KT}$
(QLRO, Approaching Transition)




$T \approx 0.92 \gtrsim T_{KT}$
(Pairs Unbinding)




$T = 1.20 > T_{KT}$
(Disordered, Free Vortices)

Spin configurations for L=20 showing the transition from quasi-long-range order with bound vortex pairs (left) to a disordered state with proliferating free vortices (right). Red/Blue circles indicate vortex/antivortex locations.

# Renormalization Group Analysis

- Mapping to 2D Coulomb gas: vortices as charges with $\ln(r)$ interaction.
- Kosterlitz RG flow equations for effective coupling $K(l) = J_{\text{eff}}(l)/(k_B T)$ and vortex fugacity $y(l)$ at length scale $l$:

$$\frac{dK^{-1}(l)}{dl} = 4\pi^3 y(l)^2 + O(y(l)^4) \tag{8}$$

$$\frac{dy(l)}{dl} = (2 - \pi K(l))y(l) + O(y(l)^3) \tag{9}$$

# Renormalization Group Analysis

- Mapping to 2D Coulomb gas: vortices as charges with $\ln(r)$ interaction.
- Kosterlitz RG flow equations for effective coupling $K(l) = J_{\text{eff}}(l)/(k_B T)$ and vortex fugacity $y(l)$ at length scale $l$:

$$\frac{dK^{-1}(l)}{dl} = 4\pi^3 y(l)^2 + O(y(l)^4) \tag{8}$$

$$\frac{dy(l)}{dl} = (2 - \pi K(l))y(l) + O(y(l)^3) \tag{9}$$

- Flow diagram shows two phases separated by a line of fixed points:
  - **Low-T** ($K > 2/\pi$): $y \to 0$ (vortices are irrelevant, QLRO prevails).
  - **High-T** ($K < 2/\pi$): $y$ increases (vortices proliferate, destroying QLRO).

# Renormalization Group Analysis

- Mapping to 2D Coulomb gas: vortices as charges with $\ln(r)$ interaction.
- Kosterlitz RG flow equations for effective coupling $K(l) = J_{\text{eff}}(l)/(k_B T)$ and vortex fugacity $y(l)$ at length scale $l$:

$$\frac{dK^{-1}(l)}{dl} = 4\pi^3 y(l)^2 + O(y(l)^4) \tag{8}$$

$$\frac{dy(l)}{dl} = (2 - \pi K(l))y(l) + O(y(l)^3) \tag{9}$$

- Flow diagram shows two phases separated by a line of fixed points:
  - **Low-T** ($K > 2/\pi$): $y \to 0$ (vortices are irrelevant, QLRO prevails).
  - **High-T** ($K < 2/\pi$): $y$ increases (vortices proliferate, destroying QLRO).
- Universal predictions:
  - Critical coupling: $K_c = 2/\pi \implies T_{KT} = \frac{\pi J}{2k_B}$.
  - **Nelson-Kosterlitz jump**: Spin stiffness $\rho_s$ (helicity modulus) jumps discontinuously at the transition:

$$\rho_s(T_{KT}^-) = \frac{2k_B T_{KT}}{\pi} \tag{10}$$

  - Correlation length above transition: $\xi(T) \sim \exp\left(b/\sqrt{T/T_{KT} - 1}\right)$

# Outline

1. Introduction

2. Theoretical Background

3. Monte Carlo Methods

4. Implementation Details

5. Results and Analysis

6. Conclusion

# Simulating the XY Model

- Challenges in simulating BKT transition:
  - Critical slowing down near phase transition
  - Need for large system sizes (finite-size effects)
  - Long correlation times

# Simulating the XY Model

- Challenges in simulating BKT transition:
  - Critical slowing down near phase transition
  - Need for large system sizes (finite-size effects)
  - Long correlation times
- Standard Metropolis algorithm:
  - Energy change for spin flip: $\Delta E = -J \sum_{\langle i,j \rangle} [\cos(\theta_i' - \theta_j) - \cos(\theta_i - \theta_j)]$
  - Accept new state with probability $P = \min(1, e^{-\beta \Delta E})$
  - Very inefficient near critical point due to large correlated regions

# Simulating the XY Model

- Challenges in simulating BKT transition:
  - Critical slowing down near phase transition
  - Need for large system sizes (finite-size effects)
  - Long correlation times
- Standard Metropolis algorithm:
  - Energy change for spin flip: $\Delta E = -J \sum_{\langle i,j \rangle} [\cos(\theta_i' - \theta_j) - \cos(\theta_i - \theta_j)]$
  - Accept new state with probability $P = \min(1, e^{-\beta \Delta E})$
  - Very inefficient near critical point due to large correlated regions
- More efficient approaches:
  - **Wolff cluster algorithm (Implemented):** Updates entire clusters of spins, significantly reducing critical slowing down.
  - *Parallel tempering (Discussed):* Exchanges configurations between different temperatures to improve sampling (beneficial but not used in our final simulation code).

# Wolff Cluster Algorithm

- Efficiently updates entire clusters of correlated spins
  1. Choose a random reflection axis (unit vector $\mathbf{r}$)
  2. Pick a random initial site for the cluster
  3. Add neighboring spins with probability $P_{\text{add}} = 1 - \exp[\min(0, 2\beta J(\mathbf{S}_i \cdot \mathbf{r})(\mathbf{S}_j \cdot \mathbf{r}))]$
  4. Reflect all spins in the cluster w.r.t. the chosen axis

# Wolff Cluster Algorithm

- Efficiently updates entire clusters of correlated spins
  1. Choose a random reflection axis (unit vector $\mathbf{r}$)
  2. Pick a random initial site for the cluster
  3. Add neighboring spins with probability $P_{\text{add}} = 1 - \exp[\min(0, 2\beta J(\mathbf{S}_i \cdot \mathbf{r})(\mathbf{S}_j \cdot \mathbf{r}))]$
  4. Reflect all spins in the cluster w.r.t. the chosen axis
- Advantages:
  - Dramatically reduces critical slowing down
  - Dynamical critical exponent $z \approx 0$ (vs. $z \approx 2$ for Metropolis)
  - Efficiently samples low-energy configurations

# Wolff Cluster Algorithm

- Efficiently updates entire clusters of correlated spins
  1. Choose a random reflection axis (unit vector $\mathbf{r}$)
  2. Pick a random initial site for the cluster
  3. Add neighboring spins with probability $P_{\text{add}} = 1 - \exp[\min(0, 2\beta J(\mathbf{S}_i \cdot \mathbf{r})(\mathbf{S}_j \cdot \mathbf{r}))]$
  4. Reflect all spins in the cluster w.r.t. the chosen axis
- Advantages:
  - Dramatically reduces critical slowing down
  - Dynamical critical exponent $z \approx 0$ (vs. $z \approx 2$ for Metropolis)
  - Efficiently samples low-energy configurations
- Our implementation (‘simulate.py‘):
  - Uses the Wolff algorithm (or Metropolis as an option).
  - Leverages Numba JIT compilation for performance.
  - Runs simulations for different temperatures in parallel using multiprocessing.
  - *Note: Does not implement Parallel Tempering (Replica Exchange).*

# Parallel Tempering

- *(Method discussed for context, not implemented in 'simulate.py' results)*
- Simulates multiple replicas at different temperatures $T_1 < T_2 < ... < T_M$
- Periodically proposes exchanges between adjacent temperatures
  - Accept exchange with probability $P = \min(1, \exp((\beta_i - \beta_j)(E_j - E_i)))$

# Parallel Tempering

- *(Method discussed for context, not implemented in 'simulate.py' results)*
- Simulates multiple replicas at different temperatures $T_1 < T_2 < ... < T_M$
- Periodically proposes exchanges between adjacent temperatures
  - Accept exchange with probability $P = \min(1, \exp((\beta_i - \beta_j)(E_j - E_i)))$
- Benefits:
  - Helps systems escape from local minima
  - Improves sampling at low temperatures
  - Configurations can diffuse across temperature space

# Parallel Tempering

- *(Method discussed for context, not implemented in 'simulate.py' results)*
- Simulates multiple replicas at different temperatures $T_1 < T_2 < ... < T_M$
- Periodically proposes exchanges between adjacent temperatures
  - Accept exchange with probability $P = \min(1, \exp((\beta_i - \beta_j)(E_j - E_i)))$
- Benefits:
  - Helps systems escape from local minima
  - Improves sampling at low temperatures
  - Configurations can diffuse across temperature space
- Implementation details (if used):
  - Alternating even/odd exchanges for better mixing
  - Temperature spacing optimized for exchange acceptance rates
  - Often combined with cluster algorithms like Wolff for optimal efficiency

# Outline

1. Introduction

2. Theoretical Background

3. Monte Carlo Methods

4. Implementation Details

5. Results and Analysis

6. Conclusion

# Our Implementation Approach

- Simulation structure:
  1. **Initial Equilibration**: Run sufficient updates to forget initial state.
  2. **Thermalization**: Ensure equilibrium is reached at each target temperature (often combined with equilibration).
  3. **Measurement**: Collect statistics over long runs in equilibrium.

# Our Implementation Approach

- Simulation structure:
  1. **Initial Equilibration**: Run sufficient updates to forget initial state.
  2. **Thermalization**: Ensure equilibrium is reached at each target temperature (often combined with equilibration).
  3. **Measurement**: Collect statistics over long runs in equilibrium.
- Optimization techniques:
  - Numba ‘@njit‘ compilation for core simulation loops (Wolff update, energy calculation).
  - Vectorized operations with NumPy where possible.
  - Pre-computation of neighbor indices.
  - (Optional: Dynamic adjustment of updates per sweep based on avg. cluster size / acceptance rates).

# Our Implementation Approach

- Simulation structure:
  1. **Initial Equilibration**: Run sufficient updates to forget initial state.
  2. **Thermalization**: Ensure equilibrium is reached at each target temperature (often combined with equilibration).
  3. **Measurement**: Collect statistics over long runs in equilibrium.
- Optimization techniques:
  - Numba ‘@njit‘ compilation for core simulation loops (Wolff update, energy calculation).
  - Vectorized operations with NumPy where possible.
  - Pre-computation of neighbor indices.
  - (Optional: Dynamic adjustment of updates per sweep based on avg. cluster size / acceptance rates).
- Simulation workflow:
  - Define temperature range and system sizes $L$.
  - Run independent simulations for each $(L, T)$ using Wolff algorithm (and potentially Parallel Tempering).
  - Thermalize, then collect measurements (Energy, Stiffness, etc.) over many sweeps.
  - Calculate averages and estimate errors (e.g., via Jackknife).
  - Perform finite-size scaling analysis to extract $T_{KT}$ and critical exponents.

# Key Observables

- Physical quantities measured in our simulation:
  - **Energy per spin**: $e = \frac{\langle H \rangle}{N} = -\frac{J}{N} \sum_{\langle i,j \rangle} \langle \cos(\theta_i - \theta_j) \rangle$
  - **Specific heat**: $c_v = \frac{\langle E^2 \rangle - \langle E \rangle^2}{NT^2}$
  - **Magnetization**: $\mathbf{m} = \frac{1}{N} \left\langle \sum_i (\cos\theta_i, \sin\theta_i) \right\rangle$
  - **Susceptibility**: $\chi = \frac{\langle M^2 \rangle - \langle M \rangle^2}{NT}$

# Key Observables

- Physical quantities measured in our simulation:
  - **Energy per spin**: $e = \frac{\langle H \rangle}{N} = -\frac{J}{N} \sum_{\langle i,j \rangle} \langle \cos(\theta_i - \theta_j) \rangle$
  - **Specific heat**: $c_v = \frac{\langle E^2 \rangle - \langle E \rangle^2}{NT^2}$
  - **Magnetization**: $\mathbf{m} = \frac{1}{N} \left\langle \sum_i (\cos\theta_i, \sin\theta_i) \right\rangle$
  - **Susceptibility**: $\chi = \frac{\langle M^2 \rangle - \langle M \rangle^2}{NT}$
  - **Spin stiffness (Helicity modulus)**:

$$\rho_s = \langle H_x \rangle - \frac{\beta}{N} \langle I_x^2 \rangle \tag{11}$$

where $H_x = \sum_{\langle i,j \rangle_x} \cos(\theta_i - \theta_j)$ and $I_x = \sum_{\langle i,j \rangle_x} \sin(\theta_i - \theta_j)$

# Key Observables

- Physical quantities measured in our simulation:
  - **Energy per spin**: $e = \frac{\langle H \rangle}{N} = -\frac{J}{N} \sum_{\langle i,j \rangle} \langle \cos(\theta_i - \theta_j) \rangle$
  - **Specific heat**: $c_v = \frac{\langle E^2 \rangle - \langle E \rangle^2}{NT^2}$
  - **Magnetization**: $\mathbf{m} = \frac{1}{N} \left\langle \sum_i (\cos\theta_i, \sin\theta_i) \right\rangle$
  - **Susceptibility**: $\chi = \frac{\langle M^2 \rangle - \langle M \rangle^2}{NT}$
  - **Spin stiffness (Helicity modulus)**:

$$\rho_s = \langle H_x \rangle - \frac{\beta}{N} \langle I_x^2 \rangle \tag{11}$$

    where $H_x = \sum_{\langle i,j \rangle_x} \cos(\theta_i - \theta_j)$ and $I_x = \sum_{\langle i,j \rangle_x} \sin(\theta_i - \theta_j)$
  - **Vortex density**: $\omega_v = \frac{1}{N} \sum_{\text{plaquettes}} \left| \frac{1}{2\pi} \sum_{\text{edges}} \Delta\theta_{ij} \right|$
- Spin stiffness is crucial for locating $T_{KT}$ through the universal jump

# Error Analysis

- Statistical errors from Monte Carlo sampling:
  - Sequential MC data is *correlated*
  - Standard error formulas for independent data are inadequate

# Error Analysis

- Statistical errors from Monte Carlo sampling:
  - Sequential MC data is *correlated*
  - Standard error formulas for independent data are inadequate
- Jackknife resampling for reliable error estimation:
  1. Divide data into $B$ blocks larger than correlation time
  2. For each block $b$, compute estimate $\bar{O}_b$ by excluding block $b$
  3. Jackknife error: $\sigma^2_{\text{JK}} = \frac{B-1}{B} \sum_b (\bar{O}_b - \bar{O})^2$

# Error Analysis

- Statistical errors from Monte Carlo sampling:
  - Sequential MC data is *correlated*
  - Standard error formulas for independent data are inadequate
- Jackknife resampling for reliable error estimation:
  1. Divide data into $B$ blocks larger than correlation time
  2. For each block $b$, compute estimate $\bar{O}_b$ by excluding block $b$
  3. Jackknife error: $\sigma^2_{\text{JK}} = \frac{B-1}{B} \sum_b (\bar{O}_b - \bar{O})^2$
- Benefits:
  - Correctly accounts for autocorrelations in MC data
  - Works well for derived quantities (non-linear functions)
  - More reliable than naive variance calculations
  - Essential for accurate error bars on quantities near transition

# Outline

1. Introduction

2. Theoretical Background

3. Monte Carlo Methods

4. Implementation Details

5. Results and Analysis

6. Conclusion

# Energy and Specific Heat

**Internal Energy per Spin ($e$)**

- Smoothly decreases with temperature.
- Weak dependence on system size $L$.
- No sharp features indicating the transition.

**Specific Heat ($c_v$)**

- Exhibits a broad peak around $T \approx 1.0J$.
- Peak height increases slightly with $L$.
- Crucially, no divergence at $T_{KT}$, consistent with infinite-order BKT transition.






# Magnetization and Susceptibility

## Magnetization ($|\mathbf{m}|$)

- Suppressed at finite $T$ for finite $L$, consistent with Mermin-Wagner.
- Slow decay with $L$ in the low-T phase, compatible with algebraic decay ($|\mathbf{m}| \sim L^{-\eta(T)/2}$) expected for QLRO.
- Faster decay in the high-T phase.

## Susceptibility ($\chi$)

- Develops a peak near the transition region.
- Peak height grows significantly with $L$.
- Diverges as $L \to \infty$ for $T \leq T_{KT}$.
- Provides an alternative route to estimate $T_{KT}$, though less precise than stiffness.






# Spin Stiffness (Helicity Modulus $\rho_s$)

## Identifying the BKT Transition

- Spin stiffness measures the free energy cost of applying a small twist to the boundary conditions.
- Decreases monotonically with temperature.
- Key indicator: Universal jump prediction by Nelson-Kosterlitz:

$$\rho_s(T_{KT}^-) = \frac{2k_B T_{KT}}{\pi}$$

- Finite-size systems show smooth curves crossing the universal line $y = 2k_B T/\pi$.
- Intersection of $\rho_s(T, L)$ with this line provides a size-dependent estimate $T_{KT}(L)$.
- The convergence point of these intersections for $L \to \infty$ yields $T_{KT}(\infty)$.
- Our data suggests $T_{KT} \approx 0.89 J/k_B$.




Plot shows $\rho_s$ vs $T$ for different $L$. The dashed line is $y = 2k_B T/\pi$.

# Vortex Density

## Role of Topological Defects

- Vortex density $\omega_v$ measures the concentration of topological defects.
- At low temperatures ($T \ll T_{KT}$):
  - Vortices exist only in tightly bound vortex-antivortex pairs.
  - Measured density is very low.
- Near $T_{KT}$:
  - Pairs start to unbind.
  - Density increases sharply.
- Above $T_{KT}$:
  - Proliferation of free vortices and antivortices.
  - Density becomes significant, screening interactions and destroying QLRO.
- Provides a microscopic view of the transition mechanism.




Plot shows average vortex density vs $T$ for different $L$.

# Finite-Size Scaling of $T_{KT}(L)$

**Extracting the Transition Temperature**

- Estimate $T_{KT}(L)$ for each size $L$ from the intersection $\rho_s(T, L) = 2k_B T/\pi$.
- BKT theory predicts the scaling towards the thermodynamic limit ($L \to \infty$):

$$T_{KT}(L) = T_{KT}(\infty) + \frac{c}{(\ln(L/L_0))^2}$$

where $c$ and $L_0$ are non-universal constants.

# Finite-Size Scaling of $T_{KT}(L)$

**Extracting the Transition Temperature**

- Estimate $T_{KT}(L)$ for each size $L$ from the intersection $\rho_s(T, L) = 2k_B T/\pi$.
- BKT theory predicts the scaling towards the thermodynamic limit ($L \to \infty$):

$$T_{KT}(L) = T_{KT}(\infty) + \frac{c}{(\ln(L/L_0))^2}$$

  where $c$ and $L_0$ are non-universal constants.
- Plotting $T_{KT}(L)$ vs $1/(\ln L)^2$ should yield a linear relationship for large $L$.
- The y-intercept gives the estimate for $T_{KT}(\infty)$.

# Finite-Size Scaling of $T_{KT}(L)$

## Extracting the Transition Temperature

- Estimate $T_{KT}(L)$ for each size $L$ from the intersection $\rho_s(T, L) = 2k_B T/\pi$.
- BKT theory predicts the scaling towards the thermodynamic limit ($L \to \infty$):

$$T_{KT}(L) = T_{KT}(\infty) + \frac{c}{(\ln(L/L_0))^2}$$

  where $c$ and $L_0$ are non-universal constants.
- Plotting $T_{KT}(L)$ vs $1/(\ln L)^2$ should yield a linear relationship for large $L$.
- The y-intercept gives the estimate for $T_{KT}(\infty)$.
- Our linear fit yields:

$$T_{KT}(\infty) \approx 0.893(4) J/k_B$$

- Result is consistent with established literature values (e.g., Ref. [1] and others cite $T_{KT} \approx 0.89289 J/k_B$).




Plot shows $T_{KT}(L)$ extracted from stiffness data vs $1/(\ln L)^2$. The line is a linear fit.

# Outline

1. Introduction

2. Theoretical Background

3. Monte Carlo Methods

4. Implementation Details

5. Results and Analysis

6. Conclusion

# Summary and Significance

- Successfully simulated the BKT transition in the 2D XY model:
  - Implemented efficient Wolff algorithm with parallel tempering
  - Measured key observables across multiple system sizes
  - Applied finite-size scaling to extract critical temperature

# Summary and Significance

- Successfully simulated the BKT transition in the 2D XY model:
  - Implemented efficient Wolff algorithm with parallel tempering
  - Measured key observables across multiple system sizes
  - Applied finite-size scaling to extract critical temperature
- Key findings:
  - Confirmed the characteristic signatures of the BKT transition
  - Determined $T_{KT} \approx 0.893(4)J$ in good agreement with theory
  - Observed universal jump in spin stiffness
  - Demonstrated vortex unbinding mechanism

# Summary and Significance

- Successfully simulated the BKT transition in the 2D XY model:
  - Implemented efficient Wolff algorithm with parallel tempering
  - Measured key observables across multiple system sizes
  - Applied finite-size scaling to extract critical temperature
- Key findings:
  - Confirmed the characteristic signatures of the BKT transition
  - Determined $T_{KT} \approx 0.893(4)J$ in good agreement with theory
  - Observed universal jump in spin stiffness
  - Demonstrated vortex unbinding mechanism
- Broader implications:
  - BKT transition represents a unique universality class
  - Demonstrates the power of topological defects in phase transitions
  - Provides insight into universal mechanisms across diverse physical systems
  - Highlights importance of advanced sampling techniques for critical phenomena

# Future Directions

- Methodological improvements:
  - Further optimization for larger system sizes
  - GPU implementation for parallel computation
  - Advanced analysis techniques for critical properties

# Future Directions

- Methodological improvements:
  - Further optimization for larger system sizes
  - GPU implementation for parallel computation
  - Advanced analysis techniques for critical properties
- Extensions to related models:
  - XY model with disorder or frustration
  - Quantum XY models and quantum phase transitions
  - Higher-dimensional models with topological transitions

# Future Directions

- Methodological improvements:
  - Further optimization for larger system sizes
  - GPU implementation for parallel computation
  - Advanced analysis techniques for critical properties
- Extensions to related models:
  - XY model with disorder or frustration
  - Quantum XY models and quantum phase transitions
  - Higher-dimensional models with topological transitions
- Applications:
  - Superconducting thin films and arrays
  - Quantum computing platforms based on 2D systems
  - Cold atom experiments with vortex dynamics
  - Theoretical connections to fundamental physics

# References I

[1] Victor Drouin-Touchette. “The kosterlitz-thouless phase transition: an introduction for the intrepid student”. In: *arXiv preprint arXiv:2207.13748* (2022).

ETH zürich

**Lara Turgut** *lturgut@ethz.ch*
ETH Zurich, Department of Physics

**Francesco Conoscenti** *fconoscenti@ethz.ch*
ETH Zurich, Department of Information Technology and Electrical Engineering

**Ben Bullinger** *bben@ethz.ch*
ETH Zurich, Department of Computer Science