

Notes on Approximate Leave-One-Out for Elastic Net

Linyun He

Wanchao Qin

Peng Xu

Yuze Zhou

July 24, 2018

1 ALO for Elastic Net, Approximation in the Primal Domain

Recall the objective function for the elastic net problem:

$$\min_{\beta} \frac{1}{2} \sum_{j=1}^n (\mathbf{x}_j^\top \beta - y_j)^2 + \lambda \left(\alpha \|\beta\|_1 + \frac{1-\alpha}{2} \|\beta\|_2^2 \right).$$

Let $A = \{i : \beta_i \neq 0, i = 1, \dots, p\}$ be the active set, we have

$$\dot{\ell}(\mathbf{x}_j^\top \beta; y_j) = \mathbf{x}_j^\top \beta - y_j, \quad \ddot{\ell}(\mathbf{x}_j^\top \beta; y_j) = 1, \quad \nabla^2 R(\hat{\beta}_A) = (1-\alpha)\lambda \mathbf{I}_{A,A}.$$

Thus, Eqn. 31 reduces to

$$\mathbf{H} = \mathbf{X}_{\cdot,A} \left[\mathbf{X}_{\cdot,A}^\top \mathbf{X}_{\cdot,A} + (1-\alpha)\lambda \mathbf{I}_{A,A} \right]^{-1} \mathbf{X}_{\cdot,A}^\top.$$

By augmenting \mathbf{X} with an extra column of 1s, adding the intercept back to the model is straightforward, as Eqn. 31 now becomes

$$\mathbf{H} = [\mathbf{1}_n, \mathbf{X}_{\cdot,A}] \left\{ [\mathbf{1}_n, \mathbf{X}_{\cdot,A}]^\top \mathbf{D} [\mathbf{1}_n, \mathbf{X}_{\cdot,A}] + \nabla^2 R(\hat{\beta}_0, \hat{\beta}_A) \right\}^{-1} [\mathbf{1}_n, \mathbf{X}_{\cdot,A}]^\top,$$

where

$$\mathbf{D} = \text{diag} \left[\ddot{\ell}(\hat{\beta}_0 + \mathbf{x}_j^\top \hat{\beta}; y_j) \right]_{j \in A} = \mathbf{I}_{A,A}, \quad \nabla^2 R(\hat{\beta}_0, \hat{\beta}_A) = \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & (1-\alpha)\lambda & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (1-\alpha)\lambda \end{bmatrix}.$$

Therefore, the ALO update is

$$\begin{bmatrix} 1 & \mathbf{x}_i^\top \end{bmatrix} \begin{bmatrix} \tilde{\beta}_0^{\setminus i} \\ \tilde{\beta}^{\setminus i} \end{bmatrix} = (\hat{\beta}_0 + \mathbf{x}_i^\top \hat{\beta}) + \frac{\mathbf{H}_{ii}}{1 - \mathbf{H}_{ii} \ddot{\ell}(\hat{\beta}_0 + \mathbf{x}_i^\top \hat{\beta}; y_i)} \dot{\ell}(\hat{\beta}_0 + \mathbf{x}_i^\top \hat{\beta}; y_i)$$

2 ALO for Elastic Net, Approximation with Proximal Formulation

For the elastic net problem, the proximal mapping is known to be

$$\mathbf{prox}_R(z) = \gamma \operatorname{sgn}(z) \odot (|z| - \lambda \mathbf{1}_p)_+, \quad \gamma = \frac{1}{1 + (1 - \alpha)\lambda}.$$

Let E be the active set, if $z_i \in E$, then

$$\frac{\partial}{\partial z_i} \gamma \operatorname{sgn}(z_i)(|z_i| - \lambda)_+ = \gamma.$$

Plug in $z = \hat{\beta} - \sum_{j=1}^n \dot{\ell}(\mathbf{x}_j^\top \hat{\beta}; y_j) \mathbf{x}_j$, Eqn. 46 thus reduce to

$$\mathbf{H} = \gamma \mathbf{X}_{\cdot, E} \left[\gamma \mathbf{X}_{\cdot, E}^\top \mathbf{X}_{\cdot, E} + (1 - \gamma) \mathbf{I}_{E, E} \right]^{-1} \mathbf{X}_{\cdot, E}^\top.$$

Bringing back the intercept term is straightforward as well. Noted that

$$\begin{bmatrix} \hat{\beta}_0^{\setminus i} \\ \hat{\beta}^{\setminus i} \end{bmatrix} = \mathbf{prox}_R(z), \quad z = \begin{bmatrix} \hat{\beta}_0^{\setminus i} \\ \hat{\beta}^{\setminus i} \end{bmatrix} - \sum_{j \neq i} \begin{bmatrix} 1 \\ \mathbf{x}_j \end{bmatrix} \dot{\ell}(\hat{\beta}_0^{\setminus i} + \mathbf{x}_j^\top \hat{\beta}^{\setminus i}; y_j).$$

Hence, from the first-order condition $\sum_{j \neq i} \dot{\ell}(\hat{\beta}_0^{\setminus i} + \mathbf{x}_j^\top \hat{\beta}^{\setminus i}; y_j) = 0$, we can derive that

$$\mathbf{J}_{E, E} = [\mathbf{J}(\mathbf{u})]_{E, E} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & (1 - \alpha)\lambda & 0 & \dots & 0 \\ 0 & 0 & (1 - \alpha)\lambda & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & (1 - \alpha)\lambda \end{bmatrix}^{-1}.$$

The ALO formula is then immediate by Thm. 5.1.

3 ALO for LASSO, with Intercept through Generalized LASSO

For the generalized LASSO problem

$$\min_{\beta} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \|\mathbf{D}\beta\|_1,$$

the dual problem is derived as:

$$\min_{\mathbf{u}} \frac{1}{2} \|\mathbf{y} - \boldsymbol{\theta}\|_2^2, \quad \boldsymbol{\theta} \in \{\mathbf{X}^\top \boldsymbol{\theta} = \mathbf{D}^\top \mathbf{u}, \|\mathbf{u}\|_\infty \leq \lambda\}.$$

The dual problem could be written in a proximal approach, such that:

$$\hat{\mathbf{u}} = \mathbf{prox}_R(\mathbf{y}), \quad R(\mathbf{u}) = \begin{cases} 0 & \boldsymbol{\theta} \in \{\mathbf{X}^\top \boldsymbol{\theta} = \mathbf{D}^\top \mathbf{u}, \|\mathbf{u}\|_\infty \leq \lambda\}, \\ \infty & \text{otherwise.} \end{cases}$$

Denote \mathbf{J} as the Jacobian of the proximal operator at the full data problem \mathbf{y} , then the ALO estimator could be obtained as:

$$\mathbf{y}^{\setminus i} = \mathbf{y}_i - \frac{\hat{\mathbf{u}}_i}{\mathbf{J}_{ii}}.$$

For the case of LASSO with an intercept, we could expand the \mathbf{X} with a column of ones in the first column, expand β with another dimension and choose $\mathbf{D} = [\mathbf{0}, \mathbf{I}]$. Let $E := \{j : |\mathbf{X}_j^\top \theta| = \lambda\}$ denote the active set. The Jacobian is locally given as the projection onto the orthogonal complement of the span of \mathbf{X}_E and the vector of ones. Further denote $\tilde{\mathbf{X}}_E = [\mathbf{1}, \mathbf{X}_E]$, then the Jacobian is given as $\mathbf{I} - \tilde{\mathbf{X}}_E(\tilde{\mathbf{X}}_E^\top \tilde{\mathbf{X}}_E)^{-1} \tilde{\mathbf{X}}_E^\top$.

4 Usage of ALO formulae with glmnet package

The glmnet package scales the elastic net loss function by a factor of $1/n$, so the ALO formulae must be adjusted accordingly, e.g. for the proximal one, we instead have:

$$\tilde{\mathbf{y}}_j^{\setminus i} = \hat{\mathbf{y}}_j + \frac{\mathbf{H}_{ii}(\hat{\mathbf{y}}_j - \mathbf{y}_j)}{n - \mathbf{H}_{ii}}, \quad \mathbf{H} = \gamma \mathbf{X}_{\cdot, E} \left[\frac{\gamma}{n} \mathbf{X}_{\cdot, E}^\top \mathbf{X}_{\cdot, E} + (1 - \gamma) \mathbf{I}_{E, E} \right]^{-1} \mathbf{X}_{\cdot, E}^\top.$$

Furthermore, glmnet implicitly “standardizes \mathbf{y} to have unit variance before computing its λ sequence (and then unstandardizes the resulting coefficients)”. So to get comparable result, it is necessary to rescale \mathbf{y} by the MLE $\hat{\sigma}_y$ before fitting the model. Figure 1 shows the comparison of the ALO and LOO for different α s. Without standardizing \mathbf{y} first, a growing discrepancy between the two curves can be observed as $\alpha \rightarrow 0$.




Figure 1: ALO vs. LOO for Elastic Net.