

The Global Biodiversity Framework supports global assessment of One Health actions

Francis Banville¹, Claire Burnel², Colin Carlson³, Elodie Eiffener⁴, Gabriel Munoz Acevedo⁵, Andrea Paz¹ and Timothée Poisot¹

¹ Université de Montréal, Département de sciences biologiques, Montréal Québec, Canada; ² Université de Montréal, Département de sociologie, Montréal Québec, Canada; ³ Yale University, Yale School of Public Health, New Haven Connecticut, USA; ⁴ Karolinska Institutet, Institute of Environmental Medicine, Stockholm, Sweden; ⁵ Canadian Institute for Health Information, Ottawa Ontario, Canada

Correspondence to Francis Banville — francis.banville@umontreal.ca

Abstract: The One Health approach promotes collaboration across disciplines to enhance the health of humans, animals, plants, and the environment. Recently developed by the Quadripartite organizations, the One Health Joint Plan of Action (2022-2026) supports countries in adopting the One Health approach through six action tracks. The tracks address multiple aspects of biodiversity, containing guiding actions on: One Health capacity-building (track 1), zoonotic diseases (tracks 2 and 3), food and agriculture (track 4), antimicrobial resistance (track 5), and environmental health (track 6). However, there are currently no indicators for monitoring the implementation, state and progress of these tracks. Our research examines the extent to which they can be assessed by the existing monitoring framework of the Kunming-Montreal Global Biodiversity Framework (KM-GBF) of the Convention on Biological Diversity (CBD). We evaluated (1) the link between each indicator of the KM-GBF monitoring framework and human, animal, plant, and environmental health, respectively, and (2) the usability of these indicators for monitoring the action tracks. We found that 75% of the KM-GBF indicators are associated with either human (56%), animal (53%), plant (51%), or environmental (63%) health, and that 91% of them can be used for monitoring at least one action track. Reusing existing indicators alleviates the costs associated with the development and measurement of new indicators, thus improving the efficiency of monitoring systems. Our work supports the idea that multilateral progress on the monitoring of biodiversity, through its indissociability from all

facets of health, provides a way to retrospectively and prospectively measure progress on ambitious One Health actions.

Keywords: biodiversity indicators, One Health Joint Plan of Action, Quadripartite organizations, human health, animal health, plant health, environmental health

1 Introduction

2 Human health responds to changes in biodiversity in complex and sometimes opposite ways
3 (Robinson et al., 2024). On one hand, biodiversity supplies food, shelter, and medicine that are
4 vital to human health (Diaz et al., 2018). On the other hand, infectious diseases originating in
5 wildlife are a serious threat to human health, the transmission of pathogens from animals to
6 humans being a major cause of disease outbreaks (K. E. Jones et al., 2008). Ecosystem services,
7 such as climate stabilization, air and water filtration, and natural hazards mitigation, support
8 human societies by maintaining safe and adequate environmental conditions (Jackson et al.,
9 2013). However, plant diseases can hinder ecosystem services, e.g. through food contamina-
10 tion or the reduction in food production (Al-Sadi, 2017). The main threats to biodiversity
11 (biological invasions, climate change, habitat loss, overexploitation, and pollution; Bellard
12 et al. (2022)) all impact human health, notably by driving the emergence of infectious dis-
13 eases(Carlson et al., 2025). Because biodiversity and human health are highly interconnected,
14 both can be monitored using a similar set of tools (Poisot et al., 2025).

15 The One Health approach (OHHLEP et al., 2022; Winkler et al., 2025) recognizes the
16 importance of biodiversity for public health (Winkler et al., 2025). It addresses complex
17 public health challenges at the interface between human, animal, plant, and environmental
18 health (Rabinowitz & Conti, 2013) by promoting collaboration between health professionals,
19 veterinarians, and biologists. The One Health Joint Plan of Action (OH JPA) (2022-2026) is an
20 action-oriented framework aimed at advancing One Health principles by 2026 (Quadripartite
21 organizations, 2022). Developed by the Quadripartite organizations, which is a strategic
22 partnership between the Food and Agriculture Organization of the United Nations (FAO),
23 the United Nations Environment Programme (UNEP), the World Organisation for Animal
24 Health (WOAH), and the World Health Organization (WHO), the OH JPA contains 19 actions
25 that countries can take to improve One Health capacity-building, political commitment,
26 governance structure, funding, and knowledge systems. These actions form six tracks that
27 collectively seek to strengthen health systems (track 1), prevent zoonotic diseases (tracks 2 and
28 3), improve food safety (track 4), reduce antimicrobial resistance (track 5), and stop environ-
29 mental degradation (track 6). As we enter the final year of the plan, there is a need to generate
30 clear and objective information regarding the progress made since its adoption in 2022. This
31 evaluation is essential given the non-binding nature of the OH JPA, as it would encourage
32 effective policy development by the States and inform the development of the next version

of the plan. However, there are currently no indicators for monitoring the implementation of this plan.

Biodiversity indicators are measures of changes in the state of biodiversity over time that track progress towards specific objectives (Hébert et al., 2025; J. P. G. Jones et al., 2011). As such, they are intended to serve several purposes, the main ones being to simplify complex phenomena which are difficult to measure directly, and to improve accountability and effectiveness through prioritization, evaluation and communication of objectives and policies (Amuasi & Winkler, 2025; Gudmundsson, 2003; Nicholson et al., 2012). A solid example of monitoring framework can be found in the Kunming-Montreal Global Biodiversity Framework (KM-GBF) (UNEP, 2022a), adopted in 2022 by the Parties to the Convention on Biological Diversity (CBD). The KM-GBF aims to halt and reverse biodiversity loss by 2030 through 4 general goals and 23 specific targets, ranging from the restoration of degraded ecosystems to the integration of biodiversity in decision-making. For the first time since the CBD was signed in 1992, the KM-GBF includes a monitoring framework (UNEP, 2025) made up of 204 indicators that countries can use to monitor these goals and targets. These indicators are classified in four groups: headline (high-level), binary (yes/no questions), component (technical), and complementary (supporting) indicators. Only two of these groups (headline and binary indicators) are mandatory when reporting national progress to the CBD. This reliance on policy evaluation and indicators is in line with a broader trend that some refer to as an “indicator industry” (Lehtonen, 2015), which aims to provide more precise and quantitative assessments of policies. The CBD serves as a good example of how such indicators constitute “soft instruments” capable of incentivizing States to act when the international agreement is not legally binding (Kirsop-Taylor, 2022).

More recently, the Parties to the CBD also adopted the Global Action Plan on Biodiversity and Health (GAP-BH), which formally recognizes the links between biodiversity and human health (UNEP, 2024). The GAP-BH groups the targets of the KM-GBF into 14 thematic categories, encompassing a variety of topics ranging from “Land and sea use” (category 1) to “Knowledge and engagement of people” (category 14). Each thematic category has its own relevance to human health. For example, the targets that fall under the “Land and sea use” category, which include the “restoration of 30% of all degraded ecosystems” (target 2) and the “conservation of 30% of land, waters and seas” (target 3), contribute to “the continued provision of nature’s contributions to people, which in turn support health and reduce disease emergence and transmission among wildlife, livestock and people” (UNEP, 2024). Because of the high relevance of all KM-GBF targets to human health as described in UNEP (2024), we

hypothesized that a large proportion of the indicators of these targets would also be relevant to health, and therefore to the retroactive monitoring of the OH JPA. Given the recent increase in prominence of the One Health concept following its introduction in the Article 5 of the WHO Pandemic Agreement (WHO, 2025), it is also crucial to anticipate on the reporting need of a future high-level document for One Health, and to suggest existing indicators and reporting processes that can support its implementation.

We argue that reusing indicators for both monitoring systems (the KM-GBF and the OH JPA) would improve their efficiency and reduce the workload of countries. During the vote of the KM-GBF monitoring framework, delegates warned that too many indicators could hinder monitoring, especially in less well-resourced nations (Carroll et al., 2024). Colinas et al. (2026) present best practices in selecting indicators for monitoring One Health. They suggest to review existing indicators rather than devoting resources to designing a new standardized and aggregated single index such as the One Health Index (de Moura et al., 2022) and the Global One Health Index (Zhang et al., 2022). Our work serves as a first step in the framework they propose by quantifying the extent to which the OH JPA is already covered by existing biodiversity indicators. Moreover, in the absence of a prior monitoring system, many of the KM-GBF indicators still lack an agreed-upon methodology, both at the national and international levels (Affinito et al., 2025): a further reason why pooling efforts would help improve both this framework and the OH JPA. Despite these difficulties, the indicators of the KM-GBF have ultimately achieved a broad global consensus, and this grants them an important legitimacy as reporting instruments in other context, whenever they are relevant. Therefore, we aimed to identify the indicators of the KM-GBF monitoring framework that are relevant to One Health monitoring. First, we assessed the link between each of these indicators and human, animal, plant, and environmental health, respectively. Then, we evaluated the usability of the indicators for monitoring the specific action tracks of the OH JPA. Our work shows that biodiversity monitoring is an essential component of effective One Health strategies.

Methods

We qualitatively assessed the relevance of all 204 KM-GBF indicators (UNEP, 2025) to health. This was done independently by at least two evaluators for each of the indicators, this process leading to a consensus among the coauthors for all indicators. Specifically, we evaluated the degree to which the indicators are linked to each of the four pillars of One Health (i.e., to human, animal, plant, and environmental health), as well as their usability for monitoring the

OH JPA. Our assessments were informed by the metadata (rationale, definition, method of computation, data sources, and scale of use) of the indicators provided by the World Conservation Monitoring Centre of the UN Environment Programme (UNEP-WCMC, 2025), when available, looking for explicit considerations of health. When such metadata was unavailable, assessments were supplemented by expert elicitation. The result of this analysis and the code to reproduce the figures are available on Zenodo (<https://doi.org/10.5281/zenodo.20085701>).

Box 1: What are animals, plants, and the environment in One Health?

The definitions of animals, plants, and the environment in One Health are based more on institutional norms than on biological realities. Here, we adhere to the delineation of these groups by mandates of the four organizations that form the Quadripartite. Humans are their own pillar of One Health and the primary focus of the WHO. Animals and plants refer to *domesticated* species only, a distinction we make to align more closely with the mandates of the WOA (animals) and FAO (plants). Therefore, we include pets, livestock, and edible marine and freshwater species captured from fisheries and aquaculture in the animal health pillar, i.e. species mainly looked after by veterinarians and food inspectors. Even though, biologically speaking, humans and wildlife are also animals, they are not part of the animal health pillar. Similarly, we include all cultivated plants used for food, fuel, and medicine into the plant health pillar. Wildlife and uncultivated plant species, such as wild fish and tree species used in forestry, are part of the environment health pillar, alongside all other species absent from the other three pillars. The environment health pillar also includes whole ecosystems (natural and artificial), which are central to UNEP's mission. These definitions are aligned with the core contributing sciences of the One Health approach, i.e. human medicine, veterinary medicine, agriculture, and ecology (Gibbs, 2014; Lerner & Berg, 2017). While it may be biologically frustrating to work within this framework, we emphasize that assigning groups of organisms to specific organizations has the direct benefit of inscribing our work within established, and multilaterally agreed-upon, systems of governance and reporting.

Box 2: What is health?

Throughout this manuscript, we use the overarching term “health” when referring to the health of different species and the environment, and refer to human, animal, plant, and environmental health when wanting to differentiate. Health manifests differently depending on the level of biological organization (Lerner & Berg, 2015). At the individual level, we use a broad definition of health that considers the overall well-being of individuals and their capacity to function normally, which includes the absence of diseases (WHO, 1948). In humans and domestic and wildlife animals, well-being is the ability for an individual to satisfy its physical, mental, and behavioral needs (Webb et al., 2019), whereas diseases are undesirable conditions often leading to pain, suffering, or death. This is addressed, in the spirit of One Health, holistically at the individual, populational, and global scale, and therefore accounts for *e.g.* health system preparedness. In plants, which are not conscious organisms (Mallatt et al., 2021), health is the extent to which they are able to function physiologically, and diseases are impediments to their normal physiological functioning often leading to death (Döring et al., 2012). Because domesticated animals and plants are mainly used for human consumption, animal and plant diseases have important consequences for food security (Garcia et al., 2020; Strange & Scott, 2005). At the ecosystem level, consisting of many species and their interactions, health is ill-defined (Schaeffer et al., 1988). The idea of ecosystem health, although ancient in Indigenous and non-Western cosmological perspectives (Descola, 2013; Hierro, 2005), has often been promoted as metaphorical (Costanza & Mageau, 1999; Rapport, 1989). We consider environmental health as the extent to which an ecosystem can maintain its biological and chemical processes and adapt to changes (Jakobsson, 2012). Disturbances, such as extreme weather events or resource exploitation, are the ecosystem analogue of diseases, and represent degradations that usually lead to a decline in ecosystem functioning. Environmental health also includes the health of species that are part of the environment pillar of One Health, *e.g.* wildlife and uncultivated plant species.

Assessing the link between KM-GBF indicators and health

We classified the degree to which the KM-GBF indicators are linked to each of the four pillars of One Health into four categories: direct connection, indirect connection, potential connection, and no connection. An indicator was classified as directly connected to health when there is a direct causal relationship between the indicator and health. This is the case,

for instance, when the indicator directly measures a condition or determinant of health. In contrast, an indicator was classified as indirectly connected to health when there is an intermediary factor between the indicator and health, and as potentially connected when there are multiple intermediary factors or when its connection to health is likely but difficult to certify. Finally, an indicator was classified as not connected to health when the connection is unlikely, unsupported by current scientific evidence, or absent. For each of the four pillars of One Health, we counted the number of indicators assigned to each category.

For example, headline indicator 7.2 “Pesticide environment concentration and/or aggregated total applied toxicity” was considered directly connected to human health because of the toxicity of many pesticides under high level of exposure (Kim et al., 2017). It was also considered directly connected to animal, plant, and environmental health because the use of pesticides can benefit the growth of cultivated plants through the reduction of herbivory and competition, while being highly harmful to animals and biodiversity (Mahmood et al., 2016). In contrast, component indicator 4.CT.1 “Number of plant and animal genetic resources for food and agriculture secured in either medium- or long-term conservation facilities” was considered indirectly connected to human health because conserving genetic resources increases the resilience of food systems to global disasters (Esquinas-Alcázar, 2005), which in turn protects human health. However, we did not find any evidence of a link between this indicator and animal, plant, or environmental health.

Assessing the usability of KM-GBF indicators for monitoring One Health action tracks

For each action track of the OH JPA, we classified the KM-GBF indicators into three categories: directly usable, usable after adaptation, and not usable. Directly usable indicators can be used in their current form to monitor the implementation of at least one action in an action track. They can detect trends relevant to the actions of the track and do not require adaptations. For example, we qualified headline indicator 7.2 “Pesticide environment concentration and/or aggregated total applied toxicity” as being directly usable for monitoring the sixth action track because it is a direct measure of environmental degradation and contamination. In contrast, indicators usable after adaptation need to be slightly modified before being used to monitor actions in an action track. These adaptations should be small changes in the scale of measurement, the data resolution, or the taxa monitored by the indicator. For instance, headline indicator 7.2 was qualified as being usable after adaptation for monitoring the fourth action track on food safety risks. To be more relevant for this action track, this indicator could refer more specifically to the use of pesticides in agriculture. Finally, not usable indicators

are outside the scope of an action track or need to be greatly modified before being used to monitor the implementation of an action track.

We assessed usability for each action track independently, i.e. an indicator can be useful for monitoring multiple action tracks. For each action track, we counted the total number of indicators in each category. We also counted the number of usable indicators for the whole plan regardless of the action tracks. Then, we partitioned these numbers based on the group of indicators in the KM-GBF monitoring framework (i.e. headline, binary, component, and complementary indicators) and the thematic categories of the GAP-BH. Finally, for each action track where an indicator was considered usable (either directly or after adaptation), we identified the most relevant action that can be monitored by the indicator, and counted the number of indicators associated with each action.

Results and discussion

Link between KM-GBF indicators and health

We found that 75% of KM-GBF indicators are either directly or indirectly connected to either human, animal, plant, or environmental health, with approximately 55% of indicators being directly connected to health. This signifies that monitoring the state, benefits, and pressure of biodiversity and our responses to biodiversity loss provides extensive health information. In Figure 1, we show the number of indicators that are directly, indirectly, potentially, and not connected to each of the four pillars of One Health. The number of directly connected indicators is, as expected, slightly higher for environmental health (41% of indicators) compared to human (27%), animal (26%), and plant (28%) health, mostly because the environment trivially contains most of biodiversity. However, this difference is less pronounced after adding indirectly connected indicators (63% for environmental health compared to 56%, 53%, and 51% respectively for human, animal, and plant health), highlighting the multiple pathways through which biodiversity and health are connected.

The fact that most KM-GBF indicators are connected to health is not surprising. The GAP-BH recognizes the relevance to health of all 23 targets of the KM-GBF, and because the indicators measure the progress made towards these targets, they also measure the progress made towards a healthier planet. However, we did not expect that comparable numbers of indicators would be linked to the four pillars of One Health, indicating that similar amount of biodiversity information can be reused, either directly or indirectly, to monitor human,




Figure 1: **Number of indicators of the Kunming-Montreal Global Biodiversity Framework linked with human, animal, plant, and environmental health.** Each indicator was classified as being either directly connected (blue), indirectly connected (green), potentially connected (orange), or not connected (red) to each of the four pillars of One Health.

animal, plant, and environmental health. This result therefore suggests that the core components of One Health can all be meaningfully informed by biodiversity monitoring.

Usability of KM-GBF indicators for monitoring One Health action tracks

The vast majority of KM-GBF indicators (91%) can be used either directly or after adaptation to monitor the OH JPA action tracks, and around 74% of indicators can be used directly. This signifies that we can already do an extensive evaluation of One Health actions with existing data and methodologies. However, we show in Figure 2 that the usability of indicators greatly differs depending on the action track. The proportion of usable indicators varies from 20% for the fifth action track on antimicrobial resistance (4% of directly usable indicators) to 79% for the sixth action track on the integration of the environment into One Health (62% of directly usable indicators). The high proportion of usable indicators for the sixth action track is not surprising given that the KM-GBF is a multilateral environmental agreement that aims to protect species and restore ecosystems. The low proportion of indicators that can monitor

actions in the fifth action track could be due to the characteristics of these actions, which are more focused on collaboration, capacity-building and awareness-raising than the decrease and prevention of antimicrobial resistance through biodiversity. Similarly, the actions in the third action track on endemic zoonotic, neglected tropical, and vector-borne diseases, which can only be monitored by 33% of the indicators, are more centered on capacity-building and less on the drivers and inhibitors of infectious diseases, especially in comparison to the actions of the second action track on zoonotic epidemics and pandemics. These differences suggest that, although the KM-GBF monitoring framework can greatly support One Health agencies by reducing the need to develop, gain expertise in, and monitor new indicators at the intersection of biodiversity and health, additional work is still needed to equally cover all aspects of the plan.




Figure 2: Number of indicators of the Kunming-Montreal Global Biodiversity Framework that can be used to monitor at least one action in each of the action tracks of the One Health Joint Plan of Action. The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The proportion of usable indicators (either directly or after adaptation) for each action track is indicated. The bottom bar and proportion designate the indicators that can monitor at least one action in the whole plan.

The number of indicators associated with each action in the OH JPA is presented in Table 1. There are high discrepancies in the usability of indicators between actions. The total number of usable indicators (either directly or after adaptation) varies from 1 (action 5.3: Strengthen global AMR governance structures) to 108 (action 6.1: Protect, restore, and prevent the degradation of ecosystems and the wider environment). Within a single action track, the biggest differences are found in the sixth action track, where only 12 indicators were associated to action 6.4 whereas 108 indicators were associated to action 6.1 This result shows that the KM-GBF monitoring framework can provide substantial information for some of the actions, but it also further highlights the need to identify or develop new indicators for specific actions of the plan, even within the action tracks that can be monitored by numerous indicators, such as action track 6.

Most KM-GBF indicators can be used to monitor the action tracks, for each of the groups of indicators in the KM-GBF monitoring framework (Figure 3). We evaluated that 100% of headline indicators (27/27), 93% of binary indicators (14/15), 94% of component indicators (49/52), and 87% of complementary indicators (96/110) can be used, either directly or after adaptation, to monitor at least one action track. Headline and binary indicators are the only two mandatory groups of indicators, i.e. that Parties need to include in their national biodiversity reports. The high proportion of headline and binary indicators is encouraging, because it indicates that the information most likely to figure in national reports will be reusable for evaluating the implementation of the OH JPA. On the other hand, the high reusability of component and complementary indicators, which are currently optional, should be seen as an additional incentive to measure them.

For each thematic category of the GAP-BH, at least 89% of indicators can be used, either directly or after adaptation, for monitoring at least one action track, except in the “Knowledge and engagement of people” category, where only 52% of the indicators can be used (Figure 4). This suggests that every sector of the KM-GBF monitoring framework, from “Nature’s contribution to people” to “Species management”, is relevant for One Health, even though there are wide variations in the action track most relevant for each category. For instance, the three indicators under “Urban areas” can only be used after adaptation to monitor the action tracks, apart from tracks 1 and 6, yet they do not capture actions related to antimicrobial resistance (track 5). Figure 4 also shows that all indicators that can directly monitor the fifth action track on antimicrobial resistance are in the “Access and benefit-sharing” and “Biosafety and biotechnology” categories, further emphasizing the importance of collaboration, capacity-building and awareness-raising set forth in this action track.

Table 1: Number of indicators of the Kunming-Montreal Global Biodiversity Framework that can be used directly and after adaptation to monitor each of the actions of the One Health Joint Plan of Action. The total number of usable indicators for each action track is indicated in the highlighted blue lines. A specific indicator can be used to monitor multiple action tracks, but only the most relevant action in each action track was identified for each indicator.

Action	Directly usable	Usable after adaptation	Total usable
1 Enhancing One Health capacities to strengthen health systems	46	61	107
1.1 Establish the foundations for One Health capacities	30	5	35
1.2 Generate mechanisms, tools, and capacities to establish a One Health competent workforce and the frameworks/processes to facilitate One Health work	8	50	58
1.3 Generate an enabling environment for the effective implementation of One Health	8	6	14
2 Reducing the risks from emerging and re-emerging zoonotic epidemics and pandemics	40	63	103
2.1 Understand the drivers of emergence, spillover, and spread of zoonotic pathogens	16	27	43
2.2 Identify and prioritize targeted, evidence-based upstream interventions to prevent the emergence, spillover, and spread of zoonotic pathogens	7	6	13
2.3 Strengthen national, regional, and global One Health surveillance, early warning, and response systems	17	30	47
3 Controlling and eliminating endemic zoonotic, neglected tropical and vector-borne diseases	25	43	68
3.1 Enable countries to develop and implement community-centric and risk-based solutions to endemic zoonotic, neglected tropical, and vector-borne disease control using a One Health approach involving all relevant stakeholders	3	14	17
3.2 Ensure the harmonized application of One Health principles at all levels by implementing practical measures to strengthen local, national, regional, and global policy frameworks for the control and prevention of endemic zoonotic, neglected tropical, and vector-borne diseases	18	10	28
3.3 Increase political commitment and investment in the control of endemic zoonotic, neglected tropical, and vector-borne diseases, by advocating for and demonstrating the value of a One Health approach	4	19	23
4 Strengthening the assessment, management and communication of food safety risks	42	70	112
4.1 Strengthen the One Health approach in national food control systems and food safety coordination	18	30	48
4.2 Utilize and improve food systems data and analysis, scientific evidence, and risk assessment in developing policy and making integrated risk management decisions	15	21	36
4.3 Foster the adoption of the One Health approach in national foodborne disease surveillance systems and research for the detection and monitoring of foodborne disease and food contamination	9	19	28
5 Curbing the silent pandemic of antimicrobial resistance (AMR)	9	31	40
5.1 Strengthen the capacity and knowledge of countries to prioritize and implement context-specific collaborative One Health work to control AMR in policy, legislation, and practice	7	17	24
5.2 Reinforce global and regional initiatives and programmes to influence and support One Health responses to AMR	2	13	15
5.3 Strengthen global AMR governance structures	0	1	1
6 Integrating the environment into One Health	126	36	162
6.1 Protect, restore, and prevent the degradation of ecosystems and the wider environment	92	16	108
6.2 Mainstream the health of the environment and ecosystems into the One Health approach	9	7	16
6.3 Integrate environmental knowledge, data, and evidence into One Health decision-making	20	6	26
6.4 Create an interoperable One Health academic and in-service training programme for environmental, medical, agricultural, and veterinary sector professionals	5	7	12




Figure 3: Proportion of indicators of the Kunming-Montreal Global Biodiversity Framework that can be used to monitor at least one action in each of the action tracks of the One Health Joint Plan of Action, partitioned by the different groups of indicators. The number of indicators in each group is indicated in parentheses. The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The gray bars designate the indicators that can monitor at least one action in the whole plan.

Conclusion

Instead of developing and measuring new indicators, which can be resource-intensive, the high reusability of the KM-GBF indicators for monitoring the implementation of the OH JPA can increase the efficiency of both monitoring systems. The environmental sector can meaningfully contribute to One Health efforts by sharing their data, tools, and expertise with national One Health agencies, which can in turn provide further incentives and means to measure relevant indicators. Making indicators serve more than one purpose can also improve the efficiency of public and private spending by avoiding duplication of work. Reusing existing indicators can thus greatly reduce the workload of countries that are part of different multilateral environmental agreements with overlapping objectives. However, additional effort is needed to identify or develop new indicators to monitor the action tracks where few KM-GBF



Figure 4: Proportion of indicators of the Kunming-Montreal Global Biodiversity Framework that can be used to monitor at least one action in each of the action tracks of the One Health Joint Plan of Action, partitioned by the different thematic categories of the Global Action Plan on Biodiversity and Health. The number of indicators in each thematic category is indicated in parentheses. The thematic category of an indicator was assigned based on the thematic category of its target. Indicators that monitor goals A, B, C, and D of the Kunming-Montreal Global Biodiversity Framework are not assigned to any thematic category. The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The gray bars designate the indicators that can monitor at least one action in the whole plan.

indicators were found to be usable. These gaps could be filled by finding existing indicators in other multilateral environmental agreements, such as the global indicator framework of the Sustainable Development Goals (SDGs). Nevertheless, the high reusability of the KM-GBF indicators can greatly facilitate the monitoring of One Health action tracks. Putting in the work to monitor the implementation of the OH JPA will be an essential step to ensure it is effectively put into action and meets its goals. As Parties to the CBD submit their National Biodiversity Strategies and Action Plans (NBSAPs) with the monitored indicators (UNEP, 2022b), large amounts of new data will become available for the retrospective assessment of the progress made on the ambitious One Health actions.

Acknowledgments

We acknowledge that this study was conducted on land within the traditional unceded and unsundered territory of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat and Omàmiwininiwak nations. TP was funded through award no. 223764/Z/21/Z from the Wellcome Trust and supported by the US National Science Foundation (grant no. DBI 2515340). This manuscript is a contribution of the GEO BON One Health working group. We thank Katerina Gapanenko, from the Canadian Institute for Health Information, for her useful feedback on the manuscript.

References

- Affinito, F., Butchart, S. H. M., Nicholson, E., Hirsch, T., Williams, J. M., Campbell, J. E., Ferrari, M. F., Gabay, M., Gorini, L., Kalamujic Stroil, B., Kohsaka, R., Painter, B., Pinto, J. C., Scholz, A. H., Straza, T. R. A., Tshidada, N., Vallecillo, S., Widdicombe, S., & Gonzalez, A. (2025). Assessing coverage of the monitoring framework of the Kunming-Montreal Global Biodiversity Framework and opportunities to fill gaps. *Nature Ecology & Evolution*, 9(7), 1280–1294. <https://doi.org/10.1038/s41559-025-02718-3>
- Al-Sadi, A. M. (2017). Impact of Plant Diseases on Human Health. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 7(2), 21. https://doi.org/10.4103/ijnpnd.ijnpnd_24_17
- Amuasi, J. H., & Winkler, A. S. (2025). Future trajectories for One Health. *The Lancet*, 406(10502), 426–429. [https://doi.org/10.1016/S0140-6736\(25\)01453-9](https://doi.org/10.1016/S0140-6736(25)01453-9)
- Bellard, C., Marino, C., & Courchamp, F. (2022). Ranking threats to biodiversity and why it doesn't matter. *Nature Communications*, 13(1), 2616. <https://doi.org/10.1038/s41467-022-30339-y>
- Carlson, C. J., Brookson, C. B., Becker, D. J., Cummings, C. A., Gibb, R., Halliday, F. W., Heckley, A. M., Huang, Z. Y. X., Lavelle, T., Robertson, H., Vicente-Santos, A., Weets, C. M., & Poisot, T. (2025). Pathogens and planetary change. *Nature Reviews Biodiversity*, 1(1), 32–49. <https://doi.org/10.1038/s44358-024-00005-w>
- Carroll, C., Hoban, S., & Ray, J. C. (2024). Lessons from COP15 on effective scientific engagement in biodiversity policy processes. *Conservation Biology*, 38(2), e14192. <https://doi.org/10.1111/cobi.14192>

Colinas, J., Boiteux, M., & Aenishaenslin, C. (2026). Issues and principles for legitimate and meaningful One Health monitoring. *International Journal of Public Health*, 71, 1609612. <https://doi.org/10.3389/ijph.2026.1609612>

Costanza, R., & Mageau, M. (1999). What is a healthy ecosystem? *Aquatic Ecology*, 33(1), 105–115. <https://doi.org/10.1023/A:1009930313242>

de Moura, R. R., Chiba de Castro, W. A., Farinhas, J. H., Pettan-Brewer, C., Kmetiuk, L. B., dos Santos, A. P., & Biondo, A. W. (2022). One Health Index (OHI) applied to Curitiba, the ninth-largest metropolitan area of Brazil, with concomitant assessment of animal, environmental, and human health indicators. *One Health*, 14, 100373. <https://doi.org/10.1016/j.onehlt.2022.100373>

Descola, P. (2013). *Beyond Nature and Culture*. University of Chicago Press.

Diaz, S., Pascual, U., Stenseke, M., Martin-Lopez, B., Watson, R. T., Molnar, Z., Hill, R., Chan, K. M. A., Baste, I. A., Brauman, K. A., Polasky, S., Church, A., Lonsdale, M., Larigauderie, A., Leadley, P. W., van Oudenhoven, A. P. E., van der Plaats, F., Schroter, M., Lavorel, S., ... Shirayama, Y. (2018). Assessing nature's contributions to people. *Science*, 359(6373), 270–272. <https://doi.org/10.1126/science.aap8826>

Döring, T. F., Pautasso, M., Finckh, M. R., & Wolfe, M. S. (2012). Concepts of plant health – reviewing and challenging the foundations of plant protection. *Plant Pathology*, 61(1), 1–15. <https://doi.org/10.1111/j.1365-3059.2011.02501.x>

Esquinas-Alcázar, J. (2005). Protecting crop genetic diversity for food security: Political, ethical and technical challenges. *Nature Reviews Genetics*, 6(12), 946–953. <https://doi.org/10.1038/nrg1729>

Garcia, S. N., Osburn, B. I., & Jay-Russell, M. T. (2020). One Health for Food Safety, Food Security, and Sustainable Food Production. *Frontiers in Sustainable Food Systems*, 4. <https://doi.org/10.3389/fsufs.2020.00001>

Gibbs, E. P. J. (2014). The evolution of One Health: A decade of progress and challenges for the future. *Veterinary Record*, 174(4), 85–91. <https://doi.org/10.1136/vr.g143>

Gudmundsson, H. (2003). The Policy Use of Environmental Indicators - Learning from Evaluation Research. *The Journal of Transdisciplinary Environmental Studies*, 2(2), 1–12.

Hébert, K., Acevedo, D. H., Cameron, V., Courant, S., Daguet, C., Gonzalez, A., Gravel, D., Huot, J., Jousse, M., Juhasz, C.-C., Lévesque, C., Serrano, J., Simard, A., & Pollock, L. (2025). Selecting indicators to track progress towards the Global Biodiversity Framework: A case study of Quebec's 2030 Nature Plan. *FACETS*, 10, 1–13. <https://doi.org/10.1139/facets-2024-0357>

443 Hierro, P. G. (2005). *The Land Within: Indigenous Territory and the Perception of the Environ-*
444 *ment*. IWGLA.

445 Jackson, L. E., Daniel, J., McCorkle, B., Sears, A., & Bush, K. F. (2013). Linking ecosystem
446 services and human health: The Eco-Health Relationship Browser. *International Journal of*
447 *Public Health*, 58(5), 747–755. <https://doi.org/10.1007/s00038-013-0482-1>

448 Jakobsson, C. (2012). *Sustainable Agriculture*. Baltic University Press.

449 Jones, J. P. G., Collen, B., Atkinson, G., Baxter, P. W. J., Bubb, P., Illian, J. B., Katzner, T. E.,
450 Keane, A., Loh, J., McDonald-Madden, E., Nicholson, E., Pereira, H. M., Possingham, H. P.,
451 Pullin, A. S., Rodrigues, A. S. L., Ruiz-Gutierrez, V., Sommerville, M., & Milner-Gulland, E. J.
452 (2011). The Why, What, and How of Global Biodiversity Indicators Beyond the 2010 Target.
453 *Conservation Biology*, 25(3), 450–457. <https://doi.org/10.1111/j.1523-1739.2010.01605.x>

454 Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., & Daszak, P.
455 (2008). Global trends in emerging infectious diseases. *Nature*, 451(7181), 990–U4. <https://doi.org/10.1038/nature06536>

457 Kim, K.-H., Kabir, E., & Jahan, S. A. (2017). Exposure to pesticides and the associated
458 human health effects. *Science of The Total Environment*, 575, 525–535. <https://doi.org/10.1016/j.scitotenv.2016.09.009>

460 Kirsop-Taylor, N. (2022). *Chapter 4: Biodiversity and sustainable development governance*
461 (Handbook on the Governance of Sustainable Development).

462 Lehtonen, M. (2015). *Chapter 4: Indicators: Tools for informing, monitoring or controlling?* (The
463 Tools of Policy Formulation).

464 Lerner, H., & Berg, C. (2015). The concept of health in One Health and some practical impli-
465 cations for research and education: What is One Health? *Infection Ecology & Epidemiology*,
466 5(1), 25300. <https://doi.org/10.3402/iee.v5.25300>

467 Lerner, H., & Berg, C. (2017). A Comparison of Three Holistic Approaches to Health: One
468 Health, EcoHealth, and Planetary Health. *Frontiers in Veterinary Science*, 4, 163. <https://doi.org/10.3389/fvets.2017.00163>

470 Mahmood, I., Imadi, S. R., Shazadi, K., Gul, A., & Hakeem, K. R. (2016). Effects of Pesticides
471 on Environment. In K. R. Hakeem, M. S. Akhtar, & S. N. A. Abdullah (Eds.), *Plant, Soil*
472 *and Microbes: Volume 1: Implications in Crop Science* (pp. 253–269). Springer International
473 Publishing. https://doi.org/10.1007/978-3-319-27455-3_13

474 Mallatt, J., Blatt, M. R., Draguhn, A., Robinson, D. G., & Taiz, L. (2021). Debunking a myth: Plant
475 consciousness. *Protoplasma*, 258(3), 459–476. <https://doi.org/10.1007/s00709-020-01579-w>

476 Nicholson, E., Collen, B., Barausse, A., Blanchard, J. L., Costelloe, B. T., Sullivan, K. M. E.,
 477 Underwood, F. M., Burn, R. W., Fritz, S., Jones, J. P. G., McRae, L., Possingham, H. P., & Milner-
 478 Gulland, E. J. (2012). Making Robust Policy Decisions Using Global Biodiversity Indicators.
 479 *PLOS ONE*, 7(7), e41128. <https://doi.org/10.1371/journal.pone.0041128>
 480 OHHLEP, Adisasmito, W. B., Almuhairi, S., Behraves, C. B., Bilivogui, P., Bukachi, S. A.,
 481 Casas, N., Becerra, N. C., Charron, D. F., Chaudhary, A., Zanella, J. R. C., Cunningham, A. A.,
 482 Dar, O., Debnath, N., Dungu, B., Farag, E., Gao, G. F., Hayman, D. T. S., Khaita, M., ... Zhou,
 483 L. (2022). One Health: A new definition for a sustainable and healthy future. *PLOS Pathogens*,
 484 18(6), e1010537. <https://doi.org/10.1371/journal.ppat.1010537>
 485 Poisot, T., Becker, D. J., Catchen, M. D., Gibb, R., Shimabukuro, P. H. F., & Carlson, C. J. (2025).
 486 Biodiversity science and biosurveillance are fellow travelers. *BioScience*, b1af091. <https://doi.org/10.1093/biosci/b1af091>
 487
 488 Quadripartite organizations. (2022). *One Health Joint Plan of Action (2022-2026): Working*
 489 *together for the health of humans, animals, plants and the environment*. World Health Organi-
 490 zation.
 491 Rabinowitz, P., & Conti, L. (2013). Links among human health, animal health, and ecosys-
 492 tem health. *Annual Review of Public Health*, 34, 189–204. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-publhealth-031912-114426)
 493 [publhealth-031912-114426](https://doi.org/10.1146/annurev-publhealth-031912-114426)
 494 Rapport, D. J. (1989). What Constitutes Ecosystem Health? *Perspectives in Biology and Medi-*
 495 *cine*, 33(1), 120–132.
 496 Robinson, J. M., Breed, A. C., Camargo, A., Redvers, N., & Breed, M. F. (2024). Biodiversity and
 497 human health: A scoping review and examples of underrepresented linkages. *Environmental*
 498 *Research*, 246, 118115. <https://doi.org/10.1016/j.envres.2024.118115>
 499 Schaeffer, D. J., Herricks, E. E., & Kerster, H. W. (1988). Ecosystem health: I. Measuring ecosys-
 500 tem health. *Environmental Management*, 12(4), 445–455. <https://doi.org/10.1007/BF01873258>
 501 Strange, R. N., & Scott, P. R. (2005). Plant Disease: A Threat to Global Food Security. *Annual*
 502 *Review of Phytopathology*, 43(Volume 43, 2005), 83–116. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-phyto.43.113004.133839)
 503 [phyto.43.113004.133839](https://doi.org/10.1146/annurev-phyto.43.113004.133839)
 504 UNEP. (2022a). *Kunming-Montreal Global Biodiversity Framework*. United Nations Environ-
 505 ment Programme. Decision no. CBD/COP/DEC/15/4.
 506 UNEP. (2022b). *Mechanisms for planning, monitoring, reporting and review*. United Nations
 507 Environment Programme. Decision no. CBD/COP/DEC/15/6.

508 UNEP. (2024). *Global Action Plan on Biodiversity and Health*. United Nations Environment
 509 Programme. Decision no. CBD/COP/DEC/16/19.
 510 UNEP. (2025). *Monitoring framework for the Kunming-Montreal Global Biodiversity Framework*.
 511 United Nations Environment Programme. Decision no. CBD/COP/16/L.26/Rev.1.
 512 UNEP-WCMC. (2025). Indicators for the Kunming – Montreal Global Biodiversity Framework.
 513 In *Indicator Repository*. <https://www.gbif-indicators.org/>.
 514 Webb, L. E., Veenhoven, R., Harfeld, J. L., & Jensen, M. B. (2019). What is animal happiness?
 515 *Annals of the New York Academy of Sciences*, 1438(1), 62–76. <https://doi.org/10.1111/nyas.13983>
 516 WHO. (1948). Preamble to the Constitution of the World Health Organization as adopted by
 517 the International Health Conference. New York, 19-22 June, 1946; signed on 22 July 1946 by
 518 the representatives of 61 States (Official Records of the World Health Organization, no. 2, p.
 519 100) and entered into force on 7 April 1948. In *Constitution of the World Health Organization*.
 520 WHO. (2025). *WHO Pandemic Agreement*. Seventy-eighth World Health Assembly. WHA78.1.
 521 Winkler, A. S., Brux, C. M., Carabin, H., Neves, C. G. das, Häsler, B., Zinsstag, J., Fèvre, E.
 522 M., Okello, A., Laing, G., Harrison, W. E., Pöntinen, A. K., Huber, A., Ruckert, A., Natterson-
 523 Horowitz, B., Abela, B., Aenishaenslin, C., Heymann, D. L., Rødland, E. K., Berthe, F. C. J., ...
 524 Amuasi, J. H. (2025). The Lancet One Health Commission: Harnessing our interconnectedness
 525 for equitable, sustainable, and healthy socioecological systems. *The Lancet*, 406(10502), 501–
 526 570. [https://doi.org/10.1016/S0140-6736\(25\)00627-0](https://doi.org/10.1016/S0140-6736(25)00627-0)
 527 Zhang, X.-X., Liu, J.-S., Han, L.-F., Xia, S., Li, S.-Z., Li, O. Y., Kassegne, K., Li, M., Yin, K., Hu,
 528 Q.-Q., Xiu, L.-S., Zhu, Y.-Z., Huang, L.-Y., Wang, X.-C., Zhang, Y., Zhao, H.-Q., Yin, J.-X., Jiang,
 529 T.-G., Li, Q., ... Zhou, X.-N. (2022). Towards a global One Health index: A potential assessment
 530 tool for One Health performance. *Infectious Diseases of Poverty*, 11(1), 57. [https://doi.org/10.](https://doi.org/10.1186/s40249-022-00979-9)
 531 1186/s40249-022-00979-9