Monitoring biodiversity for human, animal, plant and environmental health

Francis Banville[1], Colin Carlson[2], Elodie Eiffener[3], Gabriel Munoz Acevedo[4], Andrea Paz Velez[1] and Timothée Poisot[1]

[1] Université de Montréal, Département de sciences biologiques, Montréal Québec, Canada; [2] Yale University, Yale School of Public Health, New Haven Connecticut, USA; [3] Karolinska Institutet, Stockholm , Sweden; [4] Canadian Institute for Health Information, Ottawa Ontario, Canada

Correspondence to Francis Banville — francis.banville@umontreal.ca

Abstract: The One Health approach promotes collaboration across disciplines to enhance the health of humans, animals, plants, and the environment. The Quadripartite organizations, which include the Food and Agriculture Organization of the United Nations, the United Nations Environment Programme, the World Organisation for Animal Health, and the World Health Organization, developed the One Health Joint Plan of Action (OH JPA) to support countries in achieving One Health. This plan consists of six action tracks, each consisting of a set of actions for implementing One Health. By requiring knowledge on zoonotic diseases (tracks 2 and 3), food and agriculture (track 4), antimicrobial resistance (track 5), and environmental health (track 6), most of these tracks directly concern biodiversity. However, there are currently no indicators for monitoring the OH JPA. Our research examines the extent to which all six tracks are covered by the Kunming-Montreal Global Biodiversity Framework (KM-GBF) of the Convention on Biological Diversity (CBD), which contains many indicators at the intersection of biodiversity and health. We assessed (1) the link between each indicator of the KM-GBF and human, animal, plant, and environmental health and (2) the usability of these indicators for monitoring One Health actions. We found that 75% of indicators are associated with health, and that a similar proportion can be used for monitoring One Health actions. Overall, our work aims to strengthen collaboration between the CBD Secretariat and the Quadripartite Organizations by highlighting the need for shared data, policy, and governance practices.

Keywords: biodiversity indicators, Kunming-Montreal Global Biodiversity Framework, One Health, One Health Joint Plan of Action, Quadripartite Organizations

Introduction

There are multiple pathways through which human health depends on biodiversity (Robinson et al., 2024). On one hand, nature's contributions to people provide important ecosystem services, such as shelter and nutritious food, that are vital to human health. On the other hand, diseases originating from wildlife can be a major threat to public health. The health of animals, plants, and the wider environment greatly impacts the capacity of ecosystems to provide services and prevent disease emergence (**Rabinowitz2013Links?**). The relationship between biodiversity and human health is far more significant when we consider that they can both be affected by the same threats (e.g., climate change and air pollution), and benefit from the same actions that mitigate these threats. The One Health approach recognizes these interconnections between human, animal, plant, and environmental health and provides a set of principes for achieving a healthier planet by fostering collaboration across disciplines.

The One Health Joint Plan of Action (OH JPA) is an action-oriented framework aimed at advancing One Health principles in the horizon of 2026. Developed by the Quadripartite organizations, which is a strategic One Health partnership between the Food and Agriculture Organization of the United Nations, the United Nations Environment Programme, the World Organisation for Animal Health, and the World Health Organization, the OH JPA contains 19 actions that address important challenges at the interface between human, animal, plant, and environmental health. These actions form six action tracks that collectively seek to strengthen health systems and food safety capacities while preventing zoonotic diseases, antimicrobial resistance, and environmental degradation. However, while this plan does a good job at promoting capacity-building across many dimensions of One Health, there are currently no indicators that would help us monitor its implementation. As we enter the final year of the plan, the need to evaluate the progress made since its adoption in 2022 grows acute, especially as this evaluation can inform the development of the next version of the plan.

In 2022, in their fifteenth conference, the Parties to the Convention of Biological Diversity (CBD) adopted the Kunming-Montreal Global Biodiversity Framework (KM-GBF) to alt and reverse biodiversity loss by 2030. To achieve this, the KM-GBF contains 4 general goals and 23 specific targets, ranging from the restoration of degraded ecosystems to the integration of biodiversity in decision-making. The KM-GBF Monitoring Framework includes 204 indicators that countries can use to monitor these goals and targets (UNEP-WCMC, 2025). These indicators are classified in four groups: headline (high-level indicators), binary (yes/no questions),

component (technical indicators), and complementary (supporting indicators). Two of these groups (headline and binary indicators) are mandatory, i.e. that countries have the legal obligation to include in their national reports. Measuring these indicators requires significant resources and expertise, which should therefore be well allocated.

More recently, the Parties to the CBD also adopted the Global Action Plan on Biodiversity and Health (GAPBH, UNEP, 2024), which formally recognizes the links between biodiversity and Health. The GAPBH groups the targets of the KM-GBF into 14 thematic categories, ranging from Access and benefit-sharing to Knowledge and engagement of people. Each thematic category has its own relevance to Health. For example, the targets that fall under the Mainstreaming category contribute to "the consideration of biodiversity and health interlinkages in decision-making across all sectors [which] can improve awareness of the benefits of biodiversity to foster more equitable health systems." Because of the high relevance of KM-GBF targets to Health, we hypothesized that the indicators of these targets will also be of relevance to Health.

This study aims to identify the indicators of the KM-GBF Monitoring Framework that are of relevance to One Health. First, we assessed the link between each of these indicators and human, animal, plant, and environmental health. We found that 75% of indicators are directly or indirectly connected to at least one of the four pillars of One Health. This result reinforces the need for collaboration between biologists, who are experts of these indicators, and health professionals, who need information at the intersection of biodiversity and health. Second, we evaluated the usability of the KM-GBF indicators for monitoring the specific actions of the OH JPA. We found that 91% of indicators can be used, either directly or after adaptation, to monitor actions in at least one action track. Reusing indicators can greatly reduce the workload of countries and their need to develop and monitor new indicators at the intersection of human, animal, plant, and environmental health. These results highlight the need for data sharing and knowledge transfer practices between the CBD Secretariat and the Quadripartite organizations.

Methods

We qualitatively assessed the relevance of all 204 KM-GBF indicators (Conference of the Parties to the Convention on Biological Diversity, 2025) to Health. Specifically, we evaluated the degree to which they were linked to each of the four pillars of One Health (i.e., to human, animal, plant, and environmental heath), as well as their usability for monitoring the

OH JPA. Our assessments were informed by the metadata (rationale, definition, method of computation, data sources, and scale of use) of the indicators provided by the World Conservation Monitoring Centre of the UN Environment Programme (UNEP-WCMC, 2025), when available, looking for explicit considerations of Health. When such metadata was unavailable, assessments were entirely based on expert judgment. The result of this analysis and the code to reproduce the figures are available on Zenodo.

Assessing the link between indicators and Health

Assessing the link between the KM-GBF indicators and the four pillars of One Health requires a working definition of human, animal, plant, and environmental health. We defined each of these groups of species based on the main objects of interest of the Quadripartite organizations. Humans are their own pillar and the primary focus of the WHO. We considered pets, livestock, and edible marine and freshwater species captured from fisheries and aquaculture in the animal health pillar. These are species mainly looked after by veterinarians and food inspectors and that are of core concern to the WOAH and FAO. Even though humans and wildlife are also animals, we excluded them from the animal health pillar. We included all cultivated plants used for food, fuel, and medicine into the plant health pillar, i.e. plant species that are the primary focus of the FAO. Wildlife was considered in the environment health pillar, alongside all other species not included in the other three pillars. The environment health pillar also includes whole ecosystems (natural and artificial), as well as tree species used in forestry and wild fish species. It is the primary focus of the UNEP. These definitions are alined with the core contributing sciences of the One Health approach (Gibbs, 2014; Lerner & Berg, 2017).

We used a broad definition of Health that considers the overall well-being of individuals and their capacity to function normally, which includes the absence of diseases (WHO, 1948). In humans and domestic and wildlife animals, well-being is the ability for an individual to satisfy its physical, mental, and behavioral needs (Webb et al., 2019), whereas diseases are undesirable conditions often leading to pain, suffering, or death. In plants, which are not conscious organisms (Mallatt et al., 2021), health is more ill-defined (Döring et al., 2012). Here, we considered plant health as the extent to which an individual is able to function physiologically, and plant diseases as impediments to their normal physiological functioning often leading to death. Similarly, when it comes to whole ecosystems, consisting of many individuals and their interactions, health is also ill-defined (Schaeffer et al., 1988). We defined environmental health as the extent to which an ecosystem can maintain its biological and chemical processes and adapt

to changes. Disturbances are the ecosystem analogue of diseases, and represent degradations that usually lead to a decline in ecosystem functioning. When assessing environmental health, we also considered the health of species included in the environment pillar of One Health, e.g. wildlife and uncultivated plant species. These descriptions guided us in our assessments.

We classified the degree to which the indicators are linked to each of the four pillars of One Health into four categories: direct connection, indirect connection, potential connection, and no connection. An indicator was qualified as directly connected to Health when there is a direct causal relationship between the indicator and Health. This is the case when the indicator directly measures a condition or determinant of health. For example, headline indicator 7.2 "Pesticide environment concentration and/or aggregated total applied toxicity" was considered directly connected to human health because of the toxicity of many pesticides under high level of exposure (Kim et al., 2017). An indicator was qualified as indirectly connected to Health when there is an intermediary factor between the indicator and Health. For instance, component indicator 4.CT.1 "Number of plant and animal genetic resources for food and agriculture secured in either medium- or long-term conservation facilities" was considered indirectly connected to human health because conserving genetic resources increases the resilience of food systems to global disasters (Esquinas-Alcázar, 2005), which in turn protects human health. An indicator was qualified as potentially connected to Health when there are multiple intermediary factors between the indicator and Health, or when its connection to Health is likely but difficult to certify. For example, complementary indicator 2.CY.2 "Proportion of key biodiversity areas in favourable condition" was qualified as potentially connected to human health because key biodiversity areas are mainly defined with ecological criteria for conserving rare or endemic species and pristine habitats (Butchart et al., n.d.), and may or may not include areas of high importance for human health. Finally, an indicator was qualified as not connected to Health when the connection is unlikely or absent. This was the case for component indicator 21.CT.3 "Index of linguistic diversity", for which we did not find any evidence of a link to human, animal, plant or environmental health. For each of the four pillars of One Health, we counted the number of indicators assigned to each category.

Assessing the usability of indicators for monitoring the OH JPA

We evaluated the usability of the KM-GBF indicators for monitoring the OH JPA by judging if they can likely detect trends relevant to the actions of the plan. For each action track, we classified the usability of the indicators into three categories: directly usable, usable after

adaptation, and not usable. Directly usable indicators can be used in their current form to monitor at least one action in an action track. They do not require adaptations. For example, we qualified component indicator 4.CT.4 "Proportion of local breeds classified as being at risk of extinction" as being directly usable for monitoring the fourth action track because it directly measures food safety risks. In contrast, indicators usable after adaptation need to be slightly modified before they can be used to monitor actions in an action track. These adaptations should be small changes in the scale of measurement, the data resolution, or the taxa monitored by the indicator. For instance, we qualified headline indicator 9.1 "Benefits from the sustainable use of wild species" as being usable after adaptation for monitoring the second action track on zoonotic epidemics and pandemics. To be more relevant for this action track, this indicator could refer more specifically to wildlife reservoirs of zoonotic agents instead of all wild species. Finally, not usable indicators are outside the scope of an action track or need to be greatly modified before being used to monitor an action track. For example, we qualified component indicator 4.CT.3 "Human-wildlife conflict indicator" as being not usable for monitoring the fifth action track on antimicrobial resistance.

We assessed usability for each action track independently, i.e. an indicator can be useful for monitoring multiple action tracks. For each action track, we counted the total number of indicators in each category. In addition, we counted the number of usable indicators for the whole plan regardless of the action tracks. Then, we partitioned these numbers based on the group of indicator in the KM-GBF Monitoring Framework (i.e. headline, binary, component, and complementary indicators) and the thematic categories of the GAPBH. Finally, for each action track where an indicator was considered usable (either directly or after adaptation), we identified the most relevant action that can be monitored by the indicator, and counted the number of indicators associated with each action.

Results and discussion

Link between indicators and Health

We found that 75% of KM-GBF indicators are either directly or indirectly connected to human, animal, plant, or environmental health, with approximately 55% of indicators being directly connected to Health. This indicates that monitoring the state, benefits, and pressure of biodiversity and our responses to biodiversity loss gives us a lot of information on One Health. In Figure 1, we show the number of indicators that are directly, indirectly, potentially, and



Figure 1: **Number of KM-GBF indicators linked with human, animal, plant, and environmental health.** Each indicator was classified as being either directly connected (blue), indirectly connected (green), potentially connected (orange), or not connected (red) to each of the four pillars of One Health.

not connected to each of the four pillars of One Health. The number of directly connected indicators is slightly higher for environmental health (41% of indicators) compared to human (27%), animal (26%), and plant (28%) health. However, this difference shrinks after adding indirectly connected indicators (63% for environmental health compared to 56%, 53%, and 51% respectively for human, animal, and plant health). This suggests that the four pillars of One Health can all be meaningfully informed by biodiversity monitoring.

The fact that most KM-GBF indicators are connected to Health is not surprising. The GAPBH recognizes the relevance to Health of all 23 targets of the KM-GBF, and because the indicators measure the progress made towards these targets, they also measure the progress made towards a healthier planet. However, we did not expect that comparable amounts of indicators would monitor the four pillars of One Health. This indicates that there are virtually no biases in the amount of information that can be reused to monitor human, animal, plant, and environmental health. In other words, we monitor all core components of One Health when we monitor biodiversity.

Usability of indicators for monitoring the OH JPA

The vast majority of KM-GBF indicators (91%) can be used either directly or after adaptation to monitor actions in the OH JPA, and around 74% of indicators can be used directly. This signifies that we can already do an extensive evaluation of One Health actions with existing data and methodologies. However, we show in Figure 2 that the usability of indicators greatly differs depending on the action track. The proportion of usable indicators varies from 20% for the fifth action track on antimicrobial resistance (4% of directly usable indicators) to 79% for the sixth action track on the integration of the environment into One Health (62% of directly usable indicators). The high proportion of usable indicators for the sixth action track is not surprising given that the KM-GBF is a multilateral environmental agreement that aims to protect species and restaure ecosystems. The low proportion of indicators that can monitor actions in the fifth action track could be due to the characteristics of these actions, which are more focused on collaboration, capacity-building and awareness-raising than the decreasing and prevention of antimicrobial resistance through biodiversity. Similarly, the actions in the third action track on endemic zoonotic, neglected tropical, and vector-borne diseases, which can only be monitored by 33% of the indicators, are more centered on capacity-building and less on the drivers and inhibitors of infectious diseases, especially in comparison to the actions of the second action track on zoonotic epidemics and pandemics. These differences suggest that, although the KM-GBF Monitoring Framework can greatly reduce the workload of countries by reducing the need to develop new indicators at the intersection of biodiversity and health, additional work is still needed to equally cover all aspects of the plan.

The number of indicators associated with each action is presented in Table 1. There are high discrepancies in the usability of indicators between actions. The total number of usable indicators (either directly or after adaptation) varies from 1 (action 5.3: Strengthen global AMR governance structures) to 108 (action 6.1: Protect, restore, and prevent the degradation of ecosystems and the wider environment). Within a single action track, the biggest differences are found within the sixth action track, where only 12 indicators were associated to action 6.4 whereas 108 indicators were associated to action 6.1 This result shows that there is substantial information available for some of the actions, but it also further highlights the need to identify or develop new indicators for specific actions of the plan, even within the action tracks that can be monitored by numerous indicators.

Figure 3 shows the proportion of indicators that can be used to monitor each action track, for each group of indicators in the KM-GBF Monitoring Framework. We evaluated that 100%



Figure 2: **Number of KM-GBF indicators that can be used to monitor at least one action in each of the OH JPA action tracks.** The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The proportion of usable indicators (either directly or after adaptation) for each action track is indicated. The bottom bar and proportion designate the indicators that can monitor at least one action in the whole plan.

of headline indicators (27/27), 93% of binary indicators (14/15), 94% of component indicators (49/52), and 87% of complementary indicators (96/110) can be used, either directly or after adaptation, to monitor at least one action track. Headline and binary indicators are the only two groups of indicators that are mandatory, i.e. that Parties have the legal obligation to report on. The high proportion of headline and binary indicators is encouraging, because it indicates that the information most likely to figure in National Biodiversity Strategies and Action Plans (NBSAPs) will be reusable for evaluating the implementation of the OH JPA. On the other hand, the high reusability of component and complementary indicators, which are currently optional, should be seen as an additional incitative to measure them.

In Figure 4, we present the usability of indicators for each thematic category of the GAPBH. In every category, at least 89% of indicators can be used, either directly or after adaptation, for monitoring at least one action track, excepting in the Knowledge and engagement of people category, which only has 52% of usable indicators. This suggests that every sector of the KM-

Table 1: **Number of KM-GBF indicators that can be used directly and after adaptation to monitor each action of the OH JPA.** The total number of usable indicators for each action track is indicated in the highlighted blue lines. A specific indicator can be used to monitor multiple action tracks, but only the most relevant action in each action track was identified for each indicator.

Action	Directly usable	Usable after adaptation	Total usable
1 Enhancing One Health capacities to strengthen health systems	**46**	**61**	**107**
1.1 Establish the foundations for One Health capacities	30	5	35
1.2 Generate mechanisms, tools, and capacities to establish a One Health competent workforce and the frameworks/processes to facilitate One Health work	8	50	58
1.3 Generate an enabling environment for the effective implementation of One Health	8	6	14
2 Reducing the risks from emerging and re-emerging zoonotic epidemics and pandemics	**40**	**63**	**103**
2.1 Understand the drivers of emergence, spillover, and spread of zoonotic pathogens	16	27	43
2.2 Identify and prioritize targeted, evidence-based upstream interventions to prevent the emergence, spillover, and spread of zoonotic pathogens	7	6	13
2.3 Strengthen national, regional, and global One Health surveillance, early warning, and response systems	17	30	47
3 Controlling and eliminating endemic zoonotic, neglected tropical and vector-borne diseases	**25**	**43**	**68**
3.1 Enable countries to develop and implement community-centric and risk-based solutions to endemic zoonotic, neglected tropical, and vector-borne disease control using a One Health approach involving all relevant stakeholders	3	14	17
3.2 Ensure the harmonized application of One Health principles at all levels by implementing practical measures to strengthen local, national, regional, and global policy frameworks for the control and prevention of endemic zoonotic, neglected tropical, and vector-borne diseases	18	10	28
3.3 Increase political commitment and investment in the control of endemic zoonotic, neglected tropical, and vector-borne diseases, by advocating for and demonstrating the value of a One Health approach	4	19	23
4 Strengthening the assessment, management and communication of food safety risks	**42**	**70**	**112**
4.1 Strengthen the One Health approach in national food control systems and food safety coordination	18	30	48
4.2 Utilize and improve food systems data and analysis, scientific evidence, and risk assessment in developing policy and making integrated risk management decisions	15	21	36
4.3 Foster the adoption of the One Health approach in national foodborne disease surveillance systems and research for the detection and monitoring of foodborne disease and food contamination	9	19	28
5 Curbing the silent pandemic of antimicrobial resistance (AMR)	**9**	**31**	**40**
5.1 Strengthen the capacity and knowledge of countries to prioritize and implement context-specific collaborative One Health work to control AMR in policy, legislation, and practice	7	17	24
5.2 Reinforce global and regional initiatives and programmes to influence and support One Health responses to AMR	2	13	15
5.3 Strengthen global AMR governance structures	0	1	1
6 Integrating the environment into One Health	**126**	**36**	**162**
6.1 Protect, restore, and prevent the degradation of ecosystems and the wider environment	92	16	108
6.2 Mainstream the health of the environment and ecosystems into the One Health approach	9	7	16
6.3 Integrate environmental knowledge, data, and evidence into One Health decision-making	20	6	26
6.4 Create an interoperable One Health academic and in-service training programme for environmental, medical, agricultural, and veterinary sector professionals	5	7	12

GBF Monitoring Framework, from Nature's contribution to people to Species management, is relevant for One Health, even though there are wide variations in the action track most relevant for each category. This figure also shows that all indicators that can directly monitor



Figure 3: **Proportion of KM-GBF indicators that can be used to monitor at least one action in each of the OH JPA action tracks, partitioned by the different groups of indicators.** The number of indicators in each group is indicated in parentheses. The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The gray bars designate the indicators that can monitor at least one action in the whole plan.

the fifth action track on antimicrobial resistance are in the Access and benefit-sharing and Biosafety and biotechnology categories, further emphasizing the importance of collaboration, capacity-building and awareness-raising set forth in this action track.

Conclusion

- Monitoring Frameworks
  - The monitoring framework of the OH JPA can be based on indicators of the KM-GBF
  - Importance of reusing indicators to decrease workload on countries
  - Importance of sharing data, methodologies, and expertise
  - Sharing policy and governance practices
- Gaps in indicators
  - Indicators

Figure 4: **Proportion of KM-GBF indicators that can be used to monitor at least one action in each of the OH JPA action tracks, partitioned by the different thematic categories of the GAPBH.** The number of indicators in each thematic category is indicated in parentheses. The thematic category of an indicator was assigned based on the thematic category of its target. Indicators that monitor KM-GBF goals A, B, C, and D are not assigned to any thematic category. The opaque bars represent directly usable indicators, whereas the transparent bars represent those usable after adaptation. The gray bars designate the indicators that can monitor at least one action in the whole plan.

- Other indicators could be identified in other monitoring frameworks (e.g. SDG)
  - New indicators can be developed after identifying gaps

Acknowledgments

We acknowledge that this study was conducted on land within the traditional unceded territory of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat and Omàmiwininiwak nations. FB and TP were supported by the Wellcome Trust. A special thanks to the GEO BON One Health working group and Claire Burnel for their useful feedback on the manuscript.

References

Butchart, S. H. M., Crowe, O., Scott, T., Plumptre, A. J., Eldred, M., McKee, E., Waliczky, Z., Hasani, I. A., Azpiroz, A. B., Baisero, D., Barasa, F., Beresford, A. E., Bezeng, B. S., Boyd, C., Brooks, T. M., Buchanan, G., Bunting, G., Capellan, S., Cox, N., … Upgren, A. (n.d.). Extent, characteristics and policy applications of Key Biodiversity Areas. *Biological Reviews*, *n/a*(n/a). https://doi.org/10.1002/brv.70144

Conference of the Parties to the Convention on Biological Diversity. (2025). *Monitoring framework for the Kunming-Montreal Global Biodiversity Framework*. United Nations Environment Programme. Decision no. CBD/COP/16/L.26/Rev.1.

Döring, T. F., Pautasso, M., Finckh, M. R., & Wolfe, M. S. (2012). Concepts of plant health – reviewing and challenging the foundations of plant protection. *Plant Pathology*, *61*(1), 1–15. https://doi.org/10.1111/j.1365-3059.2011.02501.x

Esquinas-Alcázar, J. (2005). Protecting crop genetic diversity for food security: Political, ethical and technical challenges. *Nature Reviews Genetics*, *6*(12), 946–953. https://doi.org/10.1038/nrg1729

Gibbs, E. P. J. (2014). The evolution of One Health: A decade of progress and challenges for the future. *Veterinary Record*, *174*(4), 85–91. https://doi.org/10.1136/vr.g143

Kim, K.-H., Kabir, E., & Jahan, S. A. (2017). Exposure to pesticides and the associated human health effects. *Science of The Total Environment*, *575*, 525–535. https://doi.org/10.1016/j.scitotenv.2016.09.009

Lerner, H., & Berg, C. (2017). A Comparison of Three Holistic Approaches to Health: One Health, EcoHealth, and Planetary Health. *Frontiers in Veterinary Science*, *4*, 163. https://doi.org/10.3389/fvets.2017.00163

Mallatt, J., Blatt, M. R., Draguhn, A., Robinson, D. G., & Taiz, L. (2021). Debunking a myth: Plant consciousness. *Protoplasma*, *258*(3), 459–476. https://doi.org/10.1007/s00709-020-01579-w

Robinson, J. M., Breed, A. C., Camargo, A., Redvers, N., & Breed, M. F. (2024). Biodiversity and human health: A scoping review and examples of underrepresented linkages. *Environmental Research*, *246*, 118115. https://doi.org/10.1016/j.envres.2024.118115

Schaeffer, D. J., Herricks, E. E., & Kerster, H. W. (1988). Ecosystem health: I. Measuring ecosystem health. *Environmental Management*, *12*(4), 445–455. https://doi.org/10.1007/BF01873258

UNEP. (2024). *Global Action Plan on Biodiversity and Health*. https://www.cbd.int/health/GAP.shtml.

UNEP-WCMC. (2025). Indicators for the Kunming – Montreal Global Biodiversity Framework. In *Indicator Repository*. https://www.gbf-indicators.org/.

Webb, L. E., Veenhoven, R., Harfeld, J. L., & Jensen, M. B. (2019). What is animal happiness? *Annals of the New York Academy of Sciences*, *1438*(1), 62–76. https://doi.org/10.1111/nyas.13983

WHO. (1948). Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference. New York, 19-22 June, 1946; signed on 22 July 1946 by the representatives of 61 States (Official Records of the World Health Organization, no. 2, p. 100) and entered into force on 7 April 1948. In *Constitution of the World Health Organization*.