

Deciphering probabilistic species interaction networks

Francis Banville^{1,2,3} Tanya Strydom^{1,3} Penelope S. A. Blyth⁴ Chris Brimacombe⁵ Michael Catchen^{3,6}
Gabriel Dansereau^{1,3} Dominique Gravel^{2,3} Gracielle Higino² Thomas Malpas⁴ Hana Mayall⁴
Kari Norman¹ Timothée Poisot^{1,3}

¹ Université de Montréal ² Université de Sherbrooke ³ Quebec Centre for Biodiversity Science ⁴ University of Sheffield ⁵ University of Toronto ⁶ McGill University

Correspondance to:

Francis Banville — francis.banville@umontreal.ca

Representing species interactions probabilistically (how likely are they to occur?) as opposed to deterministically (are they occurring?) expresses uncertainties in our knowledge of interactions. The sources of uncertainty captured by interaction probabilities depend on the method used to evaluate them: uncertainty of predictive models, subjective assessment of experts, or empirical measurement of interaction spatiotemporal variability. However, guidelines for the estimation and documentation of probabilistic interaction data are still lacking. This is concerning because our understanding and analysis of interaction probabilities depend on their sometimes elusive definition and uncertainty sources. We review how probabilistic interactions are defined at different spatial scales, from local interactions to regional networks (metawebs), with a strong emphasis on host-parasite and trophic (predatory and herbivory) interactions. These definitions are based on the distinction between the realization of an interaction at a specific time and space (local) and its biological feasibility (regional). Through case studies using host-parasite interactions in Europe, we illustrate how these two network representations differ in their statistical properties, specifically: how local networks and metawebs differ in their spatial and temporal scaling of probabilistic interactions, but not in their taxonomic scaling. We present two approaches to inferring binary interactions from probabilistic ones that account for these differences and show that systematic biases arise when directly inferring local networks from metawebs. Our results underscore the importance of more rigorous descriptions of probabilistic species interaction networks that specify their type of interaction (local or regional) and uncertainty sources.

This work is released by its authors under a CC-BY 4.0 license



Last revision: April 22, 2024

1 Introduction

2 As we try to navigate global biodiversity change, filling in knowledge gaps about biodiversity becomes
3 instrumental to monitoring and mitigating those changes (Abrego *et al.* 2021; Gonzalez & Londoño 2022;
4 Hortal *et al.* 2015). However, cataloging species, populations and, in particular, ecological interactions (e.g.,
5 predation, parasitism, and pollination) is a substantial challenge (Pascual *et al.* 2006; Polis 1991). There are
6 methodological and biological constraints that hinder our ability to document species interactions, inevitably
7 leading to uncertainty in our knowledge of interactions. For example, the spatial and temporal uncoupling of
8 species (e.g., nocturnal and diurnal species coexisting in the same space with different daily activity timings,
9 Jordano 1987) and the large number of rare and cryptic interactions in a community, contribute to these
10 knowledge gaps (Jordano 2016).

11 More generally, several conditions must be satisfied for an interaction to be observed locally. First, both species
12 must have overlapping geographic ranges, i.e. they must co-occur within the region of interest (Cazelles *et al.*
13 2016; Morales-Castilla *et al.* 2015). Second, they must have some probability of meeting (Poisot *et al.* 2015).
14 Probabilities of interspecific encounters are typically low, especially for rare species with low abundances.
15 While species' absolute abundances may impact interaction frequencies (Vázquez *et al.* 2007), encounter
16 probabilities are determined by their relative abundances (Canard *et al.* 2012). The probability that species
17 meet also depends on their biology, such as their phenology (Olesen *et al.* 2010; Singer & McBride 2012) and
18 discoverability (Broom & Ruxton 2005). Finally, when species do come into contact, an interaction occurs only
19 if their traits are locally compatible (Poisot *et al.* 2015), including their phenotypes (Bolnick *et al.* 2011; Gravel
20 *et al.* 2013; Stouffer *et al.* 2011) and behavior (Choh *et al.* 2012; Pulliam 1974). Interactions may also be
21 influenced by a third species (e.g., a more profitable prey species, Golubski & Abrams 2011; Sanders & van
22 Veen 2012). Documenting the location and timing of interactions becomes even more difficult when accounting
23 for the spatiotemporal variability of ecological interactions (Poisot *et al.* 2012, 2015). Environmental factors,
24 such as temperature (Angilletta *et al.* 2004), drought (Woodward *et al.* 2012), climate change (Araujo *et al.*
25 2011; Gilman *et al.* 2010; Woodward *et al.* 2010), and habitat modifications (Tylianakis *et al.* 2007), contribute
26 to this spatiotemporal variability by impacting species abundance and traits. Even under favorable
27 circumstances, there remains a possibility that the interaction does not occur locally, either due to the intricate
28 nature of the system or simply by chance. If it does occur, it might go undetected, particularly if it happens
29 infrequently. In this context, it is unsurprising that our knowledge of ecological interactions remains limited

30 (Hortal *et al.* 2015) despite extensive biodiversity data collection (Schmeller *et al.* 2015).

31 Knowing the biological capacity of two species to interact directly (via e.g., trophic interactions) is necessary

32 but not sufficient for inferring their interaction at a specific time and space. The recognition of the intrinsic

33 variability of species interactions has led ecologists to expand their representation of ecological networks to

34 include a probabilistic view of interactions (Dallas *et al.* 2017; Fu *et al.* 2021; Poisot *et al.* 2016). This allows

35 filling in the Eltonian shortfall (Hortal *et al.* 2015) by modeling the probability of occurrence of interactions

36 (e.g., Gravel *et al.* 2019), which can be an important tool for directing efforts and taking action (Carlson *et al.*

37 2021), especially in places where access and resources for research are scarce. The probabilistic representation

38 of interactions has been applied to direct interactions, which are conceptually and mathematically analogous

39 regardless of their biological type (e.g., predation and pollination). This is in contrast with indirect interactions

40 (e.g., interspecific competition), which arise from distinct ecological processes and are often not directly

41 observable (Kéfi *et al.* 2015, 2016).

42 Representing direct interactions probabilistically can capture the spatiotemporal variability of the

43 aforementioned ecological processes and the uncertainty in our knowledge of interactions. Networks of

44 probabilistic interactions, within a Bayesian perspective, express our degree of belief (or confidence) regarding

45 the occurrence or observation of interactions. In contrast, interactions are simply regarded as either occurring or

46 not in networks of deterministic binary interactions. Based on the scale at which they are estimated, interaction

47 probabilities may reflect our level of confidence in whether interactions will be observed, realized locally, or

48 biologically feasible. As an illustration, we could outline a situation in which there is a 50% certainty that an

49 interaction occurs 50% of the time. Our level of confidence should be more definitive (approaching either 0 or

50 1) as we extend our sampling to a broader area and over a longer time period, thereby diminishing the

51 uncertainty of our knowledge of interactions (but not necessarily the estimation of their variability). In the

52 broadest sense, binary interactions are also a type of probabilistic interaction, in which the numerical value of

53 an interaction is restrained to 0 (non-occurring) or 1 (occurring). In networks of probabilistic interactions, only

54 forbidden interactions (i.e., interactions prohibited by biological traits or species absence, Jordano *et al.* 2003;

55 Olesen *et al.* 2010) have a probability value of zero, provided that intraspecific trait variability is considered

56 (Gonzalez-Varo & Traveset 2016).

57 By accounting for the uncertainty of interactions, networks of probabilistic interactions may provide a more

58 realistic portrait of species interactions. However, how the uncertainty of pairwise interactions propagates to

59 network structure (i.e., community-level properties driving the functioning, dynamics, and resilience of

60 ecosystems, McCann 2007; McCann 2011; Proulx *et al.* 2005; Rooney & McCann 2012) remains to be
61 elucidated. The application and development of computational methods in network ecology, often based on a
62 probabilistic representation of interactions, can alleviate (and guide) the sampling efforts required to document
63 species interactions (Strydom *et al.* 2021). For example, statistical models can be used to estimate the
64 uncertainty of pairwise interactions (Cirtwill *et al.* 2019) and the probability of missing (false negatives) and
65 spurious (false positives) interactions (Guimerà & Sales-Pardo 2009). Statistical models can generate
66 predictions of ecological networks without prior knowledge of pairwise interactions. They may do so using
67 body size (Gravel *et al.* 2013; Petchey *et al.* 2008), phylogeny (Elmasri *et al.* 2020; Strydom *et al.* 2022), or a
68 combination of niche and neutral processes (Bartomeus *et al.* 2016; Pomeranz *et al.* 2019) for inference.
69 Topological null models, which generate networks of probabilistic interactions by preserving chosen
70 characteristics of the adjacency matrix of binary interactions while intentionally omitting others (Bascompte *et*
71 *al.* 2003; Fortuna & Bascompte 2006), are examples of common probabilistic interaction models. Null models
72 can be used to produce underlying distributions of network measures for null hypothesis significance testing.
73 Many measures have been developed to describe the structure (Poisot *et al.* 2016) and diversity (Godsoe *et al.*
74 2022; Ohlmann *et al.* 2019) of probabilistic interaction networks. These models and measures support the use
75 of this approach for the study of a wide range of ecological questions, from making better predictions of species
76 distribution (Cazelles *et al.* 2016) to forecasting the impact of climate change on ecological networks (Gilman
77 *et al.* 2010).
78 Yet, a precise definition of probabilistic interactions appears to be lacking, making the estimation and use of
79 these data more difficult. In this manuscript, we aim to take a step back by outlining different ways in which
80 probabilistic interactions are defined and used in network ecology. We distinguish two broad categories of
81 probabilistic interaction networks that necessitate distinct approaches: local networks describing probabilities
82 of realized interactions, and regional networks (metawebs) describing probabilities of potential interactions. We
83 highlight the distinctions in the ecological meaning of these two representations and examine some of their
84 properties and relationships (particularly with space, time, and between each other) through empirical case
85 studies. Moreover, the lack of clear guidelines on the use of probabilistic interaction data is worrisome, as it
86 affects both data producers and re-users who generate and manipulate these numbers. This is concerning
87 because sampling strategies and decisions regarding network construction can affect our understanding of
88 network properties (Brimacombe *et al.* 2023). There is currently no reporting standard that could guide the
89 documentation of all types of probabilistic interactions (although see e.g., Salim *et al.* 2022 who discuss data

90 standards for deterministic mutualistic networks). Well-defined reporting standards for probabilistic interactions
91 would support more adequate manipulation and integration of interaction data from different sources and guard
92 against possible misinterpretations arising from ambiguous definitions of probabilistic interaction networks.
93 This documentation should outline the nature (i.e., local or regional) and type (e.g., predatory or pollination) of
94 the interactions, provide information regarding the taxonomic level, identities, and characteristics (e.g., life
95 stages) of the individuals involved in an interaction, present the mathematical formulation of probabilities,
96 including clearly identified conditional variables (e.g., spatial and temporal scales), and describe the methods
97 and contexts (e.g., location, time, environmental conditions) in which interactions were estimated. Inadequately
98 documented probabilistic interaction data should be used with caution when analyzing ecological networks.
99 The broad principles underlying our findings remain relevant and applicable across diverse ecological contexts.

100 **Probabilistic representations of interactions**

101 Consider a scenario where an avian predator has just established itself in a northern habitat home to a small
102 rodent. Suppose these species have never co-occurred before, and as a result, their interaction has not been
103 previously observed. What is the probability that the rodent is part of the diet of the avian predator, or put
104 differently, what is the probability that they interact? Answering this question requires some clarification, as
105 there are multiple ways to interpret and calculate interaction probabilities. We could calculate the probability
106 that the traits of these species match, i.e. that the avian predator possesses the biological attributes to capture
107 and consume the rodent. We could also calculate the probability that their traits support an interaction under the
108 typical environmental conditions of the new habitat. For example, because avian predators hunt by sight,
109 predation could be possible in the absence of snow but highly improbable when snow is present, as rodents may
110 use it as a shelter to hide from predators. Finally, we could calculate the probability that the avian predator will
111 consume the rodent at *that* particular location, for which the spatial and temporal boundaries need to be
112 specified. The estimation of interaction probabilities, whether through predictive models or prior distributions,
113 hinges on our understanding of these probabilities and the specific ecological processes we aim to capture.

114 An important aspect to consider when using probabilities of interactions is knowing if they describe potential or
115 realized interactions, as these two types of interactions have distinct conceptual underpinnings and sources of
116 uncertainty. A potential (regional) interaction is defined as the biological capacity of two taxa to interact (i.e.,
117 the probability that they interact if they were to encounter each other and given sufficient time) whereas a

realized (local) interaction is the occurrence or observation of this interaction in a well-defined space and time (i.e., the probability that they interact locally). For two co-occurring taxa and over infinite time, the probability of local interaction is equivalent to the probability of regional (potential) interaction. We use the terms *metaweb* (Dunne 2006) to designate regional networks of potential interactions and *local networks* (Poisot *et al.* 2012) for those of realized interactions. Metawebs are the network analogs of the species pool, where local networks originate from a subset of both species (nodes) and interactions (edges) of the regional metaweb (Saravia *et al.* 2022). Without clear documentation, it can be challenging to know if published probabilistic interaction networks describe local or regional interactions (Tbl. 1 provides examples of studies employing both types of probabilistic interaction networks), or if so-called probabilities are in reality a form of interaction score. When probabilistic regional interactions are used and interpreted incorrectly as local interactions (and conversely), this may generate misleading findings during data analysis. We believe that a better understanding of probabilistic local and regional interactions would prevent interpretation errors (e.g., when studying network-area relationships with metawebs or local networks) and facilitate a more adequate use of interaction data. In the following sections, we delve into these distinctions as we scale up from pairwise interactions to higher-level representations of ecological networks, i.e. local networks and metawebs.

Pairwise interactions: the building blocks of ecological networks

Local networks and metawebs, like any type of network, are made of nodes and edges that may be represented at different levels of organization. The basic units of ecological networks are individuals that interact with each other (e.g., by predation in food webs, Elton 2001), forming individual-based networks (Melián *et al.* 2011). The aggregation of these individuals into more or less homogeneous groups (e.g., populations, species, families, feeding guilds) allows us to represent nodes at broader taxonomic scales, which affects our interpretation of the properties of these systems (Guimarães 2020; Hemprich-Bennett *et al.* 2021).

Ecologists have traditionally represented interactions (edges) as binary objects that were considered realized after observing at least one individual from group i interact with at least another individual from group j . In an adjacency matrix B of binary interactions, the presence or absence of an interaction $B_{i \rightarrow j}$ between two taxa can be viewed as the result of a Bernoulli trial $B_{i \rightarrow j} \sim \text{Bernoulli}(P(B_{i \rightarrow j}))$, with $P(B_{i \rightarrow j})$ being the probability of interaction. This interaction probability characterizes our limited ecological knowledge and/or the intrinsic spatiotemporal variability of interactions. It may be estimated through predictive models (e.g., those based on

146 biological traits and species abundances) or expert (prior) knowledge about the interaction. In networks of
147 probabilistic interactions, the edge values $P(B_{i \rightarrow j})$ are probabilistic events whose only two possible outcomes
148 are the presence ($B_{i \rightarrow j} = 1$) or absence ($B_{i \rightarrow j} = 0$) of an interaction between each pair of nodes. Depending on
149 the type of probabilistic interaction network (local or metaweb), the mathematical formulation and interpretation
150 of stochastic parameters like $P(B_{i \rightarrow j})$ can be linked to environmental and biological factors such as species
151 abundance, traits, area, and time, for example using logistic regression with continuous explanatory variables.

152 Predicting the number of local networks in which the interaction between two given taxa occurs can be achieved
153 by using a Binomial distribution, assuming a constant interaction probability and independence between
154 networks (trials). When considering uncertainties around the estimation of $P(B_{i \rightarrow j})$ or its spatiotemporal
155 variability, a Beta distribution may be used to represent the relative likelihood of different probability values.
156 For example, when calculating the probability of interaction between two taxa based on their local abundances,
157 any spatiotemporal fluctuations in their abundances would introduce variability in the interaction probability at
158 the local scale. If we take into account the stochasticity of the interaction probability, a Beta-Binomial
159 distribution can be used to predict the number of networks in which the interaction occurs. Empirically
160 observing an interaction between two taxa at a given location and time provides important information that can
161 be used to update previous estimates of $P(B_{i \rightarrow j})$, informing us on the biological capacity of both taxa to interact
162 and the environmental conditions that enabled them to interact locally.

163 Even though binary interaction networks constitute a highly valuable source of ecological information (Pascual
164 *et al.* 2006), they overlook interaction strengths. Represented in a quantitative adjacency matrix W , interaction
165 strengths better describe the energy flows, demographic impacts or frequencies of interactions between nodes
166 (Berlow *et al.* 2004; Borrett & Scharler 2019), with $W_{i \rightarrow j}$ being a natural \mathbb{N} or real \mathbb{R} number depending on the
167 measure. For example, they may represent local interaction rates between pairs of taxa (e.g., the flower-visiting
168 rates of pollinators in a mutualistic network, Herrera 1989). When interaction strengths characterize predation
169 pressure on prey, they can serve as parameters in a Lotka-Volterra model (e.g., Emmerson & Raffaelli 2004).
170 The extra amount of ecological information in quantitative networks typically comes at a cost of greater
171 sampling effort and data volume in predictive models (Strydom *et al.* 2021), which can lead to relatively high
172 levels of uncertainties when inferring quantitative networks with limited data.

173 As for binary interaction networks, the uncertainty and variability of interaction strengths can be represented
174 probabilistically. However, the need to estimate the probability distribution of interaction strengths makes the
175 inference of probabilities more challenging in quantitative networks compared to binary interaction networks.

176 Interaction strengths can follow many probability distributions depending on the method. For instance, they can
177 follow a Poisson distribution $W_{i \rightarrow j} \sim \text{Poisson}(\lambda_{i \rightarrow j})$ when predicting the number of interactions between pairs
178 of nodes, with $\lambda_{i \rightarrow j}$ being the expected rate at which individuals of taxa i and j interact (e.g., the average
179 number of prey j consumed by all predators i in a given time period). The Poisson distribution can also be
180 0-inflated after initially modeling non-interacting taxa (e.g., Boulangeat *et al.* 2012 employ a 0-inflated model
181 to analyze species abundance following the modeling of species presence and absence), which constitute the
182 majority of taxa pairs in most local networks due to their typically high sparseness (Jordano 2016). Because of
183 the methodological difficulties typically encountered when building deterministic quantitative networks (which
184 are only partially mitigated by models such as Ecopath, Plagányi & Butterworth 2004), binary interaction
185 networks, which are easier to sample (Jordano 2016) and predict (Strydom *et al.* 2021), have been more
186 frequently studied and modeled. Moreover, most published probabilistic interaction networks and methods
187 describe interactions whose outcome is binary (whether interaction probabilities are regarded as constant or
188 variable, e.g. represented by a Beta distribution), which underlines the need for better guidelines regarding the
189 interpretation and manipulation of these types of networks first. For these reasons, our primary focus is
190 interpreting and using interaction probabilities in Bernoulli distributions, in both local networks and metawebs.

191 Local networks: communities interacting in space and time

192 Local networks of probabilistic interactions describe how likely taxa are to interact at a given location and time
193 period. Local interactions are contingent upon the environmental conditions of the community and the matching
194 of taxa biological traits. In local networks, edges commonly represent our degree of belief that two taxa interact
195 in nature, but can also represent the probability of empirically *observing* this interaction (Catchen *et al.* 2023).
196 We define space as the collection of geographic coordinates (x, y, z) , with (x, y) representing longitude and
197 latitude coordinates, and z denoting either altitudes or depths. These point coordinates delineate the spatial
198 boundaries of the system, which may be portrayed as a polyhedron. Ecological interactions may exhibit
199 variations along latitudinal and altitudinal gradients, as evidenced by changes in hummingbird-plant
200 interactions (Weinstein & Graham 2017a, b) and mosquito biting rates (e.g., Kulkarni *et al.* 2006) at different
201 elevations. In contrast, time is defined as the specific time period within which interactions were either observed
202 or predicted. Even though space and time are continuous variables that should yield probability *densities* of
203 interactions (i.e., relative likelihoods of interactions occurring at infinitesimal locations and instants in time),

these definitions enable them to be conceptualized as distinct patches and time segments. Treating space and time as discrete dimensions aligns with the common sampling methods of ecological networks and provides probabilities of interactions, which can be obtained by integrating probability densities over space and time. Furthermore, we can quantify both an area A and a duration t , which can be readily used in spatiotemporal analyses of ecological networks. When studying network-area relationships (NAR, Galiana *et al.* 2018), we anticipate that local probabilities of interactions scale positively with area and duration because taxa have more opportunities to interact as these dimensions expand.

The probability that two taxa i and j interact in a local network $L_{x,y,z,t}$ (spatial and temporal subscripts hereafter omitted or replaced by the shorter subscript k for clarity) can be conditioned on many environmental and biological factors. In addition to network area (or volume) and duration, they may be conditioned on taxa co-occurrence $X_{i,j,k}$, which is usually a Boolean describing if the geographic distributions of both taxa overlap within the study area. As illustrated in Box 1, co-occurrence may be modeled probabilistically, in which case it may conform to a Bernoulli distribution $X_{i,j,k} \sim \text{Bernoulli}(P(X_{i,k}, X_{j,k}))$, where $X_{i,k}$ and $X_{j,k}$ are the local occurrences of both taxa. The probability of co-occurrence $P(X_{i,k}, X_{j,k})$ can be estimated through the application of joint species distribution models (e.g., Pollock *et al.* 2014), potentially taking into account biotic interactions (Staniczenko *et al.* 2017). Given that the probability that two non-co-occurring taxa interact locally is zero (i.e., $P(L_{i \rightarrow j}|X_{i,j,k} = 0) = 0$), the probability of local interaction can be obtained by multiplying the probability of interaction given co-occurrence with the probability of co-occurrence:

$$P(L_{i \rightarrow j}) = P(L_{i \rightarrow j}|X_{i,j,k} = 1) \times P(X_{i,k} = 1, X_{j,k} = 1). \quad (1)$$

Other important factors that can impact interaction probabilities at the local scale are taxa local abundances $N_{i,k}$ and $N_{j,k}$, which affect encounter probabilities (Canard *et al.* 2012), and local traits distributions $T_{i,k}$ and $T_{j,k}$ (e.g., body mass, longevity, and habitat breadth, Caron *et al.* 2024), which determine the ability of individuals to interact after encountering each other (Poisot *et al.* 2015). Moreover, local interactions may be conditional on local environmental factors such as temperature (Angilletta *et al.* 2004), precipitation (Woodward *et al.* 2012), habitat structure (Klecka & Boukal 2014), and presence of other taxa in the network (Kéfi *et al.* 2012; Pilosof *et al.* 2017), as described above. Here, we use the variable E_k to describe the local environmental context in which interaction probabilities were estimated. For example, in a mesocosm experiment estimating interaction probabilities between predators and prey with and without shelters, E_k would represent the presence or absence

231 of these shelters. Like co-occurrence, E_k can also be modeled probabilistically when the stochasticity or
232 uncertainty of environmental factors is considered. E_k represents all environmental variables that were taken
233 into consideration when measuring interaction probabilities; it is a subset of all environmental factors acting on
234 ecological interactions. Finally, local interaction probabilities may be conditioned on higher-level properties of
235 the network, which we denote by $f(L)$. Many topological null models (i.e., statistical models that randomize
236 interactions by retaining certain properties of the network while excluding others) provide interaction
237 probabilities from selected measures of network structure, such as connectance (Fortuna & Bascompte 2006)
238 and the degree distribution (Bascompte *et al.* 2003).

239 Local interactions must be biologically feasible before occurring at a specific time and space. A local
240 probability of interaction $P(L_{i \rightarrow j})$ can be expressed as the product of the probability of local interaction given
241 that the two taxa can potentially interact $P(L_{i \rightarrow j}|M_{i \rightarrow j} = 1)$, which we sometimes denote as $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ for the
242 sake of simplicity, with their probability of regional interaction $P(M_{i \rightarrow j})$:

$$P(L_{i \rightarrow j}) = P(L_{i \rightarrow j}|M_{i \rightarrow j} = 1) \times P(M_{i \rightarrow j} = 1). \quad (2)$$

243 Low values of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ indicate that feasible interactions rarely occur locally, intermediate values around
244 50% suggest considerable spatiotemporal variability, while high values indicate that regional interactions are
245 nearly always realized locally. The local probability of interaction between a given pair of taxa is always equal
246 to or below their probability of regional interaction.

247 The probability that two taxa i and j interact in a local network L can thus be conditioned on their co-occurrence
248 $X_{i,j,k}$ (or more explicitly on their occurrences $X_{i,k}$ and $X_{j,k}$), local abundances $N_{i,k}$ and $N_{j,k}$, local traits
249 distributions $T_{i,k}$ and $T_{j,k}$, local environmental conditions E_k , network area (or volume) A , time interval t ,
250 network properties $f(L)$ and biological (or regional) feasibility $P(M_{i \rightarrow j})$.

251 Although these variables correspond to distinct ecological inquiries or mechanisms related to ecological
252 interactions, they may covary with each other, such as the possible dependence of $X_{i,j,k}$ and E_k on spatial and
253 temporal scales. When estimating interaction probabilities using e.g. a generalized linear model with multiple
254 explanatory variables that might not be independent, it may become important to address collinearity. In such a
255 case, to mitigate this issue, it may be necessary to use variable selection techniques before fitting the model to
256 data. The probability that a local interaction is realized is described by the following expression when all these
257 conditional variables are included:

$$P(L_{i \rightarrow j} | X_{i,k}, X_{j,k}, N_{i,k}, N_{j,k}, T_{i,k}, T_{j,k}, E_k, A, t, f(L), M_{i \rightarrow j}) \quad (3)$$

258 The representation of the local context in which probabilities are estimated and the variables that should be
 259 taken into consideration depend on the study system, the objectives of the study, and the resources available to
 260 the researchers. In other words, these variables do not systematically need to be accounted for. For example,
 261 Gravel *et al.* (2019) analyzed local European trophic networks of willow-galling sawflies and their natural
 262 enemies, depicting binary interactions, all referenced in space and time, to infer probabilities of local
 263 interactions between co-occurring species. This was achieved by including temperature and precipitation as
 264 conditional variables in their models. In Box 2, we show the extent of variation among these local host-parasite
 265 networks. We do so by measuring their dissimilarity with the regional network (metaweb aggregating all local
 266 interactions), both in terms of species composition and interactions. We build local probabilistic networks
 267 following eq. 2, showing that insufficient local variation (high probability of local interaction among potentially
 268 interacting species) results in an overestimation in both the number of interactions and connectance (i.e., the
 269 proportion of all of the non-forbidden links that are realized).

270 When accounted for, conditional variables should be clearly described in the documentation of the data
 271 (Brimacombe *et al.* 2023), preferentially in mathematical terms to avoid any confusion in their interpretation
 272 and to limit manipulation errors during their re-use. For instance, ecologists should be explicit about their
 273 consideration of co-occurrence in their estimation of local interaction probabilities. Indeed, it is important to
 274 specify if probability values are conditional ($P(L_{i \rightarrow j} | X_{i,j,k} = 1)$) or not ($P(L_{i \rightarrow j})$) on co-occurrence, as this can
 275 change the interpretation of the data. In Tbl. 1, we present examples of studies that used these different
 276 formulations of probabilistic interactions and conditional variables. We have included the probability of
 277 empirically observing an interaction that is realized locally $P(O_{i \rightarrow j} | L_{i \rightarrow j})$ to underscore the distinction between
 278 local observations and actual realizations of interactions, even though the focus of this manuscript is not on the
 279 observation of interactions.

Table 1: Mathematical expressions of probabilistic interactions. The probability of interaction between two taxa i and j is interpreted differently in a local network L of realized interactions and a metaweb M of potential interactions (representing the *biological* feasibility of interactions). Each expression emphasizes a different conditional variable, the ellipsis serving as a placeholder for other variables not explicitly stated in the expression. The outcome of each of these probabilistic events, along with common uncertainty sources (often stemming from the model used for estimation), is presented alongside examples of studies that employed them (with specific variables indicated in parentheses, when applicable). The study marked with an asterisk has been conducted on binary interaction networks. Note that we also included the expression for a local network O of observed interactions and a metaweb M^* of potential interactions (representing the *ecological* feasibility of interactions) as they were mentioned in the main text, even though they are not the focus of this manuscript.

Expression	Type	Outcome	Variability sources	Reference
$P(L_{i \rightarrow j} X_{i,k}, X_{j,k}, \dots)$	local	realization of the interaction	spatiotemporal	Gravel <i>et al.</i> (2019)
		given taxa co-occurrence	variability	
$P(L_{i \rightarrow j} N_{i,k}, N_{j,k}, \dots)$	local	realization of the interaction	neutral models	Canard <i>et al.</i> (2014)
		given taxa abundances		
$P(L_{i \rightarrow j} T_{i,k}, T_{j,k}, \dots)$	local	realization of the interaction	trait matching	Caron <i>et al.</i> (2024)
		given local traits	models	
$P(L_{i \rightarrow j} E_k, \dots)$	local	realization of the interaction	environmental-	Gravel <i>et al.</i> (2019)
		given local environmental	based models	
		conditions		
$P(L_{i \rightarrow j} A, \dots)$	local	realization of the interaction in	spatial models	Galiana <i>et al.</i> (2018)
		a given area or volume		
$P(L_{i \rightarrow j} t, \dots)$	local	realization of the interaction	temporal models	Weinstein & Graham (2017a)
		during a given time period		
$P(L_{i \rightarrow j} f(L), \dots)$	local	realization of the interaction	topological null	Fortuna & Bascompte (2006) (connectance)
		given network structure	models	
$P(L_{i \rightarrow j} M_{i \rightarrow j}, \dots)$	local	realization of the interaction	spatiotemporal	this study
		given that the taxa can	variability	
		biologically interact		
$P(O_{i \rightarrow j} L_{i \rightarrow j}, \dots)$	local	observation of the interaction	sampling model	Catchen <i>et al.</i> (2023)
		given that it is realized locally		

Expression	Type	Outcome	Variability sources	Reference
$P(M_{i \rightarrow j} T_i, T_j)$	regional	biological feasibility of the interaction given regional traits (non-forbiddenness)	trait matching models	Strydom <i>et al.</i> (2022)
$P(M_{i \rightarrow j}^* T_i, T_j, E)$	regional	ecological feasibility of the interaction given regional traits and environmental conditions	trait matching and environmental-based models	this study

When using multiple models to estimate local interaction probabilities, rather than selecting a single model that best fits the data, model averaging may enhance our estimations. In this approach, the weighting is based on the likelihood of each model. Model weights represent the probability that each model is the most suitable for explaining the data, and may be measured using Akaike weights (Burnham & Anderson 2004; Wagenmakers & Farrell 2004). For instance, considering two models mod_1 and mod_2 with respective probabilities (or weights) $P(mod_1)$ and $P(mod_2)$, the average probability of interaction $P(L_{i \rightarrow j} | \dots)$ can be calculated as follows:

$$P(L_{i \rightarrow j} | \dots) = P(L_{i \rightarrow j} | mod_1, \dots) \times P(mod_1) + P(L_{i \rightarrow j} | mod_2, \dots) \times P(mod_2), \quad (4)$$

where the ellipsis serves as a placeholder for the conditional variables incorporated in these models.

Box 1: A spatiotemporally explicit model of interactions

Ecologists may resort to predictive models (e.g., generative Bayesian models) to reconstruct local networks across time and space (Strydom *et al.* 2021). Uncertainty in model parameters, model structure, and input data leads to probabilistic interactions. We introduce and develop a simple generative mechanistic model for probabilistic local interactions that takes into consideration their spatiotemporal variability (i.e., a spatiotemporally explicit model of local interactions). This model is not designed for regional interactions, which do not vary spatially and temporally. Rather, it may prove valuable for predicting local interactions across time and space by generating new interaction data following parameter inference.

As indicated by Eq. 1, the probability that two taxa i and j interact at a specific location and time k may be determined by the product of their probability of interaction given co-occurrence and their probability of

co-occurrence. First, their co-occurrence probability can be calculated using their individual (marginal) occurrence probabilities $P(X_{i,k})$ and $P(X_{j,k})$. Given that taxa occurrences are not independent of each other, the joint probability of both taxa occurring together can be calculated by multiplying the probability of one taxon being present by the conditional probability of the other occurring when the first one is present. Alternatively, the joint occurrence probability can be obtained by multiplying the marginal probabilities by the strength of association γ between the occurrences of both taxa (Cazelles *et al.* 2016):

$$P(X_{i,k}, X_{j,k}) = P(X_{i,k}|X_{j,k})P(X_{j,k}) = P(X_{i,k})P(X_{j,k})\gamma. \quad (5)$$

When $\gamma > 1$, there is a positive association in the geographic distributions of both taxa, indicating that the presence of one taxon enhances the probability of occurrence of the other. Attractions may be the result of positive interactions (e.g., mutualism) or positive/negative interactions (e.g., antagonism between predators and prey, Cazelles *et al.* 2016). In empirical networks, $\gamma > 1$ holds for the majority of species pairs (Catchen *et al.* 2023). In contrast, repulsions ($\gamma < 1$) may be caused e.g. by strong interspecific competition (Cazelles *et al.* 2016). We model the co-occurrence $X_{i,k} \cap X_{j,k}$ (hereafter $X_{i,j,k}$) of both taxa as the outcome of a Bernoulli trial

$$X_{i,j,k} \sim \text{Bernoulli}(P(X_{i,k}, X_{j,k})). \quad (6)$$

Next, the probability of interaction given co-occurrence can be made temporally explicit by modeling it as a Poisson process with rate parameter λ_k . This parameter represents the local expected frequency of interaction between the taxa within a defined time interval and can be estimated using prior data on interaction strengths, when accessible. The probability that two co-occurring taxa engage in an interaction during a time period t_0 is given by:

$$P(L_{i \rightarrow j}|X_{i,k}, X_{j,k}) = 1 - e^{-\lambda_k t_0}, \quad (7)$$

which tends toward 1 as $t_0 \rightarrow \infty$ if $\lambda > 0$. In other words, two co-occurring taxa with a nonzero rate of interaction will inevitably interact in a sufficiently long time period. It is important to note that the units of λ_k and t_0 are complementary. For instance, if the duration t_0 is measured in months, λ_k would denote the expected number of interactions per month.

The occurrence of an interaction between i and j can be modeled as a Bernoulli trial with a probability of $P(L_{i \rightarrow j})$. Consequently, a Bayesian inference model can be built based on the preceding equations to estimate the value of the λ_k and γ parameters and generate novel interaction data:

$$L_{i \rightarrow j} \sim \text{Bernoulli}(P(L_{i \rightarrow j})) \quad (8)$$

$$P(L_{i \rightarrow j}) = P(X_{i,k})P(X_{j,k})\gamma(1 - e^{-\lambda_k t_0}) \quad (9)$$

$$\gamma \sim \text{Gamma}(2, 0.5) \quad (10)$$

$$\lambda_k \sim \text{Exponential}(2) \quad (11)$$

In Fig. 1, we show the variation in the probability of interaction under different parameter values. In the right panel, we notice that, irrespective of the interaction rate λ_k , the probability of interaction converges toward an asymptote determined by the probability of co-occurrence $P(X_{i,k}, X_{j,k})$ (Eq. 5). This model can be customized in different ways, such as by linking λ_k with specific environmental variables or explicitly incorporating observation errors (i.e., the probabilities of false negatives and false positives).




Figure 1: Parameters of the spatiotemporally explicit model of interactions. (a) Probability of local interaction given by the process model (Eq. 9) under different values of λ_k (interaction rate) and γ (strength of co-occurrence), with $t_0 = 1$ (duration). Parameters t_0 and λ_k have complementary units (e.g., t_0 in months and λ_k in number of interactions per month). The parameter values used in the right panel are denoted by the white stars. (b) Scaling of the probability of interaction with the duration parameter t_0 in Eq. 9, for different values of λ_k and γ . In both panels, the marginal probabilities of occurrence $P(X_{i,k})$ and $P(X_{j,k})$ are set to a constant value of 0.5.

290

Box 2: Dissimilarity of local host-parasite networks

We use the collection of tripartite host-parasite networks of Kopalke *et al.* (2017), consisting of willows, willow-galling sawflies, and their natural enemies, sampled across Europe (233 local networks used). Given its replicated networks spanning large spatiotemporal scales, this dataset is well-suited for analyzing network variability, both in terms of species composition and interactions. Further details regarding data manipulation and network construction can be found in the supplementary material. All code and data to reproduce these analyses are available at the Open Science Framework (TBD).

In Fig. 2a-b, we show how the dissimilarity between the metaweb of binary interactions and aggregated local networks varies with the number of sampled local networks. To do so, we randomly selected one local network of binary interactions and sequentially sampled additional networks while aggregating both their species and interactions. We compared the metaweb and the aggregated local networks using the dissimilarity in species composition (β_S , Fig. 2a) and the dissimilarity of interactions between common species (β_{OS} , Fig. 2b) indices (Poisot *et al.* 2012). We repeated this sampling process one hundred times and highlighted the median dissimilarity values across simulations, as well as the 50% and 95%

291

percentile intervals. Both dissimilarity indices were calculated based on the number of items shared by the two networks (c_{LM}) and the number of items unique to the metaweb (u_M) and the aggregated local network (u_L). The β_S dissimilarity index uses species (nodes) as items being compared, while the β_{OS} index assesses dissimilarity based on interactions between shared species. Both indices were calculated following the β_W index of Whittaker (1960):

$$\beta_W = \frac{c_{LM} + u_L + u_M}{(2c_{LM} + u_L + u_M)/2} - 1. \quad (12)$$




Figure 2: Network accumulation curves. (a) Dissimilarity in species composition and (b) dissimilarity of interactions between common species between aggregated local networks and the metaweb of binary host-parasite interactions. Aggregated local networks were obtained by sequentially and randomly selecting a number of local networks of binary interactions and aggregating both their species and interactions. In both panels, the colored line represents the median dissimilarity across simulations and the grey areas cover the 50% and 95% percentile intervals. (c) Scaling of the number of interactions and (d) scaling of connectance with the number of sampled (aggregated) binary and probabilistic local interaction networks. For a better comparison with binary interactions, local networks of probabilistic interactions were derived from a metaweb of probabilistic interactions with a false positive and false negative rate of zero. A specific value of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ (the local probability of interaction among potentially interacting species, indices omitted in the figure for simplicity) was used for all local networks within a particular curve. Aggregated local networks of probabilistic interactions were obtained by sequentially and randomly selecting a number of local networks and aggregating both their species and interactions (with the value of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ adjusting according to Eq. 13).

Expectingly, local networks are highly dissimilar from the metaweb in terms of species composition, especially when only a limited number of sites has been sampled. This is because species in the metaweb form the regional species pool, and few species occur locally. Moreover, we observe a peak in the dissimilarity of interactions between common species at intermediate sampling levels. This suggests that

species are collected faster than their interactions. With a limited number of sampled local networks, few regional interactions are observed locally. Adding more sites brings new species, but not always their interactions. Interactions are more comprehensively captured when the number of sampled sites is high. Quadratic relationships of network properties with sampling effort were also observed by McLeod *et al.* (2021).

Next, we investigate how the number of local interactions and connectance scale with the number of sampled (aggregated) local networks of probabilistic or binary interactions (Fig. 2c-d). We built probabilistic local networks following Eq. 2, and set $P(M_{i \rightarrow j})$ to 1 when the interaction was observed at least once, and to 0 otherwise (i.e., no false positive or false negative in the metaweb). We used a constant value of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ for all interactions.

When aggregating local networks of probabilistic interactions, the constancy of the probability of regional interaction across the entire study area means that any rise in the probability of local interaction is solely attributable to an increase in $P(L_{i \rightarrow j}|M_{i \rightarrow j})$. For example, let L_1 and L_2 be two local networks and $L_{1,2}$ the aggregated network. If $P(L_{1,i \rightarrow j}|M_{i \rightarrow j})$ and $P(L_{2,i \rightarrow j}|M_{i \rightarrow j})$ are the probabilities that two potentially interacting taxa interact respectively in L_1 and L_2 , the probability $P(L_{1,2,i \rightarrow j}|M_{i \rightarrow j})$ that these taxa interact in the aggregated network $L_{1,2}$ is obtained by:

$$P(L_{1,2,i \rightarrow j}|M_{i \rightarrow j}) = 1 - [1 - P(L_{1,i \rightarrow j}|M_{i \rightarrow j})] \times [1 - P(L_{2,i \rightarrow j}|M_{i \rightarrow j})], \quad (13)$$

assuming independence between the interaction of the two taxa in different networks. This equation represents the probability that the interaction is realized in either (1) exclusively the local network L_1 , (2) exclusively the local network L_2 or (3) both, given that the two taxa have the biological capacity to interact.

By comparing the scaling relationships observed in local networks of binary (empirical) and probabilistic (modeled) interactions, Fig. 2c-d shows that high values of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ lead to systematic overestimations in the number of interactions and connectance, especially when $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 1$ (corresponding to the scenario where local probabilities of interactions are equivalent to the probabilities of regional interactions). This suggests that high values of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ do not adequately capture the variability of local interactions. However, these biases tend to diminish as the number of sampled networks increases, indicating that most interactions are eventually captured when $P(L_{i \rightarrow j}|M_{i \rightarrow j})$ is high. In contrast,

low values of $P(L_{i \rightarrow j} | M_{i \rightarrow j})$ lead to missing interactions, resulting in an underestimation of the number of interactions and connectance. These results underscore the importance of using the appropriate level of variability when estimating local interaction probabilities.

295

296 Metawebs: regional catalogs of interactions

297 Metawebs (Dunne 2006) are networks of potential interactions over broad spatial, temporal, and taxonomic
298 scales (e.g., food webs at the continental scale). Potential interactions describe the biological capacity of taxa to
299 interact, which is typically assessed at the regional scale. Metawebs of probabilistic interactions are particularly
300 useful in situations where there is uncertainty in the ability of taxa to interact. This uncertainty frequently arises
301 due to insufficient interaction data, especially for taxa that have not yet been observed to co-occur, and
302 uncertainties in trait-matching models. As data accumulates, interactions in metawebs should tend towards
303 binarity, approaching probability values of 0 (repeatedly failing to observe an interaction between two
304 co-occurring taxa) and 1 (observing an interaction at least once). The extent of sampling effort thus influences
305 our evaluation of probabilities of potential interactions, as sampling over a larger area or for a longer duration
306 enables us to capture a greater number of regional interactions (McLeod *et al.* 2021). However, in contrast with
307 local networks of probabilistic interactions, which describe local interaction stochasticity, regional interactions
308 are not evaluated for any particular local context. In Box 3, we discuss the differences in spatial and temporal
309 scaling of regional interactions compared to local interactions. We do so by using the host-parasite networks of
310 Kopelke *et al.* (2017) as an illustration of spatial scaling. Although *neutrally* forbidden interactions (i.e.,
311 between rare species, Canard *et al.* 2012) tend to have low probability values in local networks, they may have
312 higher probabilities in the metaweb (i.e., when species' biological traits can support an interaction if they were
313 to encounter each other). Likewise, non-co-occurring taxa may have a non-zero probability of interaction in the
314 metaweb.

315 Potential interactions describe what we refer to as the *biological* feasibility of interactions, which is based solely
316 on the regional traits distributions T_i and T_j of taxa i and j , respectively. A probability of potential interaction in
317 a metaweb M describing the biological feasibility of interactions may be expressed as:

$$P(M_{i \rightarrow j} | T_i, T_j), \quad (14)$$

which, in contrast with local networks, is not conditioned on any spatial, temporal, co-occurrence or environmental variables (Tbl. 1). Regional traits may differ from local traits $T_{i,k}$ and $T_{j,k}$, which may vary spatially and temporally due to phenotypic plasticity (Berg & Ellers 2010). The taxonomic level at which interactions are evaluated influences the distribution of regional and local traits. However, as explained in Box 4, there is no fundamental difference in the taxonomic scaling of regional and local interactions (i.e., how interaction probabilities change with taxonomic level), unlike their spatial and temporal scaling.

The biological feasibility of interactions expresses our degree of belief that there exists at least one combination of phenotypes that could support an interaction if they were to encounter each other, assuming they had infinite time to interact. Evaluating this probability is conducted without incorporating the environmental conditions under which they encounter each other into the model, as the complement of the probability $P(F_{i \rightarrow j}|T_i, T_j)$ of forbidden interactions based uniquely on biological traits (i.e., the probability that their traits do not support an interaction):

$$P(M_{i \rightarrow j}|T_i, T_j) = 1 - P(F_{i \rightarrow j}|T_i, T_j). \quad (15)$$

For example, let i be a western diamondback rattlesnake (*Crotalus atrox*) and j , a wood lemming (*Myopus schisticolor*). These two taxa never co-occur, the rattlesnake being adapted to warm regions of North America (Castoe *et al.* 2007) and the lemming, to northern habitats of Eurasia (Fedorov *et al.* 2008). As we lack direct observations of an interaction between these two species, we have to rely on expert knowledge or trait-matching models to estimate their probability of potential interaction. To accurately estimate this probability using trait-matching models, it is crucial to ensure that the set of traits considered reflects the overall traits distributions of both taxa. We could for instance consider their average body mass and the average phylogenetic proximity of lemmings to rattlesnakes' prey. Doing so, we might find a high probability of potential interaction based on these traits.

The biological feasibility of interactions should not be confused with what we refer to as the *ecological* feasibility of interactions. A probability of potential interaction in a metaweb M^* describing ecological feasibility of interactions may be expressed as:

$$P(M_{i \rightarrow j}^*|T_i, T_j, E), \quad (16)$$

342 where E is the environmental conditions under which potential interactions are evaluated (Tbl. 1). Unlike E_k ,
 343 these environmental conditions do not represent conditions occurring at specific locations. Ecological
 344 feasibility represents the probability that two taxa interact if they were to encounter each other under given
 345 environmental conditions, assuming they had infinite time to interact. Incorporating environmental conditions
 346 into a trait-matching model may be important when there is high covariance between the environment and traits.
 347 For instance, in our example involving rattlesnakes and lemmings, the probability of potential interaction
 348 between these two species may be low in most environmental conditions. Western diamondback rattlesnakes
 349 may be unactive under low temperatures (Kissner *et al.* 1997), whereas wood lemmings may have low tolerance
 350 to high temperatures (Kausrud *et al.* 2008). The probability that an interaction is ecologically feasible is always
 351 lower than the probability that it is biologically feasible, even across all environmental conditions:

$$\int_E P(M_{i \rightarrow j}^* | T_i, T_j, E) dE \leq P(M_{i \rightarrow j} | T_i, T_j). \quad (17)$$

352 This is because the biological feasibility of an interaction is a prerequisite for its ecological feasibility.
 353 Biological feasibility is necessary but not sufficient for an interaction to be ecologically feasible. Our discussion
 354 of metawebs will focus on the biological feasibility of interactions since most methods developed for inferring
 355 probabilities of regional interactions do not explicitly take into account environmental conditions (e.g., Strydom
 356 *et al.* 2022).

357 Starting from a selected set of taxa, which are usually distributed within a broad region of interest, metawebs
 358 can be built using different data sources, including literature review (e.g., Maiorano *et al.* 2020), aggregated
 359 interaction data (e.g., Gravel *et al.* 2019; Saravia *et al.* 2022), trait-matching models (e.g., Shaw *et al.* 2024;
 360 Strydom *et al.* 2022), and expert knowledge. Every pair of taxa that have confidently been observed to interact
 361 at least once can be given a probability of 1 (i.e., $P(M_{i \rightarrow j}) = 1$) since we know that they *can* interact. This
 362 differs from local networks of probabilistic interactions, where interaction events may remain stochastic (i.e.,
 363 $P(L_{i \rightarrow j}) < 1$) even after empirically observing interactions due to their spatiotemporal variability. Interactions
 364 that were never observed typically have low probability values in local networks and vary from low to high
 365 values in metawebs, contingent upon taxa traits distributions (reaching 0 for forbidden links).

366 When incorporating local network data to estimate probabilities of regional interactions, repeatedly failing to
 367 observe an interaction between two co-occurring taxa (i.e., $P(O_{i \rightarrow j} = 0)$) should decrease the probability that
 368 the interaction is biologically feasible. Using Bayes' theorem, the probability that the interaction is biologically

369 feasible given that it was never observed locally, $P(M_{i \rightarrow j} = 1 | O_{i \rightarrow j} = 0, \dots)$, may be calculated as follows:

$$P(M_{i \rightarrow j} = 1 | O_{i \rightarrow j} = 0, \dots) = \frac{P(O_{i \rightarrow j} = 0 | M_{i \rightarrow j} = 1, \dots) \times P(M_{i \rightarrow j} = 1 | \dots)}{P(O_{i \rightarrow j} = 0 | \dots)}. \quad (18)$$

370 The reduction in the probability of regional interaction after considering that it was never observed locally (i.e.,
371 $P(M_{i \rightarrow j} = 1 | O_{i \rightarrow j} = 0, \dots) < P(M_{i \rightarrow j} = 1 | \dots)$) occurs because $P(O_{i \rightarrow j} = 0 | M_{i \rightarrow j} = 1, \dots)$ must be lower than
372 $P(O_{i \rightarrow j} = 0, \dots)$, i.e. there is a higher chance of observing an interaction when we know it is biologically
373 feasible.

374 Many observations of interactions are false positives because of observation errors due to taxonomic
375 misidentifications and ecological misinterpretations (e.g., due to cryptic species and interactions, Pringle &
376 Hutchinson 2020). Likewise, forbidden interactions may be false negatives, e.g. if they have been evaluated
377 based on unrepresentative or incomplete traits distributions. Employing Bayesian models proves valuable when
378 estimating interaction probabilities in metawebs (e.g., Bartomeus *et al.* 2016; Cirtwill *et al.* 2019). This
379 improvement is achieved by updating prior information regarding the feasibility of interactions (e.g., experts'
380 prior assessments of interaction probabilities) with empirical data on interactions and traits.

Box 3: Spatial and temporal scaling of interactions

Network-Area Relationships document the scaling of network properties (such as modularity and connectance) with spatial scale (e.g., Galiana *et al.* 2018; Wood *et al.* 2015). The variation in network structure across spatial scales may stem from the scaling of species richness (species-area relationships, SARs), the number of interactions (Brose *et al.* 2004), and many other higher-level properties of the system (e.g., environmental heterogeneity, Thompson & Townsend 2005) with the sampled area. Likewise, interaction accumulation curves describe the scaling of the number of observed interactions with sampling effort (Jordano 2016). Sampling effort, which may correspond to the duration of the sampling period used to construct the network, can impact connectance (Bersier *et al.* 1999) and various measures of network structure (Banašek-Richter *et al.* 2004; McLeod *et al.* 2021). Apart from sampling effort, the temporal scaling of interactions also describes how network structure changes with the temporal resolution of the network, acknowledging that distinct interactions take place over time, ranging from short-term fluctuations of interactions to long-term trends. As local networks of probabilistic interactions may explicitly account for the spatiotemporal variability of interactions, they offer a distinct approach to

investigating the scaling of network structure with space and time.

Local networks and metawebs exhibit distinct relationships with spatial and temporal scales. On one hand, metawebs of probabilistic interactions feature regional interactions that do not scale with space and time. This is because regional interactions depend solely on the biological capacity of two taxa to interact, regardless of their co-occurrence and local environmental conditions. However, probabilities of potential interactions may change, tending to become more definitive with increased sampling effort (even though they do not vary with the spatial and temporal extent (boundaries) of the network). The probability of two taxa potentially interacting should be the same in all metawebs in which they are present, provided that the data and methods used for estimation are consistent. For example, if a smaller metaweb M_1 is derived from a larger metaweb M_0 by selecting the subset of taxa present in the region described by M_1 and retaining all their interactions, their probabilities of interaction should be identical regardless of scale, i.e. $P(M_{1,i \rightarrow j}) = P(M_{0,i \rightarrow j})$. With a larger or equal number of taxa in M_0 as compared to M_1 , the total number of interactions is expected to be higher in the larger network, even though pairwise probabilities remain identical.

On the other hand, local interactions scale both spatially and temporally, given that they have more opportunities to be realized and observed in larger areas and longer durations. This is attributed to factors such as a higher number of individuals, greater trait variations, and increased opportunities for encounters, as highlighted by McLeod *et al.* (2020). For example, if a local network of probabilistic interactions L_1 with an area A_1 is derived from a larger network L_0 with an area A_0 , and A_1 is entirely nested within A_0 , interaction probabilities should be lower in the smaller network, i.e. $P(L_{1,i \rightarrow j}|A_1 < A_0) \leq P(L_{0,i \rightarrow j}|A_0)$. However, if A_1 and A_0 are disjoint, interaction probabilities could be higher in the smaller area, contingent upon their environmental and biological conditions. Likewise, interaction probabilities should be lower in networks with shorter durations when time intervals are nested. In Fig. 3, we show how the expected number of local host-parasite interactions scales with area, represented as an expanding latitudinal window, in comparison with regional interactions (data from Kopelke *et al.* 2017). Even though we employed local probabilities of interactions equal to regional interactions for the purpose of comparison (i.e., using $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 1$ here), we notice that the total number of regional interactions scales more rapidly than local interactions. This is because numerous regional interactions involve species that never co-occur, and as a result, these interactions are not captured in local networks.




Figure 3: Spatial scaling of interactions. Expected number of host-parasite interactions in a network aggregating all (a) local and (b) regional probabilistic interactions within a latitudinal window of a given width. Every dashed curve corresponds to a different window centered at a given latitude (color bar), with the pink solid line representing the median number of interactions across windows. Heatmaps of the expected number of (c) local and (d) regional interactions found in windows of specified width and central latitudes. Probabilities of regional interactions were obtained with a false positive rate of 5% and a false negative rate of 10%. Local probabilistic interactions were derived from probabilistic regional interactions by setting the value of $P(L_{i \rightarrow j} | M_{i \rightarrow j})$ (the local probability of interaction among potentially interacting species) to 1, ensuring a conservative comparison between aggregated local networks and metawebs. Aggregated local networks were obtained by aggregating both the species and interactions found within a particular latitudinal window, with the values of $P(L_{i \rightarrow j} | M_{i \rightarrow j})$ remaining at their maximum value of 1 following Eq. 13. Further details on the data and network construction are presented in the supplementary material.

Box 4: Taxonomic scaling of interactions

Probabilistic interaction networks offer a versatile approach to tackle a broad array of ecological questions, depending on their level of organization. For instance, the assemblage of interactions across ecological scales can be explored through species-based networks, while clade-based networks provide insights into macroevolutionary processes (Gomez *et al.* 2010). Given that our interpretation of the properties and dynamics of ecological networks depends on their taxonomic level (Guimarães 2020), investigating the taxonomic scaling of interactions (i.e., how interaction probabilities change with taxonomic level) emerges as a promising research avenue.

There are no inherent differences between the taxonomic scaling of local networks and metawebs. Interaction probabilities in both local networks (Eq. 3) and metawebs (Eq. 14) are not directly conditioned on taxonomic level. The taxonomic scale is tied to the definition of the event itself (i.e., the interaction between two taxa, defined at the desired taxonomical scale), not to the variables on which interaction probabilities are conditioned. However, some conditional variables (e.g. trait distribution) may vary with taxonomic scale. In such cases, interaction probabilities would vary taxonomically following the scaling of these variables.

In both types of networks, transitioning to a broader level of organization (e.g., from a species-level network S to a genus-level network G) can be accomplished directly by using probabilities from finer scales. For example, in a network with n_1 species from genus g_1 and n_2 species from genus g_2 , one can compute the probability that at least one species from genus g_1 interacts with at least one species from genus g_2 (i.e., that there is a non-zero number of species-level interactions or, equivalently, that the genus-level interaction occurs) as follows:

$$P(G_{g_1 \rightarrow g_2} = 1) = 1 - \prod_{i=1}^{n_1} \prod_{j=1}^{n_2} (1 - P(S_{g_1,i \rightarrow g_2,j} = 1)), \quad (19)$$

where $g_{1,i}$ and $g_{2,j}$ are the species of the corresponding genus and assuming independence between species-level interactions. If it is known that at least two of these species interact (i.e., $P(S_{g_1,i \rightarrow g_2,j}) = 1$ for at least one pair of $(g_{1,i}, g_{2,j})$), it implies a probability of genus interaction equal to 1. Canard *et al.* (2012) built a species-based network following a similar approach, by using simulated interactions between individuals derived from a neutral model (i.e., a model that assumed ecological equivalence among individuals).

In contrast, a more sophisticated approach is necessary when transitioning from a broader to a finer level of organization. This is because the knowledge of an interaction between two genera does not guarantee that all possible pairwise combinations of their species will also interact. One possible method is to build a finer-scale network by generating probabilities of interactions through random sampling from a beta distribution, parameterized by the broader-scale network.

Ideally, our biological interpretation of probabilistic interactions should remain consistent across a network even if it incorporates heterogeneous levels of organization, such as a network whose nodes represent both species and trophic species (groups of species sharing similar predators and prey). This is common in ecological networks where taxonomic resolution is typically low (Hemprich-Bennett *et al.* 2021; Vázquez *et al.* 2022). Interaction probabilities at broader taxonomic scales should be based on probabilities of interactions between individuals, for both local networks and metawebs. In local individual-based food webs, the probability that two individuals interact reflects our degree of belief that one individual will consume the other. In local species-based food webs, the probability that two species interact represents our degree of belief that at least one individual from the predator species consumes at least another individual from the prey species. In local clade-based food webs, the probability that two clades interact represents our degree of belief that at least two species from these clades interact with each other or, equivalently, that at least two individuals from these clades interact with each other. Fundamentally, the taxonomic scaling of interactions involves aggregating interactions between individuals into larger groups, which may exhibit varying degrees of homogeneity based on the organisms and the study system. In that regard, taxonomic scaling is analogous to the spatial and temporal scaling of interactions, as they all represent different ways to aggregate individuals into broader groups (either spatially, temporally, or taxonomically).

385

Box 5: Sampling for binary interaction networks

The prediction of binary interactions through Bernoulli trials is an important application of probabilistic interaction networks. This approach proves beneficial for analyzing the structural characteristics of probabilistic interaction networks, particularly in the absence of specific analytical formulas for measures of network structure (Poisot *et al.* 2016). By performing independent Bernoulli trials for each interaction, a network of binary interactions may be generated. A probability distribution of network properties can be obtained by measuring network structure across multiple randomly generated networks (Poisot *et al.*

386

2016). This method enables the representation of the variability or uncertainty of network structure, albeit with possible biases when connectance is low (Chagnon 2015; Poisot & Gravel 2014). Employing this strategy to generate binary interaction networks under a null model facilitates null hypothesis significance testing, wherein the observed measure is compared against the simulated distribution (e.g., Bascompte *et al.* 2003). Yet, inferring network structure through sampling for binary interactions assumes independence among interactions, which might not accurately represent reality. Covariation among interactions could exist even if we do not explicitly condition interactions on others. For example, an interaction between two taxa could be more probable when another interaction occurs. The consequences of this assumption of independence on the prediction of network structure have yet to be empirically examined.

There are at least two approaches to sampling binary interaction networks across space from probabilistic interaction networks (i.e., when predicting a binary interaction network for each location k within a given region). Both approaches assume independence between interactions. The first approach involves performing a singular Bernoulli trial for each pair of taxa based on their regional probability of interaction:

$$M_{i \rightarrow j} \sim \text{Bernoulli}(P(M_{i \rightarrow j})).$$

In employing this approach, we predict a single metaweb of binary interactions for each simulation. Every pair of taxa predicted to interact in this metaweb will be treated as interacting in all localized networks where they co-occur, i.e. $L_{k,i \rightarrow j} = M_{i \rightarrow j}$ when $X_{i,j,k} = 1$. This will result in local pairwise interactions without spatial variation.

The second approach is to independently sample each local network of probabilistic interactions:

$$L_{k,i \rightarrow j} \sim \text{Bernoulli}(P(L_{k,i \rightarrow j})).$$

This can be achieved by first generating distinct probabilistic interaction networks for each location. These local networks of probabilistic interactions may vary in taxa composition and interaction probabilities. Because binary interaction networks are sampled independently for each location, this second approach introduces spatial variation in binary interactions. This approach may more effectively capture network structure across space and time, facilitating the investigation of ecological hypotheses about interactions at broad spatial and temporal scales.

In Fig. 4, we compare the average connectance of binary interaction networks resulting from these

two sampling techniques. We drew regional and local interactions from our host-parasite networks of probabilistic interactions (Kopelke *et al.* 2017), generating a number of binary interaction network realizations for each site in the dataset. These two sampling approaches yield different outcomes, particularly for intermediate values of $P(L_{i \rightarrow j}|M_{i \rightarrow j})$, which represent instances where regional interactions do not consistently manifest locally (i.e., with the largest local variability). Small differences between these techniques are also apparent when we equate the probability of local interaction to the probability of regional interaction (i.e., when using $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 1.0$, Fig. 4a,d), especially when the number of samples of binary interaction networks for each location is low. As anticipated, we observe that sampling binary interactions from the metaweb tends to overestimate connectance on average compared to sampling them from local networks (Fig. 4a-c). Furthermore, we observe an increase in the variability of connectance when employing a single simulation (Fig. 4, cross markers), which is a more tangible representation of the process leading to the realization of local interactions in nature.




Figure 4: Connectance of sampled binary interaction networks. (a-c) Average connectance of binary interaction networks obtained from the two sampling approaches for 20 randomly selected host-parasite networks. Cross markers represent the connectance of a single sample for each network, diamond markers the average connectance across 10 samples, hexagon markers the average connectance across 50 samples, and the colored circles the average connectance across 100 samples (marker size proportional to the number of samples). (d-f) Reduction in the mean squared logarithmic error between the average connectance of binary interaction networks (all 233 host-parasite networks) obtained from these two sampling methods as the number of samples increases. The local probability of interaction between potentially interacting species was set to three different values: (a,d) $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 1.0$, (b,e) $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 0.75$, and (c,f) $P(L_{i \rightarrow j}|M_{i \rightarrow j}) = 0.50$. Probabilities of regional interactions were obtained with a false positive rate of 5% and a false negative rate of 10%. Further details on the data and network construction are presented in the supplementary material. Regional samples were obtained by randomly sampling binary interactions from the probabilistic interaction metaweb, and then propagating this result to all local networks that include the species potentially engaged in the interactions. Local binary interaction networks were generated by independently sampling binary interactions for each local network of probabilistic interactions.

The choice of sampling approach can influence the selection of grid cell size when delineating local communities within a broader region of interest. In the first approach, pairwise interactions remain constant irrespective of cell size because they are sampled only once from the metaweb. However, in the second approach, local interaction probabilities are contingent on network area. For instance, consider the local networks L_1 and L_2 with an area of $\frac{1}{2}A_0$, both nested within A_0 but disjoint from each other, forming L_0 . If we treat L_1 and L_2 as independent, the probability of interaction between taxa i and j in L_0

is given by:

$$P(L_{0,i \rightarrow j}) = 1 - [1 - P(L_{1,i \rightarrow j})] \times [1 - P(L_{2,i \rightarrow j})]. \quad (20)$$

Due to its larger area, the probability that the two taxa interact in L_0 is equal or greater than in L_1 and L_2 . When sampling binary interactions from local networks, it is crucial to sample at the same spatial scale for which probabilities were estimated. Otherwise, interaction probabilities must be adjusted to align with the intended cell size, preventing systematic biases in predictions.

390

391 Conclusion

392 In this contribution, we underline the importance of network documentation for adequately interpreting and
393 manipulating probabilistic interaction data. The mathematical representation of probabilities and their statistical
394 properties depend on the type of interactions (local or regional) and the conditions under which these
395 interactions were evaluated. We showed that local networks and metawebs of probabilistic interactions differ in
396 their relationship to spatial and temporal scales (Box 3), with regional interactions remaining consistent across
397 scales. In contrast with metawebs, local interactions are measured in a specific context (e.g., in a given area,
398 time, and biological and environmental conditions) and depend on taxa co-occurrence. These differences bring
399 to light the need to use probabilistic data with caution, for instance when generating network realizations of
400 binary interactions across space (Box 5). Clear documentation describing the type of interaction and the
401 variables used in their estimation are required to ensure adequate data manipulation. Sound data practices and
402 foundations for probabilistic thinking in network ecology facilitate reliable assessments of the spatiotemporal
403 variability and uncertainty of biotic interactions. Here we identify key research priorities for improving our
404 understanding of local and regional interactions and their stochasticity.

405 Predicting local networks from metawebs

406 Metawebs are a valuable source of ecological information for predicting local networks across time and space.
407 Local networks of binary interactions can be reconstructed by selecting a subset of taxa and interactions from
408 the metaweb (Dunne 2006). Metawebs thus contain more interactions than local networks, even though their
409 connectance is usually much smaller than the one of local networks (Gravel *et al.* 2011). Determining the list of

410 taxa to select can be achieved empirically (e.g., observed occurrence data for a site) or numerically (e.g., species
411 distribution models). As species composition is arguably easier to sample or predict than pairwise interactions,
412 the primary challenge lies in deciding which interactions to select from the metaweb. Inferring the structure of
413 local networks from the metaweb before predicting local pairwise interactions could hold promise (Strydom *et
414 al.* 2021), considering that the structure of local networks is constrained by the metaweb (Saravia *et al.* 2022).

415 Inferring local networks of probabilistic interactions from a metaweb comes with its own set of challenges. For
416 example, Dansereau *et al.* (2023) inferred spatially-explicit food webs from a metaweb of probabilistic trophic
417 interactions between Canadian mammals. Their predicted localized food webs are downscaled versions of the
418 metaweb (i.e., localized metawebs with the same interaction probabilities as those in the regional metaweb). To
419 infer local networks as defined in this manuscript (i.e., describing local realizations of interactions), local
420 interaction probabilities must be smaller than regional interaction probabilities. This decrease is due to the
421 prerequisite that two taxa must initially possess the capacity to interact before engaging in local interactions
422 (Eq. 2). Inferring local networks from a metaweb by maintaining identical interaction probability values
423 introduces systematic biases into the predictions, as discussed in Box 2 (unless networks are seen as downscaled
424 metawebs).

425 As suggested by McLeod *et al.* (2021), although metawebs do not capture the spatiotemporal variability of
426 interactions, they establish an upper limit for local interactions (similarly for metawebs of probabilistic
427 interactions, Strydom *et al.* 2023). In other words, the probability that two taxa interact at a specific location
428 and time is consistently lower or equal to the probability of their regional interaction, regardless of the
429 conditional variables considered:

$$P(L_{i \rightarrow j} | \dots) \leq P(M_{i \rightarrow j} | T_i, T_j). \quad (21)$$

430 Moreover, the probability that two taxa possess the biological capacity to interact must be higher than the
431 probability of them interacting at any location and time because they may never co-occur or encounter locally.
432 Specifically, the cumulative probability of local interaction across all spatial, temporal, and environmental
433 conditions must be less than the probability of regional interaction, i.e.

$$\int_{E_k} \int_A \int_t P(L_{i \rightarrow j} | E_k, A, t) dt dA dE_k \leq P(M_{i \rightarrow j} | T_i, T_j). \quad (22)$$

434 Estimating more precisely the probability $P(L_{i \rightarrow j} | M_{i \rightarrow j})$ that two taxa interact locally if they can potentially
435 interact allows for improved predictions of local networks from the metaweb of probabilistic interactions. This
436 task is challenging due to the variability of this probability across space and time, as well as its variability
437 across pairwise interactions within a network. Using simple models of $P(L_{i \rightarrow j} | M_{i \rightarrow j})$, as shown in Box 2,
438 represents an initial step toward the overarching objective of reconstructing local networks from metawebs.

439 **Quantifying and reducing interaction uncertainty**

440 While sampling biological communities decreases the uncertainty of interactions by accumulating evidence for
441 their feasibility and local realization, there is a limit to how much we can reduce uncertainty. In metawebs,
442 probabilities reflect our limited knowledge of interactions (i.e., our degree of belief that interactions are
443 feasible), which is expected to improve with a larger volume of data. Regional interactions should become more
444 definitive (with probabilities approaching 0 or 1) as we investigate various conditions, including different
445 combinations of species traits.

446 In comparison, although local networks can be seen as random instances of metawebs, their stochasticity cannot
447 be reduced to the same extent. Local interaction probabilities may represent both their uncertainty and
448 spatiotemporal variability. Owing to environmental heterogeneity, there will invariably be instances in which an
449 interaction occurs and others in which it does not, across different times and locations, irrespective of the extent
450 to which we can improve our knowledge of its biological feasibility and the local conditions that facilitate its
451 occurrence. When local networks describe probabilities of observing interactions rather than their actual
452 occurrence, we must also consider observation variability (sampling error) as an additional source of
453 stochasticity. Every ecological process is stochastic but there is also a possibility that a phenomenon goes
454 undetected. Quantifying and partitioning this stochasticity will enable us to make more accurate predictions
455 about ecological interactions at various spatial and temporal scales. This will prove to be of vital importance as
456 our time to understand nature runs out, especially at locations where the impacts of climate change and habitat
457 loss hit harder.

458 **Relaxing the independence assumption**

459 Estimating local interaction probabilities independently for each taxa pair and assembling them into a network
460 of probabilistic interactions comes with limitations. Predicting local networks of binary interactions based on

461 these interaction probabilities assumes independence among interactions, a condition seldom respected in
462 practice (Golubski & Abrams 2011). Relaxing this assumption is the next logical step in the stochastic
463 representation of interactions.

464 A more accurate representation of the stochasticity of ecological networks involves creating *probabilistic*
465 *networks* ($P(L|...)$ and $P(M|...)$), rather than networks of *probabilistic interactions* ($P(L_{i \rightarrow j}|...)$ and $P(M_{i \rightarrow j}|...)$).
466 Probabilistic networks describe the probability that a particular network of binary (or quantitative) interactions
467 (its whole adjacency matrix) is realized. For example, Young *et al.* (2021) used a Bayesian approach to estimate
468 the probability of different plant-pollinator network structures derived from imperfect observational data. A
469 probability distribution of ecological networks may also be derived using the principle of maximum entropy
470 given structural constrained (e.g., Cimini *et al.* 2019; Park & Newman 2004).

471 Regardless of the method employed, generating probabilistic local networks could lead to more accurate
472 predictions of local networks of binary interactions by bypassing the independence assumption. Probabilistic
473 networks could serve as an alternative to null hypothesis significance testing when comparing the structure of a
474 local network to some random expectations or, as done in Pellissier *et al.* (2018), to the metaweb. These random
475 expectations are typically derived by performing a series of Bernoulli trials on probabilistic interactions,
476 assuming independence, to generate a distribution of networks of binary interactions to calculate their structure
477 (Poisot *et al.* 2016). One could for instance compare the likelihood of an observed network to the one of the
478 most likely network structure (according to the probabilistic network distribution), thereby directly obtaining a
479 measure of discrepancy of the empirical network. Generating probabilistic ecological networks represents a
480 tangible challenge, one that, in the coming years, promises to unlock doors to more advanced and adequate
481 analyses of ecological networks.

482 Acknowledgment

483 We acknowledge that this study was conducted on land within the traditional unceded territory of the Saint
484 Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and Omàmiwininiwak nations. This work was
485 supported by the Institute for Data Valorisation (IVADO) and the Natural Sciences and Engineering Research
486 Council of Canada (NSERC) Collaborative Research and Training Experience (CREATE) program, through the
487 Computational Biodiversity Science and Services (BIOS²) program. A special thanks to all members of the
488 Black Holes and Revelations working group (organized by BIOS²) for their insightful discussions and valuable

489 feedback on this manuscript.

490 **References**

- 491 Abrego, N., Roslin, T., Huotari, T., Ji, Y., Schmidt, N.M., Wang, J., *et al.* (2021). Accounting for species
492 interactions is necessary for predicting how arctic arthropod communities respond to climate change.
493 *Ecography*, 44, 885–896.
- 494 Angilletta, M.J., Jr., Steury, T.D. & Sears, M.W. (2004). Temperature, Growth Rate, and Body Size in
495 Ectotherms: Fitting Pieces of a Life-History Puzzle¹. *Integrative and Comparative Biology*, 44, 498–509.
- 496 Araujo, M.B., Rozenfeld, A., Rahbek, C. & Marquet, P.A. (2011). Using species co-occurrence networks to
497 assess the impacts of climate change. *Ecography*, 34, 897–908.
- 498 Banašek-Richter, C., Cattin, M.-F. & Bersier, L.-F. (2004). Sampling effects and the robustness of quantitative
499 and qualitative food-web descriptors. *Journal of Theoretical Biology*, 226, 23–32.
- 500 Bartomeus, I., Gravel, D., Tylianakis, J.M., Aizen, M.A., Dickie, I.A. & Bernard-Verdier, M. (2016). A
501 common framework for identifying linkage rules across different types of interactions. *Functional Ecology*,
502 30, 1894–1903.
- 503 Bascompte, J., Jordano, P., Melián, C.J. & Olesen, J.M. (2003). The nested assembly of plant–animal
504 mutualistic networks. *Proceedings of the National Academy of Sciences*, 100, 9383–9387.
- 505 Berg, M.P. & Ellers, J. (2010). Trait plasticity in species interactions: A driving force of community dynamics.
506 *Evolutionary Ecology*, 24, 617–629.
- 507 Berlow, E.L., Neutel, A.-M., Cohen, J.E., De Ruiter, P.C., Ebenman, B., Emmerson, M., *et al.* (2004).
508 Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73, 585–598.
- 509 Bersier, L.-F., Dixon, P. & Sugihara, G. (1999). Scale-Invariant or Scale-Dependent Behavior of the Link
510 Density Property in Food Webs: A Matter of Sampling Effort? *The American Naturalist*, 153, 676–682.
- 511 Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011). Why
512 intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26, 183–192.
- 513 Borrett, S.R. & Scharler, U.M. (2019). Walk partitions of flow in Ecological Network Analysis: Review and
514 synthesis of methods and indicators. *Ecological Indicators*, 106, 105451.

- 515 Boulangeat, I., Gravel, D. & Thuiller, W. (2012). Accounting for dispersal and biotic interactions to disentangle
516 the drivers of species distributions and their abundances. *Ecology Letters*, 15, 584–593.
- 517 Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T. & Fortin, M.-J. (2023). Shortcomings of reusing
518 species interaction networks created by different sets of researchers. *PLOS Biology*, 21, e3002068.
- 519 Broom, M. & Ruxton, G.D. (2005). You can run—or you can hide: Optimal strategies for cryptic prey against
520 pursuit predators. *Behavioral Ecology*, 16, 534–540.
- 521 Brose, U., Ostling, A., Harrison, K. & Martinez, N.D. (2004). Unified spatial scaling of species and their
522 trophic interactions. *Nature*, 428, 167–171.
- 523 Burnham, K.P. & Anderson, D.R. (2004). Multimodel Inference: Understanding AIC and BIC in Model
524 Selection. *Sociological Methods & Research*, 33, 261–304.
- 525 Canard, E.F., Mouquet, N., Mouillot, D., Stanko, M., Miklisova, D. & Gravel, D. (2014). Empirical Evaluation
526 of Neutral Interactions in Host-Parasite Networks. *The American Naturalist*, 183, 468–479.
- 527 Canard, E., Mouquet, N., Marescot, L., Gaston, K.J., Gravel, D. & Mouillot, D. (2012). Emergence of
528 Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7, e38295.
- 529 Carlson, C.J., Farrell, M.J., Grange, Z., Han, B.A., Mollentze, N., Phelan, A.L., et al. (2021). The future of
530 zoonotic risk prediction. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376,
531 20200358.
- 532 Caron, D., Brose, U., Lurgi, M., Blanchet, F.G., Gravel, D. & Pollock, L.J. (2024). Trait-matching models
533 predict pairwise interactions across regions, not food web properties. *Global Ecology and Biogeography*,
534 33, e13807.
- 535 Castoe, T.A., Spencer, C.L. & Parkinson, C.L. (2007). Phylogeographic structure and historical demography of
536 the western diamondback rattlesnake (*Crotalus Atrox*): A perspective on North American desert
537 biogeography. *Molecular Phylogenetics and Evolution*, 42, 193–212.
- 538 Catchen, M.D., Poisot, T., Pollock, L.J. & Gonzalez, A. (2023). The missing link: Discerning true from false
539 negatives when sampling species interaction networks.
- 540 Cazelles, K., Araujo, M.B., Mouquet, N. & Gravel, D. (2016). A theory for species co-occurrence in interaction
541 networks. *Theoretical Ecology*, 9, 39–48.

- 542 Chagnon, P.-L. (2015). Characterizing topology of ecological networks along gradients: The limits of metrics'
543 standardization. *Ecological Complexity*, 22, 36–39.
- 544 Choh, Y., Ignacio, M., Sabelis, M.W. & Janssen, A. (2012). Predator-prey role reversals, juvenile experience
545 and adult antipredator behaviour. *Scientific Reports*, 2, 728.
- 546 Cimini, G., Squartini, T., Saracco, F., Garlaschelli, D., Gabrielli, A. & Caldarelli, G. (2019). The statistical
547 physics of real-world networks. *Nature Reviews Physics*, 1, 58–71.
- 548 Cirtwill, A.R., Eklöf, A., Roslin, T., Wootton, K. & Gravel, D. (2019). A quantitative framework for
549 investigating the reliability of empirical network construction. *Methods in Ecology and Evolution*, 10,
550 902–911.
- 551 Dallas, T., Park, A.W. & Drake, J.M. (2017). Predicting cryptic links in host-parasite networks. *PLOS
552 Computational Biology*, 13, e1005557.
- 553 Dansereau, G., Barros, C. & Poisot, T. (2023). Spatially explicit predictions of food web structure from regional
554 level data.
- 555 Dunne, J.A. (2006). The Network Structure of Food Webs. In: *Ecological networks: Linking structure and
556 dynamics* (eds. Dunne, J.A. & Pascual, M.). Oxford University Press, pp. 27–86.
- 557 Elmasri, M., Farrell, M.J., Davies, T.J. & Stephens, D.A. (2020). A hierarchical Bayesian model for predicting
558 ecological interactions using scaled evolutionary relationships. *The Annals of Applied Statistics*, 14,
559 221–240.
- 560 Elton, C.S. (2001). *Animal Ecology*. University of Chicago Press, Chicago, IL.
- 561 Emmerson, M.C. & Raffaelli, D. (2004). Predator-prey body size, interaction strength and the stability of a real
562 food web. *Journal of Animal Ecology*, 73, 399–409.
- 563 Fedorov, V.B., Goropashnaya, A.V., Boeskorov, G.G. & Cook, J.A. (2008). Comparative phylogeography and
564 demographic history of the wood lemming (*Myopus schisticolor*): Implications for late Quaternary history
565 of the taiga species in Eurasia. *Molecular Ecology*, 17, 598–610.
- 566 Fortuna, M.A. & Bascompte, J. (2006). Habitat loss and the structure of plant-animal mutualistic networks.
567 *Ecology Letters*, 9, 281–286.
- 568 Fu, X., Seo, E., Clarke, J. & Hutchinson, R.A. (2021). Link Prediction Under Imperfect Detection:
569 Collaborative Filtering for Ecological Networks. *IEEE Transactions on Knowledge and Data Engineering*,

- 570 33, 3117–3128.
- 571 Galiana, N., Lurgi, M., Claramunt-López, B., Fortin, M.-J., Leroux, S., Cazelles, K., *et al.* (2018). [The spatial scaling of species interaction networks](#). *Nature Ecology & Evolution*, 2, 782–790.
- 573 Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010). [A framework for community interactions under climate change](#). *Trends in Ecology & Evolution*, 25, 325–331.
- 575 Godsoe, W., Murray, R. & Iritani, R. (2022). [Species interactions and diversity: A unified framework using Hill numbers](#). *Oikos*, n/a, e09282.
- 577 Golubski, A.J. & Abrams, P.A. (2011). [Modifying modifiers: What happens when interspecific interactions interact?](#) *Journal of Animal Ecology*, 80, 1097–1108.
- 579 Gomez, J.M., Verdu, M. & Perfectti, F. (2010). [Ecological interactions are evolutionarily conserved across the entire tree of life](#). *Nature*, 465, 918–U6.
- 581 Gonzalez, A. & Londoño, M.C. (2022). [Monitor biodiversity for action](#). *Science*, 378, 1147–1147.
- 582 Gonzalez-Varo, J.P. & Traveset, A. (2016). [The Labile Limits of Forbidden Interactions](#). *Trends in Ecology & Evolution*, 31, 700–710.
- 584 Gravel, D., Baiser, B., Dunne, J.A., Kopalke, J.-P., Martinez, N.D., Nyman, T., *et al.* (2019). [Bringing Elton and Grinnell together: A quantitative framework to represent the biogeography of ecological interaction networks](#). *Ecography*, 42, 401–415.
- 587 Gravel, D., Massol, F., Canard, E., Mouillot, D. & Mouquet, N. (2011). [Trophic theory of island biogeography](#). *Ecology Letters*, 14, 1010–1016.
- 589 Gravel, D., Poisot, T., Albouy, C., Velez, L. & Mouillot, D. (2013). [Inferring food web structure from predator–prey body size relationships](#). *Methods in Ecology and Evolution*, 4, 1083–1090.
- 591 Guimarães, P.R. (2020). [The Structure of Ecological Networks Across Levels of Organization](#). *Annual Review of Ecology, Evolution, and Systematics*, 51, 433–460.
- 593 Guimerà, R. & Sales-Pardo, M. (2009). [Missing and spurious interactions and the reconstruction of complex networks](#). *Proceedings of the National Academy of Sciences*, 106, 22073–22078.
- 595 Hemprich-Bennett, D.R., Oliveira, H.F.M., Le Comber, S.C., Rossiter, S.J. & Clare, E.L. (2021). [Assessing the impact of taxon resolution on network structure](#). *Ecology*, 102, e03256.

- 597 Herrera, C.M. (1989). [Pollinator abundance, morphology, and flower visitation rate: Analysis of the “quantity”](#)
598 [component in a plant-pollinator system.](#) *Oecologia*, 80, 241–248.
- 599 Hortal, J., de Bello, F., Diniz-Filho, J.A.F., Lewinsohn, T.M., Lobo, J.M. & Ladle, R.J. (2015). [Seven Shortfalls](#)
600 [that Beset Large-Scale Knowledge of Biodiversity.](#) *Annual Review of Ecology, Evolution, and Systematics*,
601 46, 523–549.
- 602 Jordano, P. (1987). [Patterns of Mutualistic Interactions in Pollination and Seed Dispersal: Connectance,](#)
603 [Dependence Asymmetries, and Coevolution.](#) *The American Naturalist*, 129, 657–677.
- 604 Jordano, P. (2016). [Sampling networks of ecological interactions.](#) *Functional Ecology*, 30, 1883–1893.
- 605 Jordano, P., Bascompte, J. & Olesen, J.M. (2003). [Invariant properties in coevolutionary networks of](#)
606 [plant–animal interactions.](#) *Ecology Letters*, 6, 69–81.
- 607 Kausrud, K.L., Mysterud, A., Steen, H., Vik, J.O., Østbye, E., Cazelles, B., *et al.* (2008). [Linking climate](#)
608 [change to lemming cycles.](#) *Nature*, 456, 93–97.
- 609 Kéfi, S., Berlow, E.L., Wieters, E.A., Joppa, L.N., Wood, S.A., Brose, U., *et al.* (2015). [Network structure](#)
610 [beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky shores.](#) *Ecology*, 96,
611 291–303.
- 612 Kéfi, S., Berlow, E.L., Wieters, E.A., Navarrete, S.A., Petchey, O.L., Wood, S.A., *et al.* (2012). [More than a](#)
613 [meal... integrating non-feeding interactions into food webs.](#) *Ecology Letters*, 15, 291–300.
- 614 Kéfi, S., Miele, V., Wieters, E.A., Navarrete, S.A. & Berlow, E.L. (2016). [How Structured Is the Entangled](#)
615 [Bank? The Surprisingly Simple Organization of Multiplex Ecological Networks Leads to Increased](#)
616 [Persistence and Resilience.](#) *PLOS Biology*, 14, e1002527.
- 617 Kissner, K.J., Forbes, M.R. & Secoy, D.M. (1997). [Rattling Behavior of Prairie Rattlesnakes \(*Crotalus viridis*](#)
618 [viridis, Viperidae\) in Relation to Sex, Reproductive Status, Body Size, and Body Temperature.](#) *Ethology*,
619 103, 1042–1050.
- 620 Klecka, J. & Boukal, D.S. (2014). [The effect of habitat structure on prey mortality depends on predator and](#)
621 [prey microhabitat use.](#) *Oecologia*, 176, 183–191.
- 622 Kopelke, J.-P., Nyman, T., Cazelles, K., Gravel, D., Vissault, S. & Roslin, T. (2017). [Food-web structure of](#)
623 [willow-galling sawflies and their natural enemies across Europe.](#) *Ecology*, 98, 1730–1730.

- 624 Kulkarni, M.A., Kweka, E., Nyale, E., Lyatuu, E., Mosha, F.W., Chandramohan, D., *et al.* (2006).
625 [Entomological Evaluation of Malaria Vectors at Different Altitudes in Hai District, Northeastern Tanzania.](#)
626 *Journal of Medical Entomology*, 43, 580–588.
- 627 Maiorano, L., Montemaggiori, A., Ficetola, G.F., O'Connor, L. & Thuiller, W. (2020). [TETRA-EU 1.0: A](#)
628 [species-level trophic metaweb of European tetrapods](#). *Global Ecology and Biogeography*, 29, 1452–1457.
- 629 McCann, K. (2007). Protecting biostructure. *Nature*, 446, 29–29.
- 630 McCann, K.S. (2011). Food webs (MPB-50). In: *Food Webs (MPB-50)*. Princeton University Press.
- 631 McLeod, A., Leroux, S.J., Gravel, D., Chu, C., Cirtwill, A.R., Fortin, M.-J., *et al.* (2021). [Sampling and](#)
632 [asymptotic network properties of spatial multi-trophic networks](#). *Oikos*, n/a.
- 633 McLeod, A.M., Leroux, S.J. & Chu, C. (2020). [Effects of species traits, motif profiles, and environment on](#)
634 [spatial variation in multi-trophic antagonistic networks](#). *Ecosphere*, 11, e03018.
- 635 Melián, C.J., Vilas, C., Baldó, F., González-Ortegón, E., Drake, P. & Williams, R.J. (2011). [Eco-evolutionary](#)
636 [Dynamics of Individual-Based Food Webs](#). In: *Advances in Ecological Research*, The Role of Body Size in
637 Multispecies Systems (ed. Belgrano, A.). Academic Press, pp. 225–268.
- 638 Morales-Castilla, I., Matias, M.G., Gravel, D. & Araújo, M.B. (2015). [Inferring biotic interactions from](#)
639 [proxies](#). *Trends in Ecology & Evolution*, 30, 347–356.
- 640 Ohlmann, M., Miele, V., Dray, S., Chalmandrier, L., O'Connor, L. & Thuiller, W. (2019). [Diversity indices for](#)
641 [ecological networks: A unifying framework using Hill numbers](#). *Ecology Letters*, 22, 737–747.
- 642 Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H., Rasmussen, C. & Jordano, P. (2010). [Missing and](#)
643 [forbidden links in mutualistic networks](#). *Proceedings of the Royal Society B: Biological Sciences*, 278,
644 725–732.
- 645 Park, J. & Newman, M.E.J. (2004). [Statistical mechanics of networks](#). *Physical Review E*, 70, 066117.
- 646 Pascual, M., Dunne, J.A. & Dunne, J.A. (2006). *Ecological Networks: Linking Structure to Dynamics in Food*
647 *Webs*. Oxford University Press, USA.
- 648 Pellissier, L., Albouy, C., Bascompte, J., Farwig, N., Graham, C., Loreau, M., *et al.* (2018). [Comparing species](#)
649 [interaction networks along environmental gradients](#). *Biological Reviews*, 93, 785–800.

- 650 Petchey, O.L., Beckerman, A.P., Riede, J.O. & Warren, P.H. (2008). [Size](#), foraging, and food web structure.
651 *Proceedings of the National Academy of Sciences*, 105, 4191–4196.
- 652 Pilosof, S., Porter, M.A., Pascual, M. & Kéfi, S. (2017). [The multilayer nature of ecological networks](#). *Nature Ecology & Evolution*, 1, 1–9.
- 653
- 654 Plagányi, É.E. & Butterworth, D.S. (2004). [A critical look at the potential of Ecopath with ecosim to assist in practical fisheries management](#). *African Journal of Marine Science*, 26, 261–287.
- 655
- 656 Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). [The dissimilarity of species interaction networks](#). *Ecology Letters*, 15, 1353–1361.
- 657
- 658 Poisot, T., Cirtwill, A.R., Cazelles, K., Gravel, D., Fortin, M.-J. & Stouffer, D.B. (2016). [The structure of probabilistic networks](#). *Methods in Ecology and Evolution*, 7, 303–312.
- 659
- 660 Poisot, T. & Gravel, D. (2014). [When is an ecological network complex? Connectance drives degree distribution and emerging network properties](#). *PeerJ*, 2, e251.
- 661
- 662 Poisot, T., Stouffer, D.B. & Gravel, D. (2015). [Beyond species: Why ecological interaction networks vary through space and time](#). *Oikos*, 124, 243–251.
- 663
- 664 Polis, G.A. (1991). [Complex Trophic Interactions in Deserts: An Empirical Critique of Food-Web Theory](#). *The American Naturalist*, 138, 123–155.
- 665
- 666 Pollock, L.J., Tingley, R., Morris, W.K., Golding, N., O'Hara, R.B., Parris, K.M., et al. (2014). [Understanding co-occurrence by modelling species simultaneously with a Joint Species DistributionModel \(JSDM\)](#).
667
668 *Methods in Ecology and Evolution*, 5, 397–406.
- 669
- 670 Pomeranz, J.P.F., Thompson, R.M., Poisot, T. & Harding, J.S. (2019). [Inferring predator–prey interactions in food webs](#). *Methods in Ecology and Evolution*, 10, 356–367.
- 671
- 672 Pringle, R.M. & Hutchinson, M.C. (2020). [Resolving Food-Web Structure](#). *Annual Review of Ecology, Evolution, and Systematics*, 51, 55–80.
- 673
- 674 Proulx, S.R., Promislow, D.E.L. & Phillips, P.C. (2005). [Network thinking in ecology and evolution](#). *Trends in Ecology & Evolution*, SPECIAL ISSUE: BUMPER BOOK REVIEW, 20, 345–353.
- 675 Pulliam, H.R. (1974). [On the Theory of Optimal Diets](#). *The American Naturalist*, 108, 59–74.

- 676 Rooney, N. & McCann, K.S. (2012). [Integrating food web diversity, structure and stability](#). *Trends in Ecology and Evolution*, 27, 40–46.
- 678 Salim, J.A., Saraiva, A.M., Zermoglio, P.F., Agostini, K., Wolowski, M., Drucker, D.P., *et al.* (2022). [Data standardization of plant–pollinator interactions](#). *GigaScience*, 11, giac043.
- 680 Sanders, D. & van Veen, F.J.F. (2012). [Indirect commensalism promotes persistence of secondary consumer species](#). *Biology Letters*, 8, 960–963.
- 682 Saravia, L.A., Marina, T.I., Kristensen, N.P., De Troch, M. & Momo, F.R. (2022). [Ecological network assembly: How the regional metaweb influences local food webs](#). *Journal of Animal Ecology*, 91, 630–642.
- 684 Schmeller, D.S., Julliard, R., Bellingham, P.J., Böhm, M., Brummitt, N., Chiarucci, A., *et al.* (2015). [Towards a global terrestrial species monitoring program](#). *Journal for Nature Conservation*, 25, 51–57.
- 686 Shaw, J.O., Dunhill, A.M., Beckerman, A.P., Dunne, J.A. & Hull, P.M. (2024). [A framework for reconstructing ancient food webs using functional trait data](#).
- 688 Singer, M.C. & McBride, C.S. (2012). [Geographic mosaics of species' association: A definition and an example driven by plant–insect phenological synchrony](#). *Ecology*, 93, 2658–2673.
- 690 Staniczenko, P.P.A., Sivasubramaniam, P., Suttle, K.B. & Pearson, R.G. (2017). [Linking macroecology and community ecology: Refining predictions of species distributions using biotic interaction networks](#). *Ecology Letters*, 20, 693–707.
- 693 Stouffer, D.B., Rezende, E.L. & Amaral, L.A.N. (2011). [The role of body mass in diet contiguity and food-web structure](#). *Journal of Animal Ecology*, 80, 632–639.
- 695 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M.J., *et al.* (2022). [Food web reconstruction through phylogenetic transfer of low-rank network representation](#). *Methods in Ecology and Evolution*, 13.
- 698 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M.J., *et al.* (2023). [Graph embedding and transfer learning can help predict potential species interaction networks despite data limitations](#). *Methods in Ecology and Evolution*, 14, 2917–2930.
- 701 Strydom, T., Catchen, M.D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., *et al.* (2021). [A roadmap towards predicting species interaction networks \(across space and time\)](#). *Philosophical Transactions of the Royal Society B-Biological Sciences*, 376, 20210063.

- 704 Thompson, R.M. & Townsend, C.R. (2005). Food-Web Topology Varies with Spatial Scale in a Patchy
705 Environment. *Ecology*, 86, 1916–1925.
- 706 Tylianakis, J.M., Tscharntke, T. & Lewis, O.T. (2007). Habitat modification alters the structure of tropical
707 host-parasitoid food webs. *Nature*, 445, 202–205.
- 708 Vázquez, D.P., Melián, C.J., Williams, N.M., Blüthgen, N., Krasnov, B.R. & Poulin, R. (2007). Species
709 abundance and asymmetric interaction strength in ecological networks. *Oikos*, 116, 1120–1127.
- 710 Vázquez, D.P., Peralta, G., Cagnolo, L. & Santos, M. (2022). Ecological interaction networks. What we know,
711 what we don't, and why it matters. *Ecología Austral*, 32, 670–697.
- 712 Wagenmakers, E.-J. & Farrell, S. (2004). AIC model selection using Akaike weights. *Psychonomic Bulletin &*
713 *Review*, 11, 192–196.
- 714 Weinstein, B.G. & Graham, C.H. (2017a). On comparing traits and abundance for predicting species
715 interactions with imperfect detection. *Food Webs*, 11, 17–25.
- 716 Weinstein, B.G. & Graham, C.H. (2017b). Persistent bill and corolla matching despite shifting temporal
717 resources in tropical hummingbird-plant interactions. *Ecology Letters*, 20, 326–335.
- 718 Whittaker, R.H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological*
719 *Monographs*, 30, 279–338.
- 720 Wood, S.A., Russell, R., Hanson, D., Williams, R.J. & Dunne, J.A. (2015). Effects of spatial scale of sampling
721 on food web structure. *Ecology and Evolution*, 5, 3769–3782.
- 722 Woodward, G., Benstead, J.P., Beveridge, O.S., Blanchard, J., Brey, T., Brown, L.E., et al. (2010). Chapter 2 -
723 Ecological Networks in a Changing Climate. In: *Advances in Ecological Research*, Ecological Networks
724 (ed. Woodward, G.). Academic Press, pp. 71–138.
- 725 Woodward, G., Brown, L.E., Edwards, F.K., Hudson, L.N., Milner, A.M., Reuman, D.C., et al. (2012). Climate
726 change impacts in multispecies systems: Drought alters food web size structure in a field experiment.
727 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 2990–2997.
- 728 Young, J.-G., Valdovinos, F.S. & Newman, M.E.J. (2021). Reconstruction of plant-pollinator networks from
729 observational data. *Nature Communications*, 12, 3911.