

The ecological interpretation of probabilistic networks

Francis Banville^{1,2,3} Tanya Strydom^{1,3} Penelope S. A. Blyth⁴ Chris Brimacombe⁵ Michael Catchen^{3,6}
Gabriel Dansereau^{1,3} Dominique Gravel^{2,3} Gracielle Higino² Thomas Malpas⁴ Hana Mayall⁴
Kari Norman¹ Timothée Poisot^{1,3}

¹ Université de Montréal ² Université de Sherbrooke ³ Quebec Centre for Biodiversity Science ⁴ University of Sheffield ⁵ University of Toronto ⁶ McGill University

Correspondance to:

Francis Banville — francis.banville@umontreal.ca

The stochastic nature of ecological interactions has led biologists to adopt a probabilistic view of ecological networks. Representing species interactions probabilistically (how likely they are to occur?) as opposed to deterministically (are they occurring?) allows a better assessment of their spatiotemporal variability and accounts for inherent uncertainties in both observations and predictions. However, despite this growing interest in probabilistic networks, guidelines for the estimation and documentation of probabilistic interaction data are still lacking. This is concerning given that our biological understanding of probabilistic interactions and their appropriate analyses depend on the often elusive methods and variables used for their estimation. We review how probabilistic interactions are defined at different spatial scales, from local interactions to regional networks (metawebs), with a strong emphasis on trophic interactions (predatory, herbivory, and parasitic interactions). These definitions are based on the distinction between the realization of an interaction at a specific time and space (local web) and its biological feasibility (metaweb). We show that different network representations have different statistical properties when it comes to common ecological applications. Specifically, we argue that probabilistic local and metawebs differ in their spatial and temporal scaling of interactions, with potential interactions in metawebs remaining constant across spatial and temporal scales. This is in contrast with the taxonomic scaling of interactions, which does not qualitatively differ between both types of networks. We suggest two approaches to inferring deterministic networks from probabilistic webs that account for these differences and argue that systematic biases arise when directly inferring local webs from subsets of metawebs. To support our arguments, we develop different case studies using open empirical data on host-parasite interactions in Europe. Our results underscore the importance of clear metadata for probabilistic ecological

networks, both at the local and regional scales, to inform the appropriate reuse of interaction data. Comprehensive metadata, providing explicit details on the sources of uncertainty in probabilistic webs, not only supports more rigorous statistical analyses but also broadens the scope of applications of interaction data. This clarity may help with the identification of optimal sampling locations to effectively minimize the uncertainty of trophic interactions.

This work is released by its authors under a CC-BY 4.0 license



Last revision: *February 11, 2024*

1 Introduction

2 As we try to navigate global biodiversity change, filling in knowledge gaps about biodiversity becomes
3 instrumental to monitoring and mitigating those changes (Gonzalez & Londoño (2022), Abrego *et al.* (2021)).
4 However, cataloging species, populations and, in particular, ecological interactions (e.g., predation, parasitism,
5 and pollination) is a substantial challenge (Polis (1991), Pascual *et al.* (2006)). There are methodological and
6 biological constraints that hinder our ability to observe all ecological interactions, leading to significant
7 uncertainties in our understanding of these interactions. For example, the spatial and temporal uncoupling of
8 species (e.g., nocturnal and diurnal species coexisting in the same space without interacting due to differences
9 in the timing of their daily activities, Jordano (1987)) and the large number of rare and cryptic interactions in a
10 community contribute to these uncertainties (Jordano (2016)). More generally, a handful of conditions must be
11 satisfied for an interaction to be observed locally. First, both species must have overlapping geographic ranges,
12 i.e. they must co-occur within the region of interest (Blanchet *et al.* (2020)). Second, they must have some
13 probability of meeting. Probabilities of interspecific encounters are typically low, especially for rare species
14 with low relative abundances (Canard *et al.* (2012)). The probability that species meet each other also depends
15 on their biological characteristics, such as the synchronization of their phenology (Olesen *et al.* (2010), Singer
16 & McBride (2012)) and their discoverability (e.g., Broom & Ruxton (2005)). Finally, when species do come
17 into contact, an interaction occurs only if their traits are locally compatible (Poisot *et al.* (2015)), including but
18 not limited to their body phenotypes (Bolnick *et al.* (2011), Stouffer *et al.* (2011), Gravel *et al.* (2013)) and
19 behavioral choices (Pulliam (1974), Choh *et al.* (2012)). Interactions may also be influenced by the presence or
20 prevalence of a third species (e.g., of a more profitable prey species, Golubski & Abrams (2011), Sanders & van
21 Veen (2012)). Documenting the location and timing of interactions becomes even more difficult when
22 accounting for the spatiotemporal variability of ecological interactions (Poisot *et al.* (2012), Poisot *et al.*
23 (2015)). Environmental factors, such as temperature (Angilletta *et al.* (2004)), drought (Woodward *et al.*
24 (2012)), climate change (Gilman *et al.* (2010), Woodward *et al.* (2010), Araujo *et al.* (2011)), and habitat
25 modifications (Tylianakis *et al.* (2007)), contribute to this spatiotemporal variability by impacting species
26 abundance and traits. Even after satisfying all these conditions, there remains a possibility that the interaction
27 does not occur locally, either due to the intricate nature of the system or simply by chance. If it does occur, it
28 might still go unnoticed, particularly if it happens infrequently. In this context, it is unsurprising that our
29 knowledge of ecological interactions remains limited (Hortal *et al.* (2015)) despite extensive biodiversity data
30 collection (Schmeller *et al.* (2015)).

31 Knowing the biological capacity of two species to interact directly (via e.g., trophic interactions) is necessary
32 but not sufficient for inferring their interaction at a specific time and space. The recognition of the intrinsic
33 variability of species interactions has led ecologists to expand their representation of ecological networks (also
34 known as ecological webs) to include a probabilistic view of interactions (Poisot *et al.* (2016), Dallas *et al.*
35 (2017), Fu *et al.* (2021)). This different perspective allows us to fill in the Eltonian shortfall (Hortal *et al.*
36 (2015)) by modeling the probability of detecting interactions, which can be an important tool for directing
37 efforts and taking action, especially in places where access and resources for research are scarce. The
38 probabilistic representation of interactions is thus far limited to direct interactions, which are conceptually and
39 mathematically analogous regardless of their biological type (e.g., predation and pollination). This is in contrast
40 with indirect interactions (e.g., interspecific competition), which arise from distinct ecological processes and are
41 often not directly observable (Kéfi *et al.* (2015), Kéfi *et al.* (2016)). Representing direct interactions
42 probabilistically enables us to capture the spatiotemporal variability of the aforementioned ecological processes
43 and the uncertainties associated with their measurement. As opposed to binary deterministic webs, in which
44 interactions are regarded as either occurring or not, probabilistic webs, within a Bayesian framework, express
45 our degree of belief (or confidence) regarding the occurrence or observation of interactions. Based on the scale
46 at which they are estimated, probabilistic interactions may reflect our level of confidence in whether interactions
47 will be observed, realized locally, or biologically feasible. As an illustration, we could outline a situation in
48 which there is a 50% certainty that an interaction occurs 50% of the time, or that there is a 50% certainty that it
49 simply occurs. Our level of confidence should be more definitive (approaching either 0 or 1) as we extend our
50 sampling to a broader area and over a longer time period, thereby diminishing the uncertainty of the interactions
51 (but not necessarily the estimation of their variability). In the broadest sense, binary networks are also a type of
52 probabilistic network, in which the numerical value of an interaction is restrained to 0 (non-occurring) or 1
53 (occurring). Yet, for the sake of clarity, we omit binary networks from our discussion of probabilistic networks
54 in this contribution. In probabilistic webs, only forbidden interactions (i.e., interactions prohibited by biological
55 traits or species absence, Jordano *et al.* (2003), Olesen *et al.* (2010)) have a probability value of zero by default,
56 provided that intraspecific trait variability is considered (Gonzalez-Varo & Traveset (2016)).

57 By accounting for the uncertainty of interactions, probabilistic webs may provide a more realistic portrait of
58 species interactions and network structure (i.e. community-level properties), which are major drivers of the
59 functioning, dynamics, and resilience of ecosystems worldwide (Proulx *et al.* (2005), McCann (2007), McCann
60 (2011), Rooney & McCann (2012)). Moreover, the application and development of computational methods in

network ecology, often based on a probabilistic representation of interactions, can alleviate (and guide) the sampling efforts required to document species interactions (Strydom *et al.* (2021)). For example, statistical models can be used to estimate the uncertainty of pairwise interactions (Cirtwill *et al.* (2019)) and the probability of missing (false negatives) and spurious (false positives) interactions (Guimerà & Sales-Pardo (2009)). Considering the high rate of false negatives in species interaction data due to the difficulty of witnessing rare interactions (Catchen *et al.* (2023)), these models could inform the identification of priority sampling locations of ecological webs where data collection would yield the most valuable information, thereby reducing errors. Optimization models for sampling locations have mostly found applications in biological systems that are not networks, such as identifying priority sampling sites for disease hotspots (Andrade-Pacheco *et al.* (2020)), but there is substantial promise in applying them to probabilistic ecological interactions.

Statistical models can also be used to generate predictions of ecological webs without prior knowledge of pairwise interactions, for instance using body size (Petchey *et al.* (2008), Gravel *et al.* (2013)), phylogeny (Elmasri *et al.* (2020), Strydom *et al.* (2022)), or a combination of niche and neutral processes (Bartomeus *et al.* (2016), Pomeranz *et al.* (2019)) for inference. Topological null models, which generate probabilistic networks by preserving chosen characteristics of the binary adjacency matrix while intentionally omitting others (Bascompte *et al.* (2003), Fortuna & Bascompte (2006)), serve as other examples of common probabilistic network models. Null models can be used to produce underlying distributions of network measures for null hypothesis significance testing. Many measures have been developed to describe the structure (Poisot *et al.* (2016)) and diversity (Ohlmann *et al.* (2019), Godsoe *et al.* (2022)) of probabilistic webs. These models and measures support the use of this approach for the study of a wide range of ecological questions, from making better predictions of species distribution (Cazelles *et al.* (2016)) to forecasting the impact of climate change on ecological webs (Gilman *et al.* (2010)).

The lack of clear guidelines on the use of probabilistic interaction data is worrisome, both for data producers and re-users who generate and manipulate these numbers. This is concerning because sampling strategies and decisions regarding network construction can affect our understanding of network properties (Brimacombe *et al.* (2023)). Besides methodological difficulties that may arise when assessing probabilistic interactions, a precise definition of probabilistic interactions appears to be lacking, making the estimation and use of these data more difficult. We aim to take a step back by outlining different ways in which probabilistic interactions are defined and used in network ecology. We distinguish two broad categories of probabilistic webs that necessitate distinct approaches when applied to key ecological questions: local webs describing probabilities of realized

91 interactions, and regional webs (metawebs) describing probabilities of potential interactions. We highlight the
92 distinctions in the ecological meaning of these two representations and show that they yield different statistical
93 outcomes regarding e.g. the spatial and temporal scaling of interactions and the prediction of binary webs
94 across space. Moreover, there is currently no metadata standard that could guide the documentation of all types
95 of probabilistic interactions (although see e.g., Salim *et al.* (2022) who discuss data standards for deterministic
96 mutualistic webs). Well-defined metadata for probabilistic webs would support more adequate manipulation
97 and integration of interaction data from different sources and guard against possible misinterpretations arising
98 from ambiguous definitions of probabilistic networks. These metadata should outline the nature (i.e., local or
99 regional) and type (e.g., predatory or pollination) of the interactions, provide information regarding the
100 taxonomic level, identities, and characteristics (e.g., life stages) of the individuals involved in an interaction,
101 present the mathematical formulation of probabilities, including clearly identified conditional variables (e.g.,
102 spatial and temporal scales), and describe the methods and contexts (e.g., location, time, environmental
103 conditions) in which interactions were estimated. Inadequately documented probabilistic interaction data should
104 be used with caution when analyzing ecological webs. Our observations and advice can be applied to many
105 types of ecological networks representing direct interactions, from food webs to host-virus networks.

106 **Probabilistic representations of interactions**

107 One of the first aspects to take into consideration when estimating or interpreting probabilities of interactions is
108 knowing if they describe potential or realized interactions, as these two types of interactions have distinct
109 conceptual underpinnings and sources of uncertainty. A potential (regional) interaction is defined as the
110 biological capacity of two taxa to interact (i.e., the probability that they *can* theoretically interact) whereas a
111 realized (local) interaction is the materialization or observation of this interaction in a well-defined space and
112 time (i.e., the probability that they interact locally). Here, we use the terms *metaweb* (Dunne (2006)) to
113 designate regional webs of potential interactions and *local webs* (Poisot *et al.* (2012)) for those of realized
114 interactions. Metawebs are the network analogs of the species pool, where local webs originate from a subset of
115 both species (nodes) and interactions (edges) of the regional metaweb (Saravia *et al.* (2022)). Without clear
116 documentation, it can be challenging to know if published probabilistic webs describe local or regional
117 interactions (tbl. 1 provides examples of studies employing both types of probabilistic networks), or if so-called
118 probabilities are in reality *interaction scores* (i.e., non-probabilistic quantitative interactions analogous to

119 interaction strengths). When probabilistic regional interactions are used and interpreted incorrectly as local
120 interactions (and conversely), this may generate misleading findings during data analysis. We believe that a
121 better understanding of the differences, similarities, and interconnections between these two probabilistic
122 representations of ecological webs would alleviate interpretation errors (e.g., when studying network-area
123 relationships) and facilitate a more adequate utilization of interaction data.

124 Pairwise interactions: the building blocks of ecological networks

125 Local and metawebs, like any type of network, are made of nodes and edges that can be represented at different
126 levels of organization and precision. The basic units of ecological networks are individuals that interact with
127 each other (e.g., by predation in food webs, Elton (2001)), forming individual-based networks (Melián *et al.*
128 (2011)). The aggregation of these individuals into more or less homogeneous groups (e.g., populations, species,
129 families, feeding guilds) allows us to represent nodes at broader taxonomic scales, which affects our
130 interpretation of the properties of these systems (Guimarães (2020), Hemprich-Bennett *et al.* (2021)).

131 Edges linking nodes can describe a variety of interaction measures. Ecologists have traditionally represented
132 interactions as binary objects that were considered realized after observing at least one individual from group i
133 interact with at least another individual from group j . In a binary adjacency matrix B , the presence or absence
134 of an interaction $B_{i \rightarrow j}$ between two taxa can be viewed as the result of a Bernoulli process
135 $B_{i \rightarrow j} \sim \text{Bernoulli}(P(B_{i \rightarrow j}))$, with $P(B_{i \rightarrow j})$ being the probability of interaction that characterizes our limited
136 knowledge of the system and its intrinsic spatiotemporal variability. In probabilistic networks, $P(B_{i \rightarrow j})$ are edge
137 values, and the only two possible outcomes are the presence ($B_{i \rightarrow j} = 1$) or absence ($B_{i \rightarrow j} = 0$) of an interaction
138 between each pair of nodes. Depending on the type of probabilistic network (local or metaweb), the
139 mathematical formulation and interpretation of stochastic parameters like $P(B_{i \rightarrow j})$ can be linked to
140 environmental and biological factors such as species relative abundance, traits, area, and time (tbl. 1), for
141 example using logistic regression with a logit link function with continuous explanatory variables. Predicting
142 the number of local webs in which the interaction occurs can be achieved by using a Binomial distribution,
143 assuming a constant probability of interaction and independence between networks (trials). When considering
144 uncertainties around the estimation of $P(B_{i \rightarrow j})$, a Beta distribution can also be used to encompass all possible
145 probability values. In that case, a Beta-Binomial distribution can be used to predict the number of networks in
146 which the interaction occurs. Observing an interaction between two taxa at a given location and time provides
147 important information that can be used to update previous estimates of $P(B_{i \rightarrow j})$, informing us on the biological

capacity of both taxa to interact and the environmental conditions that enabled them to interact locally.

Even though binary webs constitute a highly valuable source of ecological information (Pascual *et al.* (2006)), they overlook important factors regarding interaction strengths. Represented in a quantitative adjacency matrix W as numbers not confined to the $[0, 1]$ range, interaction strengths better describe the energy flows, demographic impacts or frequencies of interactions between nodes (Berlow *et al.* (2004), Borrett & Scharler (2019)), with $W_{i \rightarrow j}$ being a natural \mathbb{N} or real \mathbb{R} number depending on the measure. For example, they may represent local interaction rates between pairs of taxa (e.g., the flower-visiting rates of pollinators in a mutualistic network, Herrera (1989)). When interaction strengths characterize predation pressure on prey taxa in food webs, they can serve as good estimators of the parameters describing species interactions in a Lotka-Volterra model (e.g., Emmerson & Raffaelli (2004)). The extra amount of ecological information in quantitative networks typically comes at a cost of greater sampling effort and data volume in predictive models (Strydom *et al.* (2021)), which can lead to relatively high levels of uncertainties when inferring quantitative webs with limited data.

Just like binary networks, the uncertainty and spatiotemporal variability of interaction strengths can be represented probabilistically. However, the need to estimate the probability distribution of all possible values of interaction strengths can make the inference of probabilities more challenging in quantitative webs compared to binary webs, which require only one probability estimate for each interaction. Interaction strengths can follow various probability distributions depending on the measure used. For instance, they can follow a Poisson distribution $W_{i \rightarrow j} \sim \text{Poisson}(\lambda_{i \rightarrow j})$ when predicting frequencies of interactions between pairs of nodes, with $\lambda_{i \rightarrow j}$ being the expected rate at which individuals of taxa i and j interact (e.g., the average number of prey j consumed by all predators i in a given time period). The Poisson distribution can also be 0-inflated after initially modeling non-interacting taxa (e.g., Boulangeat *et al.* (2012) employ a 0-inflated model to analyze species abundance following the modeling of species presence and absence), which constitute the majority of taxa pairs in most local webs due to their typically high sparseness (Jordano (2016)). Because of the methodological difficulties typically encountered when building deterministic quantitative webs (which are only partially mitigated by models such as Ecopath, Plagányi & Butterworth (2004)), binary webs, which are easier to sample (Jordano (2016)) and predict (Strydom *et al.* (2021)), have been more frequently studied and modeled. Moreover, most published probabilistic networks and methods describe probabilistic interactions whose outcome is binary (whether interaction probabilities are regarded as constant or variable, e.g. represented by a Beta distribution), which underlines the need for better guidelines regarding the interpretation and

178 manipulation of these types of webs first. For these reasons, our primary focus in this contribution will be on
179 addressing the challenges in interpreting and using interaction probabilities in Bernoulli distributions, in both
180 probabilistic local and metawebs.

181 Local webs: communities interacting in space and time

182 Probabilistic local webs describe how likely taxa are to interact at a given location and time period (i.e.,
183 interactions are contingent upon the environmental and biological conditions of the community). In local webs,
184 edges commonly represent our degree of belief that two taxa interact in nature, but can also represent the
185 probability of *observing* this interaction (Catchen *et al.* (2023)). For example, Gravel *et al.* (2019) used a
186 dataset of binary local European food webs of willow-galling sawflies and their natural enemies, all referenced
187 in space and time and consisting of similar species, to infer the probabilities of locally observing interactions
188 between co-occurring species. This was achieved by situating local webs within the context of environmental
189 variables, i.e. by including temperature and precipitation as conditional variables in some of their models.

190 We define space as the collection of geographic coordinates (x, y, z) , with (x, y) representing longitude and
191 latitude coordinates, and z denoting either altitudes or depths. These point coordinates delineate the spatial
192 boundaries of the system, which may be portrayed as a polyhedron. Ecological interactions may exhibit
193 variations along latitudinal and altitudinal gradients, as evidenced by changes in hummingbird-plant
194 interactions (Weinstein & Graham (2017b), Weinstein & Graham (2017a)) and mosquito biting rates (e.g.,
195 Kulkarni *et al.* (2006)) at different elevations. In contrast, time is defined as the specific time period within
196 which interactions were either observed or predicted. Even though space and time are continuous variables that
197 should yield probability *densities* of interactions (i.e., relative likelihoods of interactions occurring at
198 infinitesimal locations and instants in time), these definitions enable them to be conceptualised as distinct
199 patches and time segments. Treating space and time as discrete dimensions aligns with the common sampling
200 methods of ecological networks and provides actual probabilities of interactions, which can be obtained by
201 integrating probability densities. Furthermore, we can quantify both an area A and a duration t , which can be
202 readily used in spatiotemporal analyses of ecological networks. For example, when studying network-area
203 relationships (NAR, Galiana *et al.* (2018)), we anticipate that local probabilities of interactions scale positively
204 with area and duration because taxa have more opportunities to interact as these dimensions expand.

205 The probability that two taxa i and j interact in a local web $L_{x,y,z,t}$ (spatial and temporal subscripts hereafter

206 omitted for clarity) can be conditional on many environmental and biological factors. One of these is their
207 co-occurrence $C_{i,j}$, which is usually a Boolean describing if the geographic distributions of both taxa overlap
208 within the study area. Co-occurrence can be modeled probabilistically, in which case it may conform to a
209 Bernoulli distribution $C_{i,j} \sim \text{Bernoulli}(P(C_{i,j}))$, where the probability of co-occurrence $P(C_{i,j})$ can be
210 estimated through the application of joint species distribution models (e.g., Pollock *et al.* (2014)). Considering
211 that the probability that two non-co-occurring taxa interact locally is zero (i.e., $P(L_{i \rightarrow j}|C_{i,j} = 0) = 0$), the
212 probability of local interaction can be obtained by multiplying the probability of interaction given
213 co-occurrence with the probability of co-occurrence:

$$P(L_{i \rightarrow j}) = P(L_{i \rightarrow j}|C_{i,j} = 1) \times P(C_{i,j} = 1). \quad (1)$$

214 Other important factors that can impact interaction probabilities at the local scale are taxa relative abundance
215 (Canard *et al.* (2012)) and traits (Poisot *et al.* (2015)), as well as environmental factors such as temperature
216 (Angilletta *et al.* (2004)), precipitation (Woodward *et al.* (2012)), habitat structure (Klecka & Boukal (2014)),
217 and presence of other interacting taxa in the network (Pilosof *et al.* (2017), Kéfi *et al.* (2012)), as described
218 above. Here, we use the variable $\Omega_{x,y,z,t}$ (hereafter simply Ω) to describe the biological and ecological context
219 in which interaction probabilities were estimated. For example, if a research team conducts a mesocosm
220 experiment to estimate interaction probabilities between predators and prey with and without shelters (a place
221 that offers refuge and protection for prey, shielding them from predators), Ω would represent the presence or
222 absence of these shelters. Like co-occurrence, Ω can also be modeled probabilistically when the stochasticity or
223 uncertainty of environmental and biological factors is considered. In sum, Ω represents all ecological and
224 biological variables that were taken into consideration when measuring interaction probabilities and is,
225 therefore, a subset of all factors impacting ecological interactions. It is a vector of random variables whose
226 values determine the overall uncertainty of an interaction.

227 The probability that two taxa i and j interact in a local web L can thus be conditional on the area (or volume) A ,
228 the time interval t , their co-occurrence $C_{i,j}$ and chosen environmental and biological conditions Ω . Although
229 these variables are associated with distinct questions or mechanisms related to ecological interactions, they may
230 covary with each other, such as the possible dependence of $C_{i,j}$ and Ω on spatial and temporal scales. When
231 estimating interaction probabilities using e.g. a generalized linear model with multiple explanatory variables
232 that might not be independent, it may become important to address collinearity. Using variable selection

233 techniques, for instance, may be necessary before fitting the model to mitigate this issue. The probability of
 234 local interaction is described by the following expression when all these conditional variables are included:

$$P(L_{i \rightarrow j} | A, t, C_{i,j}, \Omega) \quad (2)$$

235 The representation of the local context in which probabilities are estimated and the variables that should be
 236 taken into consideration depend on the study system, the objectives of the study, and the resources available to
 237 the researchers. In other words, these variables do not systematically need to be accounted for. For example, in
 238 Gravel *et al.* (2019), for the purpose of model comparison, some models incorporated environmental variables
 239 as conditional factors to estimate interaction probabilities, while others did not. When accounted for, these
 240 variables should be clearly described in the documentation of the data (Brimacombe *et al.* (2023)),
 241 preferentially in mathematical terms to avoid any confusion in their interpretation and to limit manipulation
 242 errors during their re-use. For instance, ecologists should be explicit about their consideration of co-occurrence
 243 in their estimation of local interaction probabilities. Indeed, it is important to specify if probability values are
 244 conditional $P(L_{i \rightarrow j} | C_{i,j} = 1)$ or not $P(L_{i \rightarrow j})$ on co-occurrence since this can significantly impact the
 245 interpretation and analysis of the data. In [tbl. 1](#), we present examples of studies that used these diverse
 246 formulations of probabilistic interactions and conditional variables.

Table 1: Mathematical expression of probabilistic interactions. The probability of interaction between two taxa i and j is interpreted differently in a metaweb M of potential interactions and a local web L of realized interactions. Each expression includes a different conditional variable, when applicable. An example of a study employing each of these notations and conditional variables is provided, with the specific variables used indicated in parentheses. The study marked with an asterisk has been carried out on binary webs. Note that interaction probabilities can be contingent upon multiple conditional variables, or none at all. Additionally, local interaction probabilities might represent the probability of observing an interaction, not necessarily its actual occurrence.

| Expression | Type | Outcome | Reference |
|----------------------------|----------|---|--|
| $P(M_{i \rightarrow j})$ | regional | biological feasibility of the interaction | Strydom <i>et al.</i> (2022) |
| $P(L_{i \rightarrow j})$ | local | realization of the interaction | Fortuna & Bascompte (2006) (null model) |
| $P(L_{i \rightarrow j} A)$ | local | realization of the interaction in a given area or volume | Galiana <i>et al.</i> (2018) * |
| $P(L_{i \rightarrow j} t)$ | local | realization of the interaction during a given time period | Weinstein & Graham (2017a) |

| Expression | Type | Outcome | Reference |
|--|-------|--|---|
| $P(L_{i \rightarrow j} C_{i,j})$ | local | realization of the interaction given that the taxa co-occur | Gravel <i>et al.</i> (2019) |
| $P(L_{i \rightarrow j} \Omega)$ | local | realization of the interaction given environmental conditions | Gravel <i>et al.</i> (2019) (temperature and precipitation) |
| $P(L_{i \rightarrow j} M_{i \rightarrow j})$ | local | realization of the interaction given that the taxa can biologically interact | this study |

247 Metawebs: regional catalogs of interactions

248 Metawebs are networks of potential interactions that have been designed for broad spatial, temporal, and
 249 taxonomic scales (e.g., species food webs at the continental scale). They represent the probability that taxa can
 250 biologically interact regardless of their co-occurrence and local environmental conditions. Regional interactions
 251 are by definition context-independent, i.e. they are not measured at a specific location and time. In contrast with
 252 probabilistic local webs, which represent the stochasticity of interactions occurring in nature, probabilistic
 253 metawebs measure our degree of belief in the capacity of two taxa to interact (i.e., the probability that their
 254 traits could support an interaction in the right conditions). Consequently, although *neutrally* forbidden
 255 interactions (i.e., improbable interactions between rare species, Canard *et al.* (2012)) tend to have low
 256 probability values in local webs, they may exhibit a higher probability in the metaweb. Potential interactions
 257 describe the probability that there exists at least one combination of phenotypes of taxa i and j that can interact
 258 with each other if they were to encounter. For example, a piscivorous species may have the capacity to interact
 259 with a potential prey species if they share at least one matching combination of body sizes, such as an adult
 260 predator being able to consume juvenile prey. To reduce any biases in our calculation of this probability, it is
 261 crucial to ensure that the set of traits sampled or considered accurately reflects the overall trait distribution in
 262 both taxa. This enhances our confidence in concluding that the taxa cannot interact when no positive
 263 observation has been made and increases our capacity to detect an interaction when it is biologically feasible. In
 264 addition to pairwise interactions, the extent of sampling effort also contributes to shaping our assessment of
 265 metaweb properties, as sampling a larger number of local webs allows us to capture more regional interactions
 266 (McLeod *et al.* (2021)). A probability of potential interaction in a metaweb M can be expressed as

$$P(M_{i \rightarrow j}), \quad (3)$$

267 which, in contrast with local webs, is never conditional on any spatial, temporal, co-occurrence or
268 environmental variables (tbl. 1).

269 Starting from a selected set of taxa, which are usually distributed within a broad region of interest, metawebs
270 can be built using different data sources, including literature review (e.g., Maiorano2020Tetraeu), aggregated
271 local web data (e.g., Gravel *et al.* (2019), Saravia *et al.* (2022)), and predictive models (e.g., Strydom *et al.*
272 (2022), Shaw *et al.* (2024)). Every pair of taxa that have confidently been observed to interact at least once can
273 be given a probability of 1 (i.e., $P(M_{i \rightarrow j}) = 1$) since we know that they *can* interact. This is not the case in local
274 probabilistic webs, in which interaction events usually remain stochastic (i.e., $P(L_{i \rightarrow j}) < 1$) after empirically
275 observing interactions because of their intrinsic spatiotemporal variability. Similarly, although rare interactions
276 typically have low probabilities in local webs, they can have high probabilities in metawebs when the traits of
277 both taxa are congruent. On the other hand, interactions that were never observed can have low probability
278 values in both local and metawebs, going as low as 0 for forbidden links (i.e., interactions forbidden by
279 biological traits). However, many observations of interactions are in reality false positives because of
280 observation errors due to taxonomic misidentifications and ecological misinterpretations (e.g., due to cryptic
281 species and interactions, Pringle & Hutchinson (2020)). Likewise, forbidden interactions can be false negatives
282 in metawebs, e.g. if they have been assessed for specific phenotypes, locations or time. Employing Bayesian
283 models, whether they are mechanistic or phenomenological, could improve the accuracy of our estimation of
284 interaction probabilities within both local and regional webs (e.g., Bartomeus *et al.* (2016), Cirtwill *et al.*
285 (2019)). This improvement is achieved in metawebs by leveraging prior information regarding the feasibility of
286 interactions along with the empirical data on observed interactions.

287 Properties of probabilistic webs

288 Probabilistic local and metawebs differ in their type of interactions (i.e., realized or potential) and in the
289 conditional variables upon which interaction values depend. These differences are significant as they influence
290 the characteristics of probabilistic networks. Neglecting to consider them may result in misleading results and
291 interpretation errors when analyzing the properties of probabilistic webs, which could be particularly

292 problematic when addressing crucial ecological questions about networks. Here we compare the characteristics
293 of local and metawebs through the presentation of four applications of probabilistic interactions: (1) describing
294 their spatial and temporal scaling, (2) describing their taxonomic scaling, (3) sampling for binary webs, and (4)
295 reconstructing probabilistic local webs from metawebs. All code and data to reproduce these analyses are
296 available at the Open Science Framework (TBD).

297 Host-parasite network data

298 We use the collection of tripartite host-parasite networks sampled across Europe, created by Kopelke *et al.*
299 (2017), in the following case studies. This dataset contains well-resolved binary local interactions between
300 willows (52 species), willow-galling sawflies (96 species), and their parasitoids (126 species). Given its
301 replicated webs spanning large spatiotemporal scales, this dataset is well-suited for analyzing a variety of
302 ecological hypotheses and processes. Out of a total of 374 local webs, we retained those containing at least 5
303 species, resulting in a set of 233 georeferenced local webs (networks sampled within areas of 0.1 to 0.3 km²
304 during June and/or July spanning 29 years). We built a binary metaweb by aggregating all local interactions,
305 which gave us a regional web composed of 274 species and 1080 interactions. In the first two panels of fig. 1,
306 we show how the dissimilarity of interactions between common species (β_{OS}) and the dissimilarity in species
307 composition (β_S) between the metaweb and aggregated local webs (Poisot *et al.* (2012)) vary with the number
308 of sampled local webs. This shows that networks of local interactions are highly dissimilar from the metaweb,
309 both in terms of species and interactions, especially when only a limited number of sites has been sampled.
310 Both dissimilarity indices were calculated based on the number of items shared by the two webs (c_{LM}) and the
311 number of items unique to the metaweb (u_M) and to the aggregated local web (u_L). The β_S dissimilarity index
312 uses species (nodes) as items being compared, while the β_{OS} index assesses dissimilarity based on interactions
313 between shared species (Poisot *et al.* (2012)). Both indices were calculated following the β_W index of
314 Whittaker (1960):

$$\beta_W = \frac{c_{LM} + u_L + u_M}{(2c_{LM} + u_L + u_M)/2} - 1. \quad (4)$$

315 [Figure 1 about here.]

316 We converted these binary networks into probabilistic ones using models based on simple assumptions. Our
317 models are not designed to estimate the exact values of probabilistic interactions. Instead, their purpose is to

318 create plausible networks that serve as illustrative examples to highlight distinctions between probabilistic local
319 and metawebs. We created two probabilistic metawebs by employing constant false positive and false negative
320 rates for all regional interactions. In the first metaweb, we set both false positive and false negative rates to zero
321 to prevent artificially inflating the total number of links, enabling a more accurate comparison with binary webs.
322 This gave us a probability of regional interaction of 1 when at least one interaction has been observed and of 0
323 in the absence of any observed interaction between a given pair of species. In the second metaweb, we
324 introduced a 5% false positive rate to account for spurious interactions and a 10% false negative rate to address
325 the elevated occurrence of missing interactions in ecological networks (Catchen *et al.* (2023)). We believe these
326 rates represent reasonable estimates of missing and spurious potential interactions, but confirming their
327 accuracy is challenging due to the unavailability of data on the actual feasibility of interaction. Observed
328 interactions were thus given a probability of regional interaction of 95%, whereas unobserved ones were
329 assigned a probability of 10%.

330 To build probabilistic local webs, we first recognize that local interactions must initially be biologically feasible
331 before occurring at a specific time and space. A local probability of interaction $P(L_{i \rightarrow j})$ can be expressed as the
332 product of the probability of local interaction given that the two taxa can potentially interact $P(L_{i \rightarrow j}|M_{i \rightarrow j} = 1)$,
333 which we denote as $P(L|M)$ for the sake of simplicity, with their probability of regional interaction $P(M_{i \rightarrow j})$:

$$P(L_{i \rightarrow j}) = P(L_{i \rightarrow j}|M_{i \rightarrow j} = 1) \times P(M_{i \rightarrow j} = 1). \quad (5)$$

334 We built the probabilistic local webs from the binary ones by using the probabilistic metawebs and a constant
335 value of $P(L|M)$ across interactions. Low values of $P(L|M)$ indicate that feasible interactions rarely occur
336 locally, intermediate values around 50% suggest considerable spatiotemporal variability, while high values
337 indicate that regional interactions are nearly always realized locally. Following eq. 5, the local probability of
338 interaction between a given pair of taxa consistently remained equal to or below their probability of regional
339 interaction.

340 In the last two panels of fig. 1, we show how the aggregated number of links and connectance (i.e., the
341 proportion of all of the non-forbidden links that are realized) scale with the number of sampled local
342 probabilistic webs, according to different values of $P(L|M)$. When aggregating local probabilistic webs, the
343 constancy of the probability of regional interaction across the entire study area means that any rise in the
344 probability of local interaction is solely attributable to an increase in $P(L|M)$. The probability $P(L_{1,2}|M)$ of

345 local interaction among potentially interacting species in an aggregated web $L_{1,2}$ is obtained by:

$$P(L_{1,2}|M) = 1 - (1 - P(L_1|M)) \times (1 - P(L_2|M)), \quad (6)$$

346 where $P(L_1|M)$ and $P(L_2|M)$ are the probabilities of local interaction among two potentially interacting species
347 in the subnetworks L_1 and L_2 , respectively. This equation provides the probability that the interaction is realized
348 in either (1) exclusively the local web L_1 , (2) exclusively the local web L_2 or (3) both local webs, given that the
349 two taxa have the biological capacity to interact.

350 By comparing the scaling relationships observed in binary and probabilistic webs, fig. 1 illustrates that high
351 values of $P(L|M)$ lead to systematic overestimations in the number of links and connectance, especially when
352 $P(L|M) = 1$ (corresponding to the scenario where local probabilities of interactions are equivalent to the
353 probabilities of regional interactions). However, these biases tend to diminish as the number of sampled webs
354 increases.

355 Spatial and temporal scaling of interactions

356 The investigation of network-area relationships and interaction accumulation curves is an important area of
357 research in network ecology. First, network-area relationships elucidate the scaling of network properties (such
358 as modularity and trophic chain lengths) with spatial scale (e.g., Wood *et al.* (2015), Galiana *et al.* (2018)). The
359 variations in network structure across spatial scales may stem from the scaling of species richness (species-area
360 relationships, SARs) and the number of interactions (Brose *et al.* (2004)) with the sampled area. Additionally,
361 ecological processes unfolding at distinct spatial scales, such as the spatial variability in local community
362 composition resulting from different sequences of extinction and colonization events, can also contribute to this
363 variation (Galiana *et al.* (2018)). Next, interaction accumulation curves describe the scaling of the number of
364 observed interactions with sampling effort (Jordano (2016)). Sampling effort, which may correspond to the
365 duration of the sampling period used to construct the network, can impact connectance (Bersier *et al.* (1999))
366 and various measures of network structure (Banašek-Richter *et al.* (2004), McLeod *et al.* (2021)). Apart from
367 sampling effort, the temporal scaling of interactions also elucidates how network structure changes with the
368 temporal resolution of the network (Poisot *et al.* (2012)), acknowledging that distinct interactions take place
369 over time, ranging from short-term fluctuations of interactions to long-term trends. As probabilistic local webs
370 may explicitly account for the spatiotemporal variability of interactions, they offer a distinct approach to

371 investigating the scaling of network structure with space and time, in contrast to binary and quantitative webs,
372 by making the stochasticity of interactions the focal point of the modeling process.

373 Local and metawebs exhibit distinct relationships with spatial and temporal scales. On one hand, probabilistic
374 metawebs, being independent of any local context, feature regional interactions that do not scale with space and
375 time. This is because regional interactions depend solely on the biological capacity of two taxa to interact,
376 regardless of their co-occurrence and specific environmental conditions. However, probabilities of potential
377 interactions may change (tending to become more definitive) upon updating previous estimates with increased
378 sampling effort, even though they do not vary in a specific direction with the spatial and temporal extent
379 (boundaries) of the network. The probability of two taxa potentially interacting should theoretically be the same
380 in all metawebs in which they are present, provided that the data and methods used for estimation are consistent.

381 For example, if a smaller metaweb M_1 is derived from a larger metaweb M_0 by selecting the subset of taxa
382 present in the region described by M_1 and retaining all their interactions, their probabilities of interaction
383 should be identical regardless of scale, i.e. $P(M_{1,i \rightarrow j}) = P(M_{0,i \rightarrow j})$. With a larger or equal number of taxa in M_0
384 as compared to M_1 , the total number of interactions is expected to be higher or at least equal in the larger web,
385 even though pairwise probabilities remain identical.

386 On the other hand, local interactions scale both spatially and temporally, given that they have more
387 opportunities to be realized and observed in larger areas and longer durations. This is attributed to factors such
388 as a higher number of individuals, greater trait variations, and increased opportunities for encounters, as
389 highlighted by McLeod *et al.* (2020). For example, if a local probabilistic web L_1 with an area A_1 is derived
390 from a larger web L_0 with an area A_0 , and A_1 is entirely nested within A_0 , interaction probabilities should be
391 lower in the smaller web, i.e. $P(L_{1,i \rightarrow j}|A_1 < A_0) \leq P(L_{0,i \rightarrow j}|A_0)$. However, if A_1 and A_0 are disjoint, interaction
392 probabilities could be higher in the smaller area, contingent upon their environmental and biological conditions.
393 Likewise, interaction probabilities are expected to be lower in webs with shorter durations when time intervals
394 are nested. In fig. 2, we show how the expected number of local host-parasite interactions scales with area,
395 represented as an expanding latitudinal window, in comparison with regional interactions. Even though we
396 employed local probabilities of interactions equal to regional interactions for the purpose of comparison (i.e.,
397 using $P(L|M) = 1$ here), we notice that the total number of regional interactions scales more rapidly than local
398 interactions. This is because numerous regional interactions involve species that never co-occur, and as a result,
399 these interactions are not captured in local webs.

401 **Box 1: A spatiotemporally explicit model of interactions**

402 Predicting local webs across time and space is a pivotal goal of network ecology, especially given the scarcity of
 403 interaction data (Strydom *et al.* (2021)). Ecologists may resort to predictive models (e.g., generative Bayesian
 404 and machine learning models) to reconstruct networks at fine spatial and temporal scales with limited
 405 interaction data. For instance, real-time biomonitoring data coupled with appropriate numerical models (Bohan
 406 *et al.* (2017)) can be employed to reconstruct local ecological webs, opening avenues for in-depth studies on
 407 local ecosystem functioning and dynamics. The probabilistic representation of interactions acknowledges the
 408 inherent uncertainty in these models, typically expressed through probability distributions. We introduce and
 409 develop a simple generative mechanistic model for probabilistic local interactions that takes into consideration
 410 their spatiotemporal variability (i.e. a spatiotemporally explicit model of local interactions). It is essential to
 411 note that our model is not designed for regional interactions, which are scale-independent. Rather, it could
 412 prove valuable for predicting local interactions across time and space by generating new interaction data
 413 following parameter inference.

414 As indicated by eq. 1, the probability that two taxa i and j interact at a specific location (x, y, z) may be
 415 determined by the product of their probability of interaction given co-occurrence and their probability of
 416 co-occurrence. First, their co-occurrence probability can be calculated using their individual occurrence
 417 probabilities $P(X_i)$ and $P(X_j)$, along with the strength of association γ between their occurrences and
 418 co-occurrence (Cazelles *et al.* (2016)):

$$P(C_{i,j}) = P(X_i, X_j) = P(X_i)P(X_j)\gamma. \quad (7)$$

419 When $\gamma > 1$, it signifies a positive association in the geographic distributions of both taxa, indicating that the
 420 presence of one taxon enhances the probability of occurrence of the other. Attractions may be the result of
 421 positive interactions (e.g., mutualism) or positive/negative interactions (e.g., antagonism between predators and
 422 prey, Cazelles *et al.* (2016)). In empirical webs, $\gamma > 1$ holds true for the majority of species pairs (Catchen *et*
 423 *al.* (2023)). In contrast, repulsions ($\gamma < 1$) may be caused by strong interspecific competition (Cazelles *et al.*
 424 (2016)). We model the co-occurrence of both taxa as the outcome of a Bernoulli trial

$$C_{i,j} \sim \text{Bernoulli}(P(X_i, X_j)). \quad (8)$$

425 Next, the probability of interaction given co-occurrence can be made temporally explicit by modeling it as a
 426 Poisson process with rate parameter λ . This parameter represents the expected frequency of interaction between
 427 the taxa within a defined time interval and can be estimated using prior data on interaction strengths, when
 428 accessible. The probability that two co-occurring taxa engage in an interaction during a time period t_0 is given
 429 by:

$$P(L_{i \rightarrow j} | C_{i,j} = 1) = 1 - e^{-\lambda t_0}, \quad (9)$$

430 which tends toward 1 as $t_0 \rightarrow \infty$. It is important to note that the units of λ and t_0 are complementary. For
 431 instance, if the duration t_0 is measured in months, λ denote the expected number of interactions per month.
 432 The occurrence of an interaction between i and j can be modeled as a Bernoulli trial with a probability of
 433 $P(L_{i \rightarrow j})$. Consequently, a Bayesian inference model can be built based on the preceding equations to estimate
 434 the value of the λ and γ parameters and generate novel interaction data:

$$L_{i \rightarrow j} \sim \text{Bernoulli}(P(L_{i \rightarrow j})) \quad (10)$$

$$P(L_{i \rightarrow j}) = P(X_i)P(X_j)\gamma(1 - e^{-\lambda t_0}) \quad (11)$$

$$\gamma \sim \text{Gamma}(2, 0.5) \quad (12)$$

$$\lambda \sim \text{Exponential}(2) \quad (13)$$

435 In fig. 3, we show the variation in the probability of interaction under different parameter values. In the right
 436 panel, we notice that, irrespective of the interaction rate λ , the probability of interaction converges toward an
 437 asymptote determined by the co-occurrence $P(C_{i,j})$ (eq. 7). This model can be customized in different ways,

438 such as by linking λ with specific environmental variables or explicitly incorporating observation errors (i.e.,
439 the probabilities of false negatives and false positives).

440 [Figure 3 about here.]

441 **Taxonomic scaling of interactions**

442 Probabilistic webs offer a versatile approach to tackle a broad array of ecological questions, depending on their
443 level of organization. For instance, the assemblage of interactions across ecological scales can be explored
444 through species-based networks, while clade-based networks provide insights into macroevolutionary processes
445 (e.g., Gomez *et al.* (2010)). Given that our interpretation of the properties and dynamics of ecological webs
446 depends on their taxonomic scale (Guimarães (2020)), investigating the phylogenetic scaling of network
447 structure emerges as a promising research avenue. Examining the same system at various taxonomic scales can
448 yield meaningful and complementary ecological information, and, in our perspective, employing probabilistic
449 webs is an effective approach for such analyses.

450 There are no inherent differences in the taxonomic scaling between local and metawebs, as only the nodes are
451 defined taxonomically. In other words, the probability values of edges in both local (eq. 2) and metawebs (eq. 3)
452 are not conditional on any taxonomic scale. The phylogenetic scale is tied to the definition of the event itself
453 (i.e., the interaction between two taxa), not to the conditional variables. In both types of webs, transitioning to a
454 broader level of organization (e.g., from a species-level web S to a genus-level web G) can be accomplished
455 directly by using probabilities from finer scales. For example, in a network with n_1 species from genus g_1 and
456 n_2 species from genus g_2 , one can compute the probability that at least one species from genus g_1 interacts with
457 at least one species from genus g_2 as follows:

$$P(G_{g_1 \rightarrow g_2}) = 1 - \prod_{i=1}^{n_1} \prod_{j=1}^{n_2} (1 - P(S_{g_{1i} \rightarrow g_{2j}})), \quad (14)$$

458 where g_{1i} and g_{2j} are the species of the corresponding genus. If it is known that at least two of these species
459 interact (i.e., $P(S_{g_{1i} \rightarrow g_{2j}}) = 1$ for at least one pair of (g_{1i}, g_{2j})), it implies a probability of genus interaction
460 equal to 1. Canard *et al.* (2012) built a species-based network following a similar approach, by using simulated
461 interactions between individuals derived from a neutral model (i.e., a model that assumed ecological
462 equivalence among individuals). In contrast, a more sophisticated approach is necessary when transitioning

463 from a broader to a finer level of organization. This is because knowledge of an interaction between two genera
464 does not guarantee that all possible pairwise combinations of their species will also interact. One possible
465 method is to build a finer-scale network by generating probabilities of interactions through random sampling
466 from a beta distribution, parameterized by the broader-scale network.

467 Ideally, our biological interpretation of probabilistic interactions should remain consistent across a network
468 even if it incorporates heterogeneous levels of organization, such as a network whose nodes represent both
469 species and trophic species (groups of species sharing similar predators and prey). This is common in
470 ecological webs where taxonomic resolution is typically low (Hemprich-Bennett *et al.* (2021), Vázquez§ *et al.*
471 (2022)). Interaction probabilities at broader taxonomic scales should be based on probabilities of interactions
472 between individuals, for both local and metawebs. For instance, in local individual-based food webs, the
473 probability that two individuals interact reflects our degree of belief that one individual will consume the other.
474 Similarly, in local species-based food webs, the probability that two species interact represents our degree of
475 belief that at least one individual from the predator species consumes at least another individual from the prey
476 species. Furthermore, in local clade-based food webs, the probability that two clades interact represents our
477 degree of belief that at least two species from these clades interact with each other or, equivalently, that at least
478 two individuals from these clades interact with each other. Fundamentally, the taxonomic scaling of interactions
479 involves aggregating interactions between individuals into larger groups, which may exhibit varying degrees of
480 homogeneity based on the organisms and the study system. In that regard, taxonomic scaling is analogous to the
481 spatial and temporal scaling of interactions, as they all represent different ways to aggregate individuals into
482 broader groups (either spatially, temporally, or taxonomically).

483 Sampling for binary webs

484 The prediction of binary interactions through Bernoulli trials is an important application of probabilistic webs.
485 This approach proves beneficial for analyzing the structural characteristics of probabilistic webs, particularly in
486 the absence of specific analytical measures. By performing independent Bernoulli trials for each interaction in a
487 probabilistic web, a binary network may be generated. A probability distribution of network properties can be
488 obtained by measuring network structure across multiple randomly generated networks (Poisot *et al.* (2016)).
489 This method enables the representation of the variability of network structure, albeit with possible biases when
490 connectance is low (Poisot & Gravel (2014), Chagnon (2015)). Employing this strategy to generate binary
491 networks under a null model facilitates null hypothesis significance testing, wherein the observed measure is

492 compared against the simulated distribution (e.g., Bascompte *et al.* (2003)). Additionally, the random
493 generation of binary networks, from a probabilistic web that accounts for the spatiotemporal variability of
494 interactions, may effectively capture network structure across space and time. This facilitates the investigation
495 of ecological hypotheses about interactions at broad spatial and temporal scales.

496 There are at least two distinct approaches to sample binary networks from probabilistic webs across space, for
497 example, when attempting to predict a binary network for each of a number of locations within a given region.
498 The first approach involves performing a singular Bernoulli trial for each pair of taxa based on their regional
499 probability of interaction. In employing this approach, every pair of taxa predicted to interact in the binary
500 metaweb will be treated as interacting in all local webs where they co-occur. This will result in local pairwise
501 interactions without spatial variation. The second approach is to independently sample each of the local
502 probabilistic webs, which can be achieved by first generating distinct probabilistic networks for each location.
503 These local probabilistic webs may vary in taxa composition and interaction probabilities. Subsequently, binary
504 networks can be independently sampled for each location, introducing spatial variation in binary interactions.

505 In fig. 4, we compare the average connectance of binary networks resulting from these two sampling techniques,
506 where regional and local interactions are drawn from our host-parasite probabilistic networks, generating a
507 number of binary web realizations for each site in the dataset. These two sampling approaches yield different
508 outcomes, particularly for lower values of $P(L|M)$, which denote instances where regional interactions do not
509 consistently manifest locally. Small discrepancies are also apparent between these techniques when we equate
510 the probability of local interaction to the probability of regional interaction (i.e., when using $P(L|M) = 1.0$ in
511 eq. 5), especially when the number of binary network samples for each location is low. As anticipated, we
512 observe that sampling binary interactions from the metaweb tends to overestimate connectance on average
513 compared to sampling them from local webs. Furthermore, we observe an increase in the variability of
514 connectance when employing a single sample, representing what we consider as a more tangible process
515 leading to the realization of local and regional interactions in nature.

516 [Figure 4 about here.]

517 The choice of a sampling approach can influence the selection of grid cell size when delineating local
518 communities within a broader region of interest. In the first approach, pairwise interactions remain constant
519 irrespective of cell size since they are sampled only once from the metaweb. However, in the second approach,
520 local interaction probabilities are contingent on the network area. For instance, consider the local webs L_1 and

521 L_2 with an area of $\frac{1}{2}A_0$, both nested within A_0 but disjoint from each other, forming L_0 . If we treat L_1 and L_2 as
522 independent, the probability of interaction between taxa i and j in L_0 is given by:

$$P(L_{0,i \rightarrow j}) = 1 - (1 - P(L_{1,i \rightarrow j}) \times P(L_{2,i \rightarrow j})). \quad (15)$$

523 Due to its larger area, the probability that the two taxa interact in L_0 is equal or greater than in L_1 and L_2 . When
524 sampling binary interactions from local webs, it is crucial to sample at the same spatial scale for which
525 probabilities were estimated. Otherwise, interaction probabilities must be adjusted to align with the intended
526 cell size, preventing systematic biases in predictions.

527 Prediction of local webs from metawebs

528 Metawebs serve as a valuable source of ecological information for predicting local webs across time and space.
529 Binary local webs can be reconstructed by selecting a subset of taxa and interactions from the metaweb (Dunne
530 (2006)). This implies that metawebs consistently contain more interactions than their corresponding local webs.
531 Determining the list of taxa to select can be achieved empirically or through methods like range maps or species
532 distribution models. As the species composition of a community is arguably less difficult to sample or predict
533 than its interactions, the primary challenge lies in deciding which interactions to select from the metaweb, a task
534 that may necessitate advanced statistical models and ecological expertise. Inferring the structure of local webs
535 from the metaweb before predicting specific local pairwise interactions could hold promise (Strydom *et al.*
536 (2021)), considering that the structure of local webs is constrained by the metaweb (Saravia *et al.* (2022)).

537 Building probabilistic local webs from a probabilistic metaweb involves a reduction in the value of pairwise
538 interaction probabilities. This decrease is attributed to the prerequisite that two taxa must initially possess the
539 capacity to interact before engaging in local interactions (eq. 5). Therefore, inferring local webs from their
540 metaweb while maintaining identical interaction probability values would introduce systematic biases into the
541 predictions. In such cases, these networks would essentially represent smaller-scale metawebs of potential
542 interactions, possibly leading to misinterpretations by being perceived as local interactions. As proposed by
543 McLeod *et al.* (2021), although metawebs do not capture the spatiotemporal variability of interactions, they
544 establish an upper limit for local interactions. In other words, the probability of two taxa interacting at a specific
545 location and time is consistently lower or equal to the probability of their regional interaction:

$$P(L_{i \rightarrow j}|A, t, C, \Omega) \leq P(M_{i \rightarrow j}). \quad (16)$$

546 Moreover, the probability that two taxa possess the biological capacity to interact must exceed the probability of
 547 them interacting at any location and time because they may never co-occur or encounter locally. Specifically,
 548 the cumulative probability of local interactions across all spatial, temporal, and environmental conditions must
 549 be less than the probability of regional interaction, i.e.

$$\int_{\Omega} \int_A \int_t P(L_{i \rightarrow j}|A, t, \Omega) dt dA d\Omega \leq P(M_{i \rightarrow j}). \quad (17)$$

550 Estimating more precisely the probability $P(L|M)$ that two taxa interact locally if they can potentially interact
 551 allows for improved predictions of local webs from a probabilistic metaweb. This task is challenging due to the
 552 variability of this probability across space and time, as well as its variability across pairwise interactions within
 553 a network. Using simple models of $P(L|M)$, as demonstrated in our case studies, represents an initial step
 554 toward the overarching objective of reconstructing local webs from metawebs.

555 Conclusion

556 In this contribution, we underline the importance of network metadata for adequately interpreting and
 557 manipulating probabilistic interaction data. The mathematical representation of probabilities and their statistical
 558 properties depend on the type of interactions (local or regional) and the conditions under which these
 559 interactions were evaluated. We showed that probabilistic local and metawebs differ in their relationship to
 560 spatial and temporal scales, with regional interactions remaining consistent across scales. In contrast with
 561 metawebs, local interactions are measured in a specific context (e.g., in a given area, time, and biological and
 562 environmental conditions) and depend on taxa co-occurrence. These important conceptual differences bring to
 563 light the need to use probabilistic data with caution, for instance when generating binary network realizations
 564 across space and predicting local webs from metawebs. Clear metadata describing the type of interaction and
 565 the variables used in their estimation are required to ensure adequate data manipulation. Better data practices
 566 and foundations for probabilistic thinking in network ecology could enable more reliable assessments of the
 567 spatiotemporal variability and uncertainty of biotic interactions.
 568 It is essential to enhance our comprehension of both regional and local interactions, especially considering the

569 current scarcity of interaction data. However, while sampling biological communities does decrease the
570 uncertainty of interactions by accumulating evidence for their feasibility and local realization, there is a limit to
571 how much we can diminish their inherent randomness. In metawebs, probabilities reflect our limited knowledge
572 of interactions (i.e., our degree of belief that interactions are feasible), which is expected to improve with a
573 larger volume of data. We should anticipate that regional interactions will become more definitive (with
574 probabilities approaching 0 or 1) as we investigate various conditions, including different combinations of
575 species traits. Conversely, in the case of local webs, which can be seen as random instances of metawebs,
576 randomness cannot be as reduced. In local interactions, probabilities can be divided into a component
577 representing uncertainty and another representing spatiotemporal variability. Owing to environmental
578 heterogeneity, there will invariably be instances in which an interaction occurs and others in which it does not,
579 across different times and locations, irrespective of the extent to which we can improve our knowledge of its
580 biological feasibility and the local conditions that facilitate its occurrence. When local webs depict probabilities
581 of observing interactions rather than just their actual occurrence, we must also consider the observation
582 variability as an additional source of randomness. Every ecological process is stochastic but there is also a
583 possibility that a phenomenon goes undetected. Quantifying and partitioning this stochasticity will enable us to
584 make more accurate predictions about ecological interactions at various spatial and temporal scales. This will
585 prove to be of vital importance as our time to understand nature runs out, especially at the places where the
586 impacts of climate change and habitat loss hit harder.

587 Acknowledgment

588 We acknowledge that this study was conducted on land within the traditional unceded territory of the Saint
589 Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and Omàmiwininiwak nations. This work was
590 supported by the Institute for Data Valorisation (IVADO) and the Natural Sciences and Engineering Research
591 Council of Canada (NSERC) Collaborative Research and Training Experience (CREATE) program, through the
592 Computational Biodiversity Science and Services (BIOS²) program. A special thanks to all members of the
593 Black Holes and Revelations working group (organized by BIOS²) for their insightful discussions and valuable
594 feedback on this manuscript.

595 **References**

- 596 Abrego, N., Roslin, T., Huotari, T., Ji, Y., Schmidt, N.M., Wang, J., *et al.* (2021). Accounting for species
597 interactions is necessary for predicting how arctic arthropod communities respond to climate change.
598 *Ecography*, 44, 885–896.
- 599 Andrade-Pacheco, R., Rerolle, F., Lemoine, J., Hernandez, L., Meité, A., Juziwelo, L., *et al.* (2020). Finding
600 hotspots: Development of an adaptive spatial sampling approach. *Scientific Reports*, 10, 10939.
- 601 Angilletta, M.J., Jr., Steury, T.D. & Sears, M.W. (2004). Temperature, Growth Rate, and Body Size in
602 Ectotherms: Fitting Pieces of a Life-History Puzzle¹. *Integrative and Comparative Biology*, 44, 498–509.
- 603 Araujo, M.B., Rozenfeld, A., Rahbek, C. & Marquet, P.A. (2011). Using species co-occurrence networks to
604 assess the impacts of climate change. *Ecography*, 34, 897–908.
- 605 Banašek-Richter, C., Cattin, M.-F. & Bersier, L.-F. (2004). Sampling effects and the robustness of quantitative
606 and qualitative food-web descriptors. *Journal of Theoretical Biology*, 226, 23–32.
- 607 Bartomeus, I., Gravel, D., Tylianakis, J.M., Aizen, M.A., Dickie, I.A. & Bernard-Verdier, M. (2016). A
608 common framework for identifying linkage rules across different types of interactions. *Functional Ecology*,
609 30, 1894–1903.
- 610 Bascompte, J., Jordano, P., Melián, C.J. & Olesen, J.M. (2003). The nested assembly of plantanimal mutualistic
611 networks. *Proceedings of the National Academy of Sciences*, 100, 9383–9387.
- 612 Berlow, E.L., Neutel, A.-M., Cohen, J.E., De Ruiter, P.C., Ebenman, B., Emmerson, M., *et al.* (2004).
613 Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73, 585–598.
- 614 Bersier, L.-F., Dixon, P. & Sugihara, G. (1999). Scale-Invariant or Scale-Dependent Behavior of the Link
615 Density Property in Food Webs: A Matter of Sampling Effort? *The American Naturalist*, 153, 676–682.
- 616 Blanchet, F.G., Cazelles, K. & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions.
617 *Ecology Letters*, 23, 1050–1063.
- 618 Bohan, D.A., Vacher, C., Tamaddoni-Nezhad, A., Raybould, A., Dumbrell, A.J. & Woodward, G. (2017).
619 Next-Generation Global Biomonitoring: Large-scale, Automated Reconstruction of Ecological Networks.
620 *Trends in Ecology & Evolution*, 32, 477–487.

- 621 Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011). Why
622 **intraspecific trait variation matters in community ecology.** *Trends in Ecology & Evolution*, 26, 183–192.
- 623 Borrett, S.R. & Scharler, U.M. (2019). **Walk partitions of flow in Ecological Network Analysis: Review and**
624 **synthesis of methods and indicators.** *Ecological Indicators*, 106, 105451.
- 625 Boulangeat, I., Gravel, D. & Thuiller, W. (2012). **Accounting for dispersal and biotic interactions to disentangle**
626 **the drivers of species distributions and their abundances.** *Ecology Letters*, 15, 584–593.
- 627 Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T. & Fortin, M.-J. (2023). **Shortcomings of reusing**
628 **species interaction networks created by different sets of researchers.** *PLOS Biology*, 21, e3002068.
- 629 Broom, M. & Ruxton, G.D. (2005). **You can run or you can hide: Optimal strategies for cryptic prey against**
630 **pursuit predators.** *Behavioral Ecology*, 16, 534–540.
- 631 Brose, U., Ostling, A., Harrison, K. & Martinez, N.D. (2004). **Unified spatial scaling of species and their**
632 **trophic interactions.** *Nature*, 428, 167–171.
- 633 Canard, E., Mouquet, N., Marescot, L., Gaston, K.J., Gravel, D. & Mouillot, D. (2012). **Emergence of**
634 **Structural Patterns in Neutral Trophic Networks.** *PLOS ONE*, 7, e38295.
- 635 Catchen, M.D., Poisot, T., Pollock, L.J. & Gonzalez, A. (2023). **The missing link: Discerning true from false**
636 **negatives when sampling species interaction networks.**
- 637 Cazelles, K., Araujo, M.B., Mouquet, N. & Gravel, D. (2016). **A theory for species co-occurrence in interaction**
638 **networks.** *Theoretical Ecology*, 9, 39–48.
- 639 Chagnon, P.-L. (2015). **Characterizing topology of ecological networks along gradients: The limits of metrics'**
640 **standardization.** *Ecological Complexity*, 22, 36–39.
- 641 Choh, Y., Ignacio, M., Sabelis, M.W. & Janssen, A. (2012). **Predator-prey role reversals, juvenile experience**
642 **and adult antipredator behaviour.** *Scientific Reports*, 2, 728.
- 643 Cirtwill, A.R., Eklöf, A., Roslin, T., Wootton, K. & Gravel, D. (2019). **A quantitative framework for**
644 **investigating the reliability of empirical network construction.** *Methods in Ecology and Evolution*, 10,
645 902–911.
- 646 Dallas, T., Park, A.W. & Drake, J.M. (2017). **Predicting cryptic links in host-parasite networks.** *PLOS*
647 *Computational Biology*, 13, e1005557.

- 648 Dunne, J.A. (2006). The Network Structure of Food Webs. In: *Ecological networks: Linking structure and*
649 *dynamics* (eds. Dunne, J.A. & Pascual, M.). Oxford University Press, pp. 27–86.
- 650 Elmasri, M., Farrell, M.J., Davies, T.J. & Stephens, D.A. (2020). [A hierarchical Bayesian model for predicting](#)
651 [ecological interactions using scaled evolutionary relationships](#). *The Annals of Applied Statistics*, 14,
652 221–240.
- 653 Elton, C.S. (2001). *Animal Ecology*. University of Chicago Press, Chicago, IL.
- 654 Emmerson, M.C. & Raffaelli, D. (2004). [Predator-prey body size, interaction strength and the stability of a real](#)
655 [food web](#). *Journal of Animal Ecology*, 73, 399–409.
- 656 Fortuna, M.A. & Bascompte, J. (2006). [Habitat loss and the structure of plant-animal mutualistic networks](#).
657 *Ecology Letters*, 9, 281–286.
- 658 Fu, X., Seo, E., Clarke, J. & Hutchinson, R.A. (2021). [Link Prediction Under Imperfect Detection: Collaborative Filtering for Ecological Networks](#). *IEEE Transactions on Knowledge and Data Engineering*,
659 33, 3117–3128.
- 660 Galiana, N., Lurgi, M., Claramunt-López, B., Fortin, M.-J., Leroux, S., Cazelles, K., et al. (2018). [The spatial scaling of species interaction networks](#). *Nature Ecology & Evolution*, 2, 782–790.
- 661 Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010). [A framework for community interactions under climate change](#). *Trends in Ecology & Evolution*, 25, 325–331.
- 662 Godsoe, W., Murray, R. & Iritani, R. (2022). [Species interactions and diversity: A unified framework using Hill numbers](#). *Oikos*, n/a, e09282.
- 663 Golubski, A.J. & Abrams, P.A. (2011). [Modifying modifiers: What happens when interspecific interactions interact?](#) *Journal of Animal Ecology*, 80, 1097–1108.
- 664 Gomez, J.M., Verdu, M. & Perfectti, F. (2010). [Ecological interactions are evolutionarily conserved across the entire tree of life](#). *Nature*, 465, 918–U6.
- 665 Gonzalez, A. & Londoño, M.C. (2022). [Monitor biodiversity for action](#). *Science*, 378, 1147–1147.
- 666 Gonzalez-Varo, J.P. & Traveset, A. (2016). [The Labile Limits of Forbidden Interactions](#). *Trends in Ecology & Evolution*, 31, 700–710.

- 674 Gravel, D., Baiser, B., Dunne, J.A., Kopalke, J.-P., Martinez, N.D., Nyman, T., *et al.* (2019). Bringing Elton
675 and Grinnell together: A quantitative framework to represent the biogeography of ecological interaction
676 networks. *Ecography*, 42, 401–415.
- 677 Gravel, D., Poisot, T., Albouy, C., Velez, L. & Mouillot, D. (2013). Inferring food web structure from
678 predator-prey body size relationships. *Methods in Ecology and Evolution*, 4, 1083–1090.
- 679 Guimarães, P.R. (2020). The Structure of Ecological Networks Across Levels of Organization. *Annual Review*
680 of Ecology, Evolution, and Systematics, 51, 433–460.
- 681 Guimerà, R. & Sales-Pardo, M. (2009). Missing and spurious interactions and the reconstruction of complex
682 networks. *Proceedings of the National Academy of Sciences*, 106, 22073–22078.
- 683 Hemprich-Bennett, D.R., Oliveira, H.F.M., Le Comber, S.C., Rossiter, S.J. & Clare, E.L. (2021). Assessing the
684 impact of taxon resolution on network structure. *Ecology*, 102, e03256.
- 685 Herrera, C.M. (1989). Pollinator abundance, morphology, and flower visitation rate: Analysis of the “quantity”
686 component in a plant-pollinator system. *Oecologia*, 80, 241–248.
- 687 Hortal, J., de Bello, F., Diniz-Filho, J.A.F., Lewinsohn, T.M., Lobo, J.M. & Ladle, R.J. (2015). Seven Shortfalls
688 that Beset Large-Scale Knowledge of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*,
689 46, 523–549.
- 690 Jordano, P. (1987). Patterns of Mutualistic Interactions in Pollination and Seed Dispersal: Connectance,
691 Dependence Asymmetries, and Coevolution. *The American Naturalist*, 129, 657–677.
- 692 Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*, 30, 1883–1893.
- 693 Jordano, P., Bascompte, J. & Olesen, J.M. (2003). Invariant properties in coevolutionary networks of
694 plant-animal interactions. *Ecology Letters*, 6, 69–81.
- 695 Kéfi, S., Berlow, E.L., Wieters, E.A., Joppa, L.N., Wood, S.A., Brose, U., *et al.* (2015). Network structure
696 beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky shores. *Ecology*, 96,
697 291–303.
- 698 Kéfi, S., Berlow, E.L., Wieters, E.A., Navarrete, S.A., Petchey, O.L., Wood, S.A., *et al.* (2012). More than a
699 meal... integrating non-feeding interactions into food webs. *Ecology Letters*, 15, 291–300.
- 700 Kéfi, S., Miele, V., Wieters, E.A., Navarrete, S.A. & Berlow, E.L. (2016). How Structured Is the Entangled
701 Bank? The Surprisingly Simple Organization of Multiplex Ecological Networks Leads to Increased

- 702 Persistence and Resilience. *PLOS Biology*, 14, e1002527.
- 703 Klecka, J. & Boukal, D.S. (2014). The effect of habitat structure on prey mortality depends on predator and
704 prey microhabitat use. *Oecologia*, 176, 183–191.
- 705 Kopelke, J.-P., Nyman, T., Cazelles, K., Gravel, D., Vissault, S. & Roslin, T. (2017). Food-web structure of
706 willow-galling sawflies and their natural enemies across Europe. *Ecology*, 98, 1730–1730.
- 707 Kulkarni, M.A., Kweka, E., Nyale, E., Lyatuu, E., Mosha, F.W., Chandramohan, D., et al. (2006).
708 Entomological Evaluation of Malaria Vectors at Different Altitudes in Hai District, Northeastern Tanzania.
709 *Journal of Medical Entomology*, 43, 580–588.
- 710 McCann, K. (2007). Protecting biostructure. *Nature*, 446, 29–29.
- 711 McCann, K.S. (2011). Food webs (MPB-50). In: *Food Webs (MPB-50)*. Princeton University Press.
- 712 McLeod, A., Leroux, S.J., Gravel, D., Chu, C., Cirtwill, A.R., Fortin, M.-J., et al. (2021). Sampling and
713 asymptotic network properties of spatial multi-trophic networks. *Oikos*, n/a.
- 714 McLeod, A.M., Leroux, S.J. & Chu, C. (2020). Effects of species traits, motif profiles, and environment on
715 spatial variation in multi-trophic antagonistic networks. *Ecosphere*, 11, e03018.
- 716 Melián, C.J., Vilas, C., Baldó, F., González-Ortegón, E., Drake, P. & Williams, R.J. (2011). Eco-evolutionary
717 Dynamics of Individual-Based Food Webs. In: *Advances in Ecological Research*, The Role of Body Size in
718 Multispecies Systems (ed. Belgrano, A.). Academic Press, pp. 225–268.
- 719 Ohlmann, M., Miele, V., Dray, S., Chalmandrier, L., O'Connor, L. & Thuiller, W. (2019). Diversity indices for
720 ecological networks: A unifying framework using Hill numbers. *Ecology Letters*, 22, 737–747.
- 721 Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H., Rasmussen, C. & Jordano, P. (2010). Missing and
722 forbidden links in mutualistic networks. *Proceedings of the Royal Society B: Biological Sciences*, 278,
723 725–732.
- 724 Pascual, M., Dunne, J.A. & Dunne, J.A. (2006). *Ecological Networks: Linking Structure to Dynamics in Food
725 Webs*. Oxford University Press, USA.
- 726 Petchey, O.L., Beckerman, A.P., Riede, J.O. & Warren, P.H. (2008). Size, foraging, and food web structure.
727 *Proceedings of the National Academy of Sciences*, 105, 4191–4196.

- 728 Pilosof, S., Porter, M.A., Pascual, M. & Kéfi, S. (2017). [The multilayer nature of ecological networks](#). *Nature Ecology & Evolution*, 1, 1–9.
- 729
- 730 Plagányi, É.E. & Butterworth, D.S. (2004). [A critical look at the potential of Ecopath with ecosim to assist in practical fisheries management](#). *African Journal of Marine Science*, 26, 261–287.
- 731
- 732 Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). [The dissimilarity of species interaction networks](#). *Ecology Letters*, 15, 1353–1361.
- 733
- 734 Poisot, T., Cirtwill, A.R., Cazelles, K., Gravel, D., Fortin, M.-J. & Stouffer, D.B. (2016). [The structure of probabilistic networks](#). *Methods in Ecology and Evolution*, 7, 303–312.
- 735
- 736 Poisot, T. & Gravel, D. (2014). [When is an ecological network complex? Connectance drives degree distribution and emerging network properties](#). *PeerJ*, 2, e251.
- 737
- 738 Poisot, T., Stouffer, D.B. & Gravel, D. (2015). [Beyond species: Why ecological interaction networks vary through space and time](#). *Oikos*, 124, 243–251.
- 739
- 740 Polis, G.A. (1991). [Complex Trophic Interactions in Deserts: An Empirical Critique of Food-Web Theory](#). *The American Naturalist*, 138, 123–155.
- 741
- 742 Pollock, L.J., Tingley, R., Morris, W.K., Golding, N., O’Hara, R.B., Parris, K.M., *et al.* (2014). [Understanding co-occurrence by modelling species simultaneously with a Joint Species DistributionModel \(JSDM\)](#).
- 743
- 744 *Methods in Ecology and Evolution*, 5, 397–406.
- 745
- 746 Pomeranz, J.P.F., Thompson, R.M., Poisot, T. & Harding, J.S. (2019). [Inferring predatorprey interactions in food webs](#). *Methods in Ecology and Evolution*, 10, 356–367.
- 747
- 748 Pringle, R.M. & Hutchinson, M.C. (2020). [Resolving Food-Web Structure](#). *Annual Review of Ecology, Evolution, and Systematics*, 51, 55–80.
- 749
- 750 Proulx, S.R., Promislow, D.E.L. & Phillips, P.C. (2005). [Network thinking in ecology and evolution](#). *Trends in Ecology & Evolution*, SPECIAL ISSUE: BUMPER BOOK REVIEW, 20, 345–353.
- 751
- Pulliam, H.R. (1974). [On the Theory of Optimal Diets](#). *The American Naturalist*, 108, 59–74.
- 752
- 753 Rooney, N. & McCann, K.S. (2012). [Integrating food web diversity, structure and stability](#). *Trends in Ecology & Evolution*, 27, 40–46.

- 754 Salim, J.A., Saraiva, A.M., Zermoglio, P.F., Agostini, K., Wolowski, M., Drucker, D.P., *et al.* (2022). Data
755 standardization of plantpollinator interactions. *GigaScience*, 11, giac043.
- 756 Sanders, D. & van Veen, F.J.F. (2012). Indirect commensalism promotes persistence of secondary consumer
757 species. *Biology Letters*, 8, 960–963.
- 758 Saravia, L.A., Marina, T.I., Kristensen, N.P., De Troch, M. & Momo, F.R. (2022). Ecological network
759 assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, 91, 630–642.
- 760 Schmeller, D.S., Julliard, R., Bellingham, P.J., Böhm, M., Brummitt, N., Chiarucci, A., *et al.* (2015). Towards a
761 global terrestrial species monitoring program. *Journal for Nature Conservation*, 25, 51–57.
- 762 Shaw, J.O., Dunhill, A.M., Beckerman, A.P., Dunne, J.A. & Hull, P.M. (2024). A framework for reconstructing
763 ancient food webs using functional trait data.
- 764 Singer, M.C. & McBride, C.S. (2012). Geographic mosaics of species' association: A definition and an
765 example driven by plantinsect phenological synchrony. *Ecology*, 93, 2658–2673.
- 766 Stouffer, D.B., Rezende, E.L. & Amaral, L.A.N. (2011). The role of body mass in diet contiguity and food-web
767 structure. *Journal of Animal Ecology*, 80, 632–639.
- 768 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M.J., *et al.* (2022). Food web
769 reconstruction through phylogenetic transfer of low-rank network representation. *Methods in Ecology and
770 Evolution*, 13.
- 771 Strydom, T., Catchen, M.D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., *et al.* (2021). A
772 roadmap towards predicting species interaction networks (across space and time). *Philosophical
773 Transactions of the Royal Society B-Biological Sciences*, 376, 20210063.
- 774 Tylianakis, J.M., Tscharntke, T. & Lewis, O.T. (2007). Habitat modification alters the structure of tropical
775 hostparasitoid food webs. *Nature*, 445, 202–205.
- 776 Vázquez§, D.P., Peralta§, G., Cagnolo, L., Santos, M. & Igual, §.E. autores contribuyeron por. (2022).
777 Ecological interaction networks. What we know, what we don't, and why it matters. *Ecología Austral*, 32,
778 670–697.
- 779 Weinstein, B.G. & Graham, C.H. (2017a). On comparing traits and abundance for predicting species
780 interactions with imperfect detection. *Food Webs*, 11, 17–25.

- 781 Weinstein, B.G. & Graham, C.H. (2017b). Persistent bill and corolla matching despite shifting temporal
782 resources in tropical hummingbird-plant interactions. *Ecology Letters*, 20, 326–335.
- 783 Whittaker, R.H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological
784 Monographs*, 30, 279–338.
- 785 Wood, S.A., Russell, R., Hanson, D., Williams, R.J. & Dunne, J.A. (2015). Effects of spatial scale of sampling
786 on food web structure. *Ecology and Evolution*, 5, 3769–3782.
- 787 Woodward, G., Benstead, J.P., Beveridge, O.S., Blanchard, J., Brey, T., Brown, L.E., et al. (2010). Chapter 2 -
788 Ecological Networks in a Changing Climate. In: *Advances in Ecological Research*, Ecological Networks
789 (ed. Woodward, G.). Academic Press, pp. 71–138.
- 790 Woodward, G., Brown, L.E., Edwards, F.K., Hudson, L.N., Milner, A.M., Reuman, D.C., et al. (2012). Climate
791 change impacts in multispecies systems: Drought alters food web size structure in a field experiment.
792 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 2990–2997.




Figure 1: Network accumulation curves. (a) Dissimilarity of interactions between common species and (b) dissimilarity in species composition between aggregated local webs and the binary metaweb of host-parasite interactions. Aggregated local webs were obtained by sequentially and randomly selecting a number of binary local webs and aggregating both their species and interactions. In both panels, the colored line represents the median dissimilarity across simulations and the grey areas cover the 50% and 95% percentile intervals. (c) Scaling of the number of links and (d) scaling of connectance with the number of sampled binary and probabilistic local webs. For a better comparison with binary webs, local probabilistic webs were derived from the probabilistic metaweb with a false positive and false negative rate of zero. A specific value of $P(L|M)$ (the local probability of interaction among potentially interacting species) was used for all local webs within a particular curve. Aggregated probabilistic local webs were obtained by sequentially and randomly selecting a number of probabilistic local webs and aggregating both their species and interactions (with the value of $P(L|M)$ adjusting according to eq. 6).




Figure 2: Spatial scaling of interactions. Expected number of host-parasite interactions in a network aggregating all probabilistic (a) local and (b) regional interactions within a latitudinal window of a given length. Every dashed curve corresponds to a different expanding window characterized by a different central latitude, with the colored solid line representing the median number of interactions. Heatmaps of the expected number of (c) local and (d) regional interactions found in windows of specified length and central latitudes. Probabilities of regional interactions were obtained with a false positive rate of 5% and a false negative rate of 10%. Local probabilistic interactions were derived from the probabilistic metaweb by setting the value of $P(L|M)$ (the local probability of interaction among potentially interacting species) to 1, ensuring a conservative comparison between aggregated local webs and metawebs. Aggregated local webs were obtained by aggregating both the species and interactions found within a particular latitudinal window, with the values of $P(L|M)$ remaining at their maximum value of 1 following eq. 6.




Figure 3: Parameters of the spatiotemporally explicit model of interactions. (a) Probability of local interaction given by the process model (eq. 11) under different values of λ and γ , with $t_0 = 1$. The parameter values used in the right panel are denoted by the white stars. (b) Scaling of the probability of interaction with the duration parameter t_0 in eq. 11, for different values of λ and γ . In both panels, the individual probabilities of occurrence $P_i(x, y, z)$ and $P_j(x, y, z)$ are set to a constant value of 0.5.




Figure 4: Connectance of sampled binary webs. Comparison between the average connectance of binary network samples obtained from the probabilistic local and metawebs. Each dot corresponds to a different site. The local probability of interaction between potentially interacting species was set to three different values: (a) $P(L|M) = 1.0$, (b) $P(L|M) = 0.75$, and (c) $P(L|M) = 0.50$. Grey dots represent the outcome of a single trial, while colored dots represent the average connectance of each network across 100 trials. (d) Reduction in the mean squared logarithmic error between the average connectance of binary networks obtained from these two sampling methods as the number of trials increases, for the same values of $P(L|M)$ used in panels a-c. Probabilities of regional interactions were obtained with a false positive rate of 5% and a false negative rate of 10%. Metaweb samples were obtained by randomly sampling binary interactions from the probabilistic metaweb, and then propagating this result to all local webs that include the species potentially engaged in the interactions. Local binary webs were generated by independently sampling binary interactions for each local probabilistic web.