

# **What constrains food webs? A maximum entropy framework for predicting their structure with minimal biases**

Francis Banville<sup>1,2,3</sup> Dominique Gravel<sup>2,3</sup> Timothée Poisot<sup>1,3</sup>

<sup>1</sup> Université de Montréal <sup>2</sup> Université de Sherbrooke <sup>3</sup> Quebec Centre for Biodiversity Science

## **Correspondance to:**

Francis Banville — francis.banville@umontreal.ca

**Abstract:** Food webs are complex ecological networks whose structure is both ecologically and statistically constrained, with many network properties being correlated with each other. Despite the recognition of these invariable relationships in food webs, the use of the principle of maximum entropy (MaxEnt) in network ecology is still rare. This is surprising considering that MaxEnt is a renowned and rigorous statistical tool precisely designed for understanding and predicting many different types of constrained systems. Precisely, this principle asserts that the least-biased probability distribution of a system's property, constrained by prior knowledge about that system, is the one with maximum information entropy. MaxEnt has been proven useful in many ecological modelling problems, but its application in food webs and other ecological networks is limited. Here we show how MaxEnt can be used to derive many food-web properties both analytically and heuristically. First, we show how the joint degree distribution (the joint probability distribution of the numbers of prey and predators for each species in the network) can be derived analytically using the number of species and the number of interactions in food webs. Second, we present a heuristic and flexible approach of finding a network's adjacency matrix (the network's representation in matrix format) based on simulated annealing and SVD entropy. We built two heuristic models using the connectance and the joint degree sequence as statistical constraints, respectively. We compared both models' predictions against corresponding null and neutral models commonly used in network ecology using open access data of terrestrial and aquatic food webs sampled globally ( $N = 257$ ). We found that the heuristic model constrained by the joint degree sequence was a good predictor of many measures of food-web structure, especially the nestedness and motifs distribution. Specifically, our results suggest that the structure of terrestrial and aquatic food webs is mainly driven by their joint degree distribution.

## **1 Introduction**

### **2 The constrained structure of ecological networks**

3 A variety of measures of the structure of ecological networks have been used to describe the  
4 organization of species interactions in a biological community (Delmas et al. 2019). These  
5 measures provide valuable information on the functioning of ecosystems and their responses  
6 to environmental change (e.g., Pascual and Dunne 2006; Gómez, Perfectti, and Jordano 2011).  
7 For instance, Bascompte et al. (2003) suggest that plant–pollinator and seed-disperser networks  
8 have a highly nested structure that can promote species persistence. Another example, in food  
9 webs, shows that a high connectance can promote the robustness of the system to species lost  
10 (Dunne, Williams, and Martinez 2002). However, despite the growing literature on the ecologi-  
11 cal implications of network structure, the association between many of these measures impedes  
12 our ability to fully understand what drives the structure and behavior of ecological networks. In  
13 particular, nestedness and modularity are strongly associated in ecological networks (Fortuna et  
14 al. 2010), and network connectance has been shown to be an important driver of many other  
15 emerging network properties (Timothée Poisot and Gravel 2014). In light of these observations,  
16 it is difficult to assess whether attributed effects of given properties are the artifacts of other,  
17 perhaps simpler, measures.

18 One way to tackle this issue is first to recognize that food webs and other ecological networks  
19 are constrained systems. In other words, the space of possible network configurations shrinks  
20 as we know more about a network structure. For example, there is a finite number of networks  
21 with specified numbers of nodes and edges. Indeed, the structure of ecological networks is first  
22 and above all constrained by the number of species, or nodes, present. Food webs with high  
23 species richness typically have a lower connectance (MacDonald, Banville, and Poisot 2020)  
24 than smaller networks. This is because the number of realized interactions in empirical food  
25 webs scales slower than the number of possible species pairs (MacDonald, Banville, and Poisot  
26 2020). As shown by Timothée Poisot and Gravel (2014), connectance itself can constrain differ-  
27 ent aspects of network structure such as the degree distribution (i.e. the probability distribution  
28 of the number of interspecific interactions realized by a species). Other measures, such as the

29 maximum trophic level, can also constrain the space of feasible networks.

30 Prior knowledge on the structure of ecological networks is thus especially useful in the current  
31 context of data scarcity about species interactions (Jordano 2016). Indeed, this Eltonian shortfall  
32 (Hortal et al. 2015) can be partially alleviated using known information about an ecological  
33 network. As suggested by Strydom et al. (2021), network structure can be used to improve the  
34 prediction of pairwise species interactions when data is lacking by constraining the space of  
35 feasible networks. Similarly, partial knowledge on the structure of an ecological network can  
36 also be used to predict others of its properties by constraining their range of possible values.  
37 This is important given that many aspects of network structure cannot be measured empirically  
38 without data on pairwise species interactions, a prevailing situation worldwide (Timothée Poisot  
39 et al. 2021).

40 Understanding the ecological constraints that shape species interactions networks and predicting  
41 their emerging structure are thus two complementary aims of network ecology. This distinction  
42 between understanding and predicting is essential when using statistical and mathematical mod-  
43 els in network ecology and interpreting them. On one hand, null models help us identify potential  
44 ecological mechanisms that drive species interactions and constrain ecological networks. Null  
45 models generate a distribution for a target measure using a set of rules that exclude the mecha-  
46 nism of interest (Fortuna and Bascompte 2006; Delmas et al. 2019). The deviation between the  
47 model and empirical data helps us evaluate the effect of this ecological process in nature. On  
48 the other hand, predictive models can help fill many gaps on species interactions data. A variety  
49 of such models have recently been developed using machine learning and other statistical tools,  
50 most of which are presented in Strydom et al. (2021). However, given the constrained nature of  
51 ecological networks, it is surprising that the principle of maximum entropy, a rigorous mathe-  
52 matical method designed for both the analysis and prediction of constrained systems, has been  
53 barely used in network ecology.

#### 54 **The principle of maximum entropy: A primer for ecologists**

55 The principle of maximum entropy (MaxEnt) is a mathematical method of finding probabil-  
56 ity distributions, strongly rooted in statistical mechanics and information theory (Jaynes 1957a,

57 1957b; Harremoës and Topsøe 2001). Starting from a set of constraints given by prior knowl-  
58 edge of a system (i.e. what we call state variables), this method helps us find least-biased prob-  
59 ability distributions subject to the constraints. These probability distributions are guaranteed to  
60 be unique given our prior knowledge and represent the most we can say about a system without  
61 making more assumptions. For example, if the only thing we know about a biological commu-  
62 nity is its average number of individuals per species, the least-biased inference we could make on  
63 its species abundance distribution is the exponential distribution (Frank and Smith 2011; Harte  
64 and Newman 2014). However, this does not imply that this distribution will be the best fit to  
65 empirical data. The challenge is to find the right set of constraints that would best reproduce  
66 distributions found in nature.

67 MaxEnt states that the least-biased probability distribution given the constraints used is the one  
68 with the highest entropy among all probability distributions that satisfy these constraints. En-  
69 tropy is a measure of the average amount of information given by the outcome of a random  
70 variable. Many measures of entropy have been developed in physics (Beck 2009), but only  
71 a fraction of them could be used as an optimization measure with the principle of maximum  
72 entropy. According to Beck (2009) and Khinchin (2013), a measure of entropy  $H$  should sat-  
73 isfy four properties in the discrete case: (1) it should be a function of a probability distribution  
74  $p(n)$  only; (2) it should be maximized when  $p(n)$  is uniform; (3) it should not be influenced by  
75 outcomes with a null probability; and (4) it should be independent of the order of information  
76 acquisition. The Shannon's entropy (Shannon 1948)

$$H = - \sum_n p(n) \log p(n) \quad (1)$$

77 satisfies all of these properties. Finding the probability distribution  $p(n)$  that maximizes  $H$  under  
78 a set of  $m$  constraints  $g$  can be done using the method of Lagrange multipliers. These constraints  
79 could include one or many properties of the probability distribution (e.g., its mean, variance, and  
80 range). However, the normalization constraint always need to be included in  $g$  in order to make  
81 sure that  $p(n)$  sums to 1. The objective is then to find the values of the Lagrange multipliers  $\lambda_i$   
82 that optimize a function  $F$ :

$$F = H - \sum_{i=1}^m \lambda_i(g_i - c_i), \quad (2)$$

83 where  $g_i$  is the mathematical formulation of the constraint  $i$  and  $c_i$ , its value. Note that  $F$  is just  
 84 Shannon's entropy to which we added terms that each sums to zero ( $g_i = c_i$ ).  $F$  is maximized  
 85 by setting to 0 its partial derivative with respect to  $p(n)$ .

86 The principle of maximum entropy has been used in a wide range of disciplines, from thermo-  
 87 dynamics, chemistry and biology (Martyushev and Seleznev 2006) to graph and network theory  
 88 (e.g., Park and Newman 2004; van der Hoorn, Lippner, and Krioukov 2018). It has also been  
 89 proven useful in ecology, e.g. in species distribution models (Phillips, Anderson, and Schapire  
 90 2006) and macroecological models (Harte et al. 2008; Harte and Newman 2014). In network  
 91 ecology, MaxEnt has been used to predict the degree distribution of bipartite networks from the  
 92 number of species and the number of interactions (Williams 2011) and to predict interaction  
 93 strengths between species pairs using their relative abundances within an optimal transportation  
 94 theory regularized with information entropy (Stock, Poisot, and De Baets 2021). However, to  
 95 the best of our knowledge, MaxEnt has never been used to predict food-web structure directly,  
 96 even though food webs are among the most documented and widespread ecological networks  
 97 (Ings et al. 2009).

98 Food-web properties that can be derived using MaxEnt are varied and pertain to different ele-  
 99 ments of the network (i.e. at the species (node), the interaction (edge) or the community (net-  
 100 work) levels). Because MaxEnt is a method of finding least-biased probability distributions  
 101 given partial knowledge about a system, these properties need to be represented probabilistically.  
 102 For example, at the species level, MaxEnt can be used to predict the distribution of trophic levels  
 103 among species, as well as the distribution of species' vulnerability (number of predators) and  
 104 generality (number of prey). By contrast, at the interaction level, predictions can be made on  
 105 the distribution of interaction strengths in weighted food webs. At the community level, it can  
 106 generate probability distributions of many measures of their emerging structure and of networks  
 107 themselves (i.e. a probability distribution that specific network configurations are realized given  
 108 the model and constraints). Overall, the potential of this method in the study of food webs is  
 109 broad. The applicability and performance of MaxEnt mostly depend on the ecological infor-

110 mation available and on our capacity to find the right set of state variables that best represent  
111 natural systems and to translate them into appropriate statistical constraints. Having a validated  
112 maximum entropy model for the system at hand allows us to make least-biased predictions using  
113 a minimal amount of data, as well as identify the most important ecological processes shaping  
114 that system. In other words, MaxEnt is a valuable and underexploited tool in network ecology  
115 that allows us to better understand and predict the structure of ecological networks.

## 116 Analytical and heuristic approaches

117 In this contribution, we used two complementary approaches to predict the structure of food  
118 webs using the principle of maximum entropy. The first approach consists in deriving con-  
119 strained probability distributions of given network properties analytically, whereas the second  
120 approach consists in finding the adjacency matrix of maximum entropy heuristically, from which  
121 network properties can be measured. We compared our predictions against empirical data and  
122 null and neutral models commonly used in network ecology. We focus on deterministic and  
123 unweighted (Boolean) food webs in both approaches for data availability reasons. However, our  
124 framework can be applied to all types of ecological networks and a wide variety of measures.

125 For the first approach (analytic), we focus on species level properties. Specifically, we derived  
126 the joint degree distribution (i.e. the joint probability distribution that a species has a given  
127 number of prey and predators in its network) of maximum entropy using only the number of  
128 species  $S$  and the number of interactions  $L$  as state variables. Then, we predicted the degree  
129 distribution of maximum entropy directly from the joint degree distribution since the first is the  
130 sum of the marginal distributions of the second. Because of the scarcity of empirical data on  
131 the number of interactions in food webs, we present a method to predict  $L$  from  $S$  (Box 1), thus  
132 allowing the prediction of the joint degree distribution from  $S$  solely.

133 For the second approach (heuristic), we focus on network level properties. We used a flexible and  
134 heuristic model based on simulated annealing (an optimization algorithm) to find the network  
135 configuration *close* to maximum entropy and measured its structure. We developed this heuristic  
136 model because the analytical derivation of a maximum entropy graph model of food webs is  
137 difficult, and because this model is readily applicable to other types of ecological networks and

measures. Indeed, the mathematical representation of food webs (i.e. directed simple graphs frequently having self-loops) makes the optimization of maximum entropy graph models more complicated than with many other types of (non-ecological) networks. In other words, deriving a probability distribution on the graphs themselves is difficult when working with food webs.

We built two types of heuristic MaxEnt models depending on the constraint used. Our type I MaxEnt model uses the connectance of the network (i.e. the ratio  $L/S^2$ ) as a constraint, whereas our type II MaxEnt model uses the whole joint degree sequence as a constraint.

## 145 Data and code

We tested our analytical and heuristic MaxEnt models against open food-web data queried from three different sources and integrated into what we call our *complete dataset*. These sources include (1) terrestrial and aquatic food webs sampled globally and archived on the ecological interactions database Mangal; (2) freshwater stream food webs from New Zealand; and (3) aquatic food webs from Tuesday Lake (Michigan, United States). First, all food webs archived on `mangal.io` (Timothée Poisot et al. 2016; Banville, Vissault, and Poisot 2021) were directly queried from the database ( $N = 235$ ). Most ecological networks archived on Mangal are multi-layer networks, i.e. networks that describe different types of interactions. We kept all networks whose interactions were mainly of predation and herbivory types, and removed the largest network ( $S = 714$ ) for computational efficiency reasons. Then, to this set we added food webs from two different sources: the New Zealand dataset ( $N = 21$ ; J. P. Pomeranz et al. 2018) and the Tuesday Lake dataset ( $N = 2$ ; Cohen, Jonsson, and Carpenter 2003). Our complete dataset thus contained a total of 257 food webs. Of the New Zealand and Tuesday Lake datasets, 19 networks had data on species' relative abundances that were used in the neutral model (Box 2). These 19 food webs constitute our *abundance dataset*, which is a subset of our complete dataset.

All code and data to reproduce this article are available at the Open Science Framework. Data cleaning, simulations and analyses were conducted in Julia v1.6.1.

163 **Analytical maximum entropy models**

164 Our analytical approach is the most common way to use and develop maximum entropy models.  
165 As shown above, starting from a defined set of constraints, a mathematical expression for the  
166 target distribution is derived using the method of Lagrange multipliers. The derived distribution  
167 of maximum entropy is unique and least biased given the constraints used. Although we refer to  
168 this approach as analytic, finding the values of the Lagrange multipliers usually requires the use  
169 of numerical methods. Here we use MaxEnt to derive two species level properties in food webs:  
170 the joint degree distribution and the degree distribution. The degree distribution has driven the  
171 attention of ecologists because of its role in determining the assembly of ecological networks  
172 (Vázquez 2005), shaping their emerging structure (Fortuna et al. 2010), and understanding  
173 interaction mechanisms (Williams 2011). As noted above, although the degree distribution of  
174 maximum entropy has already been derived in bipartite networks (Williams 2011), we show  
175 in much greater details its mathematical derivation in food webs. But first, we derive the joint  
176 degree distribution, a related property that holds significantly more ecological information than  
177 the degree distribution.

178 **Joint degree distribution**

179 The joint degree distribution  $p(k_{in}, k_{out})$  of a food web with  $S$  species is a joint discrete proba-  
180 bility distribution describing the probability that a species has  $k_{in}$  predators and  $k_{out}$  prey, with  
181  $k_{in}$  and  $k_{out} \in [0, S]$ . Basal species (e.g., plants) have a  $k_{out}$  of 0, whereas top predators have  
182 a  $k_{in}$  of 0. In contrast, the maximum number of prey and predators a species can have is set  
183 by the number of species in the food web. Here we show how the joint degree distribution of  
184 maximum entropy can be obtained given knowledge of the number of species  $S$  and the number  
185 of interactions  $L$ .

186 We want to maximize Shannon's entropy

$$H = - \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S p(k_{in}, k_{out}) \log p(k_{in}, k_{out}) \quad (3)$$

<sup>187</sup> subject to the following constraints:

$$g_1 = \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S p(k_{in}, k_{out}) = 1; \quad (4)$$

$$g_2 = \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S k_{in} p(k_{in}, k_{out}) = \langle k_{in} \rangle = \frac{L}{S}; \quad (5)$$

$$g_3 = \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S k_{out} p(k_{in}, k_{out}) = \langle k_{out} \rangle = \frac{L}{S}. \quad (6)$$

<sup>188</sup> The first constraint  $g_1$  is our normalizing constraint, whereas the other two ( $g_2$  and  $g_3$ ) fix the  
<sup>189</sup> average of the marginal distributions of  $k_{in}$  and  $k_{out}$  to the linkage density  $L/S$ . It is important  
<sup>190</sup> to notice that  $\langle k_{in} \rangle = \langle k_{out} \rangle$  because every edge is associated to a predator and a prey. Therefore,  
<sup>191</sup> without using any further constraints, we would expect the joint degree distribution of maximum  
<sup>192</sup> entropy to be a symmetric probability distribution with regards to  $k_{in}$  and  $k_{out}$ . However, this  
<sup>193</sup> does not mean that the joint degree *sequence* will be symmetric, since the joint degree sequence  
<sup>194</sup> is essentially a random realization of its probabilistic counterpart.

<sup>195</sup> The joint probability distribution of maximum entropy given these constraints is found using the  
<sup>196</sup> method of Lagrange multipliers. To do so, we seek to maximize the following expression:

$$F = H - \lambda_1(g_1 - 1) - \lambda_2 \left( g_2 - \frac{L}{S} \right) - \lambda_3 \left( g_3 - \frac{L}{S} \right), \quad (7)$$

<sup>197</sup> where  $\lambda_1$ ,  $\lambda_2$ , and  $\lambda_3$  are the Lagrange multipliers. The probability distribution that maximizes  
<sup>198</sup> entropy is obtained by finding these values. As pointed out above,  $F$  is just Shannon's entropy  
<sup>199</sup> to which we added terms that each sums to zero (our constraints).  $F$  is maximized by setting to  
<sup>200</sup> 0 its partial derivative with respect to  $p(k_{in}, k_{out})$ . Because the derivative of a constant is zero,  
<sup>201</sup> this gives us:

$$\frac{\partial H}{\partial p(k_{in}, k_{out})} = \lambda_1 \frac{\partial g_1}{\partial p(k_{in}, k_{out})} + \lambda_2 \frac{\partial g_2}{\partial p(k_{in}, k_{out})} + \lambda_3 \frac{\partial g_3}{\partial p(k_{in}, k_{out})}. \quad (8)$$

202 Evaluating the partial derivatives with respect to  $p(k_{in}, k_{out})$ , we obtain:

$$-\log p(k_{in}, k_{out}) - 1 = \lambda_1 + \lambda_2 k_{in} + \lambda_3 k_{out}. \quad (9)$$

203 Then, solving eq. 9 for  $p(k_{in}, k_{out})$ , we obtain:

$$p(k_{in}, k_{out}) = \frac{e^{-\lambda_2 k_{in} - \lambda_3 k_{out}}}{Z}, \quad (10)$$

204 where  $Z = e^{1+\lambda_1}$  is called the partition function. The partition function ensures that probabilities  
205 sum to 1 (our normalization constraint). It can be expressed in terms of  $\lambda_2$  and  $\lambda_3$  as follows:

$$Z = \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S e^{-\lambda_2 k_{in} - \lambda_3 k_{out}}. \quad (11)$$

206 After substituting  $p(k_{in}, k_{out})$  in eq. 5 and eq. 6, we get a nonlinear system of two equations and  
207 two unknowns:

$$\frac{1}{Z} \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S k_{in} e^{-\lambda_2 k_{in} - \lambda_3 k_{out}} = \frac{L}{S}; \quad (12)$$

$$\frac{1}{Z} \sum_{k_{in}=0}^S \sum_{k_{out}=0}^S k_{out} e^{-\lambda_2 k_{in} - \lambda_3 k_{out}} = \frac{L}{S}. \quad (13)$$

208 We solved eq. 12 and eq. 13 numerically using the Julia library JuMP.jl v0.21.8 (Dunning,  
209 Huchette, and Lubin 2017). JuMP.jl supports nonlinear optimization problems by providing  
210 exact second derivatives that increase the accuracy and performance of its solvers. The estimated  
211 values of  $\lambda_2$  and  $\lambda_3$  can be substituted in eq. 10 to have a more workable expression for the joint  
212 degree distribution.

213 We assessed the empirical support of this expression using all food webs in our complete dataset.  
214 First, we predicted the joint degree distribution of maximum entropy for each of these food webs,  
215 i.e. using their number of species and number of interactions as state variables. Then, we sam-  
216 pled one realization of the joint degree sequence for each network using the probabilities given

217 by the joint degree distribution of maximum entropy, while fixing the total number of interac-  
218 tions. This gave us a random realization of the number of prey and predators for each species in  
219 each network. We standardized the predicted  $k_{out}$  and  $k_{in}$  of each species by the total number of  
220 species in their network to generate relative values, which can be compared across networks. In  
221 fig. 1 (left panels), we show the relationship between these relative  $k_{out}$  and  $k_{in}$  obtained from the  
222 joint degree distributions of maximum entropy (bottom panel) and this relationship using em-  
223 pirical values (top panel). We observe that our model predicts higher values of generality and  
224 vulnerability compared to empirical food webs (i.e. relative values of  $k_{out}$  and  $k_{in}$  both closer to  
225 1) for many species. In other words, our model predicts that species that have many predators  
226 also have more prey than what is observed empirically (and conversely). This is not surprising,  
227 given that our model did not include biological factors preventing generalist predators from hav-  
228 ing many prey. Nevertheless, with the exception of these generalist species, MaxEnt adequately  
229 predicts that most species have low generality and vulnerability values.

230 Examining the difference between predicted and empirical values for each species gives a slightly  
231 different perspective (right panel of fig. 1). To do so, we must first associate each of our predic-  
232 tions to a specific species in a network in order to make that comparison. Indeed, our predicted  
233 joint degree sequences have the same number of species (elements) as their empirical counter-  
234 parts, but they are species agnostic. In other words, instead of predicting a value for each species  
235 directly, we predicted the entire joint degree sequence without taking into account species' iden-  
236 tity. The challenge is thus to adequately associate predictions with empirical data. In fig. 1, we  
237 present these differences when species are ordered by their total degree in their respective net-  
238 works (i.e. by the sum of their in and out-degrees). This means that the species with the highest  
239 total degree in its network will be associated with the highest prediction, and so forth. Doing so,  
240 we see that species predicted to have a higher number of predators than what is observed gen-  
241 erally have a lower number of prey than what is observed (and conversely). This is because the  
242 difference in total degree ( $k_{out} + k_{in}$ ) between predictions and empirical data is minimized when  
243 species are ranked by their total degree (i.e. the average deviation of the sum of relative  $k_{out}$  and  
244  $k_{in}$  is close to 0 across all species). This result thus shows that the difference between predicted  
245 and empirical total degrees is low for most species when ordered by their total degrees. There

246 are no apparent biases towards in or out degrees. In fig. S1, we show how these differences  
247 change when species are instead ordered by their out-degrees (left panel) and in-degrees (right  
248 panel), respectively.

249 [Figure 1 about here.]

250 Another way to evaluate the empirical support of the predicted joint degree sequences is to com-  
251 pare their shape with the ones of empirical food webs. We can describe the shape of a joint degree  
252 sequence by comparing its marginal distributions with one another. To do so, we calculated the  
253 Kullback–Leibler (KL) divergence between the in and out-degree sequences sampled from the  
254 joint degree distribution of maximum entropy. Similarly, we calculated the divergence between  
255 the in and out-degree sequences obtained empirically. This allows us to compare the symmetry  
256 of empirical and predicted joint degree sequences (left panel of fig. 2). As we expected, our  
257 model predicts more similar in-degree and out-degree sequences than empirical data (shown  
258 by lower KL divergence values). However, this difference decreases with connectance (right  
259 panel of fig. 2). This might be due to the fact that food webs with a low connectance are harder  
260 to predict than food webs with a high connectance. Indeed, in low connectance systems, what  
261 makes two species interact might be more important for prediction than in high connectance  
262 systems, in which what prevents species from interacting might be more meaningful. This im-  
263 plies that more ecological information might be needed in food webs with a low connectance  
264 because more ecological processes determine interactions compared to non-interactions. There-  
265 fore, other ecological constraints might be needed to account for the asymmetry of the joint  
266 degree distribution, especially for networks with a lower connectance. However, our MaxEnt  
267 model was able to capture quite well the shape of the joint degree sequence for networks having  
268 a high connectance.

269 [Figure 2 about here.]

270 **Degree distribution**

271 The joint degree distribution derived using MaxEnt (eq. 10) can be used to obtain the degree  
272 distribution of maximum entropy. Indeed, the degree distribution  $p(k)$  represents the probability

273 that a species has  $k$  interactions in its food web, with  $k = k_{in} + k_{out}$ . It can thus be obtained  
 274 from the joint degree distribution as follows:

$$p(k) = \sum_{i=0}^k p(k_{in} = k - i, k_{out} = i).$$

275 The degree distribution could have also been obtained directly using the principle of maximum  
 276 entropy, as discussed in Williams (2011). This gives the following distribution:

$$p(k) = \frac{e^{-\lambda_2 k}}{Z}, \quad (14)$$

277 with  $Z = \sum_{k=0}^S e^{-\lambda_2 k}$ .

278 This can be solved numerically using the constraint of the average degree  $\langle k \rangle = \frac{2L}{S}$  of a species,  
 279 yielding an identical solution to the one obtained using the joint degree distribution as an inter-  
 280 mediate. Note that the mean degree is twice the value of the linkage density, because every link  
 281 must be counted twice when we add in and out-degrees together.

$$\frac{1}{Z} \sum_{k=0}^S k e^{-\lambda_2 k} = \frac{2L}{S} \quad (15)$$

282 One aspect of the degree distribution that informs us of its ecological realism is the number of  
 283 isolated species it predicts. As MacDonald, Banville, and Poisot (2020) pointed out, the size of  
 284 food webs should at least be of  $S - 1$  interactions, since a lower number would yield isolated  
 285 species, i.e. species without any predators or prey. Because non-basal species must eat to survive,  
 286 isolated species could indicate that other species are missing or they could simply be removed  
 287 from the food web. In fig. S2, we show that the degree distribution of maximum entropy, given  
 288  $S$  and  $L$ , gives very low probabilities that a species will be isolated in its food web (i.e. having  
 289  $k = 0$ ) above the  $S - 1$  threshold. However, under our purely information-theoretic model,  
 290 the probability that a species is isolated is quite high when the total number of interactions is  
 291 below  $S - 1$ . Moreover, the expected proportion of isolated species rapidly declines by orders  
 292 of magnitude with increasing numbers of species and interactions. This supports the ecological  
 293 realism of the degree distribution of maximum entropy derived above. Nevertheless, ecologists

294 wanting to model a system without allowing isolated species could simply change the lower limit  
295 of  $k$  to 1 in eq. 15 and solve the resulting equation numerically.

## 296 Box 1 - Working with predicted numbers of interactions

297 Our analytical MaxEnt models require information on the number of species and the number of  
298 interactions. However, since the later is rarely measured empirically, ecologists might need to  
299 use predictive models to estimate the total number of interactions in a food web before using  
300 MaxEnt. Here we illustrate how this can be done by combining both models sequentially.

301 We used the flexible links model of MacDonald, Banville, and Poisot (2020) to predict the  
302 number of interactions from the number of species. The flexible links model, in contrast to other  
303 predictive models of the number of interactions, incorporates meaningful ecological constraints  
304 into the prediction of  $L$ , namely the minimum  $S - 1$  and maximum  $S^2$  numbers of interactions  
305 in food webs. It estimates the proportion of the  $S^2 - (S - 1)$  *flexible links* that are realized.  
306 More precisely, this model states that the number of *realized* flexible links (or interactions)  
307  $L_{FL}$  in a food web represents the number of realized interactions above the minimum (i.e.  $L =$   
308  $L_{FL} + S - 1$ ) and is obtained from a beta-binomial distribution with  $S^2 - (S - 1)$  trials and  
309 parameters  $\alpha = \mu e^\phi$  and  $\beta = (1 - \mu) e^\phi$ :

$$L_{FL} \sim \text{BB}(S^2 - (S - 1), \mu e^\phi, (1 - \mu) e^\phi), \quad (16)$$

310 where  $\mu$  is the average probability across food webs that a flexible link is realized, and  $\phi$  is the  
311 concentration parameter around  $\mu$ .

312 We fitted the flexible links model on all food webs in our complete dataset, and estimated the  
313 parameters of eq. 16 using a Hamiltonian Monte Carlo sampler with static trajectory (1 chain  
314 and 3000 iterations):

$$[\mu, \phi | \mathbf{L}, \mathbf{S}] \propto \prod_{i=1}^m \text{BB}(L_i - (S_i - 1) | S_i^2 - (S_i - 1)), \mu e^\phi, (1 - \mu) e^\phi) \times \text{B}(\mu | 3, 7) \times \mathcal{N}(\phi | 3, 0.5), \quad (17)$$

315 where  $m$  is the number of food webs ( $m = 257$ ) and  $\mathbf{L}$  and  $\mathbf{S}$  are respectively the vectors of their  
 316 numbers of interactions and numbers of species. Our weakly-informative prior distributions  
 317 were chosen following MacDonald, Banville, and Poisot (2020), i.e. a beta distribution for  $\mu$   
 318 and a normal distribution for  $\phi$ . The Monte Carlo sampling of the posterior distribution was  
 319 conducted using the Julia library Turing v0.15.12.

320 The flexible links model is a generative model, i.e. it can generate plausible values of the pre-  
 321 dicted variable. We thus simulated 1000 values of  $L$  for different values of  $S$  using the joint  
 322 posterior distribution of our model parameters (eq. 17), and calculated the mean degree for each  
 323 simulated value. The resulting distributions are shown in the left panel of fig. 3 for three different  
 324 values of species richness. In the right panel of fig. 3, we show how the probability distribution  
 325 for the mean degree constraints can be used to generate a distribution of maximum entropy de-  
 326 gree distributions, since each simulated value of mean degree generates a different maximum  
 327 entropy degree distribution (eq. 14 and eq. 15).

328 [Figure 3 about here.]

## 329 **Box 2 - Corresponding null and neutral models**

### 330 **Null models (types I and II)**

331 The predictions of our heuristic maximum entropy models were compared against two topologi-  
 332 cal null models. These null models use the same ecological information as our heuristic models  
 333 and thus constitute an adequate baseline for comparison. The first is the type I null model of  
 334 Fortuna and Bascompte (2006), in which the probability that a species  $i$  predares on another  
 335 species  $j$  is given by

$$p(i \rightarrow j) = \frac{L}{S^2}. \quad (18)$$

336 The second is the type II null model of Bascompte et al. (2003), in which the probability of  
 337 interaction is instead given by

$$p(i \rightarrow j) = \frac{1}{2} \left( \frac{k_{in}(j)}{S} + \frac{k_{out}(i)}{S} \right), \quad (19)$$

338 where  $k_{in}(j)$  and  $k_{out}(i)$  are the in and out-degrees of species  $j$  and  $i$ , respectively. The type I  
 339 null model is based on connectance, whereas the type II null model is based on the joint degree  
 340 sequence. Therefore, the type I and II topological null models correspond to our type I and II  
 341 heuristic MaxEnt models, respectively, since they use similar constraints.

342 We generated probabilistic networks using both types of null models for all empirical food webs  
 343 in our complete dataset. Then, we converted these networks to adjacency matrices of Boolean  
 344 values by generating 100 random networks for each of these probabilistic webs, and kept the  $L$   
 345 entries that were sampled the most amount of times, with  $L$  given by the number of interactions  
 346 in each food web. This ensured that the resulting null networks had the same number of inter-  
 347 actions as their empirical counterparts. Thus, for each null model, we ended up with one null  
 348 adjacency matrix for each empirical network.

### 349 Neutral model

350 We also compared our heuristic MaxEnt models with a neutral model of relative abundances, in  
 351 which the probabilities of interaction are given by

$$p(i \rightarrow j) \propto \frac{n_i}{N} \times \frac{n_j}{N}, \quad (20)$$

352 where  $n_i$  and  $n_j$  are the abundances (or biomass) of both species, and  $N$  is the total abundance  
 353 (or biomass) of all species in the network. We generated neutral abundance matrices for all em-  
 354 pirical food webs in our abundance dataset, and converted these weighted networks to adjacency

355 matrices of Boolean values using the same method as the one we used for our null models.

356 **Heuristic maximum entropy models**

357 With the analytical approach, we showed how important measures of food-web structure (e.g.,  
358 the degree distribution and the joint degree distribution) can be derived with the principle of  
359 maximum entropy using minimal knowledge about a biological community. This type of mod-  
360 els, although useful to make least-biased predictions of many network properties, can be hard  
361 to apply for other measures. Indeed, there are dozens of measures of network structure (Del-  
362 mas et al. 2019) and many are not directly calculated with mathematical equations, but are  
363 instead estimated algorithmically. Moreover, the applicability of this method to empirical sys-  
364 tems is limited by the state variables we can actually measure and use. Here, we propose a more  
365 flexible method to predict many measures of network structure simultaneously, i.e. by finding  
366 heuristically the network configuration having maximum entropy given partial knowledge of its  
367 emerging structure.

368 **From Shannon's to SVD entropy**

369 The principle of maximum entropy can be applied on the network itself if we decompose its  
370 adjacency matrix into a non-zero vector of relative values. This is a necessary step when work-  
371 ing with food webs, which are frequently expressed as a matrix  $A = [a_{ij}]$  of Boolean values  
372 representing the presence ( $a_{ij} = 1$ ) or absence ( $a_{ij} = 0$ ) of an interaction between two species  $i$   
373 and  $j$ . Knowing one or many properties of a food web of interest (e.g., its number of species and  
374 number of interactions), we can simulate its adjacency matrix randomly by using these known  
375 ecological information to constrain the space of potential networks. The entropy of this hypo-  
376 thetical matrix can then be measured after decomposing it into appropriate values. Simulating  
377 a series of networks until we find the one having the highest entropy allows us to search for the  
378 most complex food-web configuration given the ecological constraints used. This configuration  
379 is the least biased one considering the information available. In other words, the most we can  
380 say about a network's adjacency matrix, without making more assumptions than the ones given

381 by our incomplete knowledge of its structure, is the one of maximum entropy. Generating the  
382 most complex network that corresponds to this structure allows us to explore more easily other  
383 properties of food webs under MaxEnt.

384 Shannon's entropy can only be calculated on conventional probability distributions such as the  
385 joint degree distribution. This is an issue when working with the adjacency matrix of ecological  
386 networks. For this reason, we need to use another measure of entropy if we want to predict a  
387 network's configuration directly using MaxEnt. We used the SVD entropy as our measure of  
388 entropy, which is an application of Shannon's entropy to the relative non-zero singular values  
389 of a truncated singular value decomposition (t-SVD; Strydom, Dalla Riva, and Poisot 2021) of  
390 a food web's Boolean adjacency matrix. We measured SVD entropy as follows:

$$J = - \sum_{i=1}^R s_i \log s_i, \quad (21)$$

391 where  $s_i$  are the relative singular values of the adjacency matrix ( $s_i = \sigma_i / \sum_{i=1}^R \sigma_i$ , where  $\sigma_i$   
392 are the singular values). Note that the distribution of relative singular values is analogous to  
393 a probability distribution, with  $0 < s_i < 1$  and  $\sum s_i = 1$ . This measure also satisfies all four  
394 properties of an appropriate entropy measure above-mentioned, while being a proper measure of  
395 the internal complexity of food webs (Strydom, Dalla Riva, and Poisot 2021). Following Stry-  
396 dom, Dalla Riva, and Poisot (2021), we standardized this measure with the rank  $R$  of the matrix  
397 (i.e.  $J / \ln(R)$ ) to account for the difference in dimensions between networks (*sensu* Pielou's  
398 evenness; Pielou 1975).

### 399 **Types I and II heuristic MaxEnt models**

400 We used SVD entropy to predict the network configuration of maximum entropy (i.e. of max-  
401 imum complexity) heuristically given different constraints for all food webs in our complete  
402 dataset. We built two types of heuristic MaxEnt models that differ on the constraint used. The  
403 type I heuristic MaxEnt model is based on connectance, whereas the type II heuristic MaxEnt  
404 model is based on the joint degree sequence. These models are thus based on the same con-  
405 straints as the types I (Fortuna and Bascompte 2006) and II (Bascompte et al. 2003) null models

406 (Box 2) frequently used to generate random networks topologically. This allows direct com-  
407 parison of the performance of null and heuristic MaxEnt models in reproducing the emerging  
408 structure of empirical food webs.

409 For each network in our complete dataset, we estimated their configuration of maximum entropy  
410 given each of these constraints. For both types of heuristic MaxEnt models, we used a simulated  
411 annealing algorithm with 4 chains, 2000 steps and an initial temperature of 0.2. For each food  
412 web, we first generated one random Boolean matrix per chain while fixing the number of species.  
413 We also maintained the total number of interactions (i.e. the sum of all elements in the matrix) in  
414 the type I MaxEnt model and the joint degree sequence (i.e. the rows and columns sums) in the  
415 type II MaxEnt model. These were our initial configurations. Then, we swapped interactions se-  
416 quentially while maintaining the original connectance or joint degree sequence. Configurations  
417 with a higher SVD entropy than the previous one in the chain were always accepted, whereas  
418 they were accepted with a probability conditional to a decreasing temperature when lower. The  
419 final configuration with the highest SVD entropy among the four chains constitute the estimated  
420 maximum entropy configuration of a food web given the constraint used.

## 421 Structure of MaxEnt food webs

422 We measured various properties of these configurations of maximum entropy to evaluate how  
423 well they reproduce the structure of sampled food webs. Specifically, we evaluated their nest-  
424 edness  $\rho$ , their maximum trophic level  $maxtl$ , their network diameter  $diam$ , their average maxi-  
425 mum similarity between species pairs  $MxSim$  (Williams and Martinez 2000), their proportion  
426 of cannibal species  $Cannib$ , their proportion of omnivorous species  $Omniv$ , their SVD entropy,  
427 and their motifs profile. Nestedness indicates how much the diet of specialist species is a subset  
428 of the one of generalists (Delmas et al. 2019) and was measured using the spectral radius of  
429 the adjacency matrix (Staniczenko, Kopp, and Allesina 2013). In turn, the network diameter  
430 represents the longest of the shortest paths between all species pair (Albert and Barabasi 2002).  
431 Further, cannibal species are species that can eat individuals of their own species (i.e. species  
432 having self loops), whereas omnivorous species can prey on different trophic levels (Williams  
433 and Martinez 2000). Finally, a motifs profile represents the proportion of three-species motifs

434 (subnetworks), which can be considered as simple building blocks of ecological networks (Milo  
435 et al. 2002; Stouffer et al. 2007). All of these properties are relatively easy to measure and were  
436 chosen based on their ecological importance and prevalent use in network ecology (McCann  
437 2011; Delmas et al. 2019).

438 We compared the performance of both heuristic MaxEnt models in predicting these measures to  
439 the one of the null (tbl. 1) and neutral models (tbl. 2). Overall, we found that the models based  
440 on the joint degree sequence (i.e. the type II null and heuristic MaxEnt models) reproduced the  
441 structure of empirical food webs much better than the ones based on connectance (i.e. the type  
442 I null and heuristic MaxEnt models). This suggests that the predictive power of connectance  
443 might be more limited than what was previously suggested (Timothée Poisot and Gravel 2014).  
444 On the other hand, the neutral model of relative abundances was surprisingly good at predict-  
445 ing the maximum trophic level and the network diameter. However, with the exception of the  
446 network diameter, the type II heuristic MaxEnt model was better at predicting network structure  
447 than the neutral model for most measures considered. This might be because, although neutral  
448 processes are important, they act in concert with niche processes in determining species inter-  
449 actions Canard et al. (2014). The joint degree sequence encodes information on both neutral  
450 and niche processes because the number of prey and predators a species has is determined by its  
451 relative abundance and biological traits. These results thus show that having information on the  
452 number of prey and predators for each species substantially improves the prediction of food-web  
453 structure, both compared to models solely based on connectance and to the ones solely based on  
454 species relative abundances.

455 Next, the predictions of the type II heuristic MaxEnt model can be compared to its null model  
456 counterpart. On average, the type II heuristic MaxEnt model was better at predicting nestedness  
457 ( $0.62 \pm 0.08$ ) than its corresponding null model ( $0.73 \pm 0.05$ ; empirical networks:  $0.63 \pm 0.09$ ) for  
458 networks in our complete dataset (tbl. 1). This might in part be due to the fact that nestedness was  
459 calculated using the spectral radius of the adjacency matrix, which directly leverages information  
460 on the network itself just like the heuristic MaxEnt model. The proportion of self-loops (cannibal  
461 species) was also better predicted by the type II heuristic MaxEnt model in comparison to the  
462 type II null model. However, the type II null model was better at predicting network diameter

463 and average maximum similarity between species pairs, and predictions of the maximum trophic  
 464 level and the proportion of omnivorous species were similar between both types of models. We  
 465 believe that this is because increasing the complexity of a food web might increase its average  
 466 and maximum food-chain lengths. In comparison, the null model was more stochastic and does  
 467 not necessarily produce more complex food webs with longer food-chain lengths.

Table 1: Standardized mean difference between predicted network measures and empirical data for all food webs in our complete dataset ( $N = 257$ ). Positive (negative) values indicate that the measure is overestimated (underestimated) on average. Empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake food webs. Null 1: Type I null model based on connectance. MaxEnt 1: Type I heuristic MaxEnt model based on connectance. Null 2: Type II null model based on the joint degree sequence. MaxEnt 2: Type II heuristic MaxEnt model based on the joint degree sequence.  $\rho$ : nestedness measured by the spectral radius of the adjacency matrix. *maxtl*: maximum trophic level. *diam*: network diameter. *MxSim*: average maximum similarity between species pairs. *Cannib*: proportion of cannibal species (self loops). *Omniv*: proportion of omnivorous species. *entropy*: SVD entropy.

model	rho	maxtl	diam	MxSim	Cannib	Omniv	entropy
null 1	-0.167	0.980	1.428	-0.502	2.007	1.493	0.056
MaxEnt 1	-0.226	0.831	1.274	-0.524	1.982	1.863	0.106
null 2	0.160	-0.125	0.016	0.007	1.078	0.559	-0.023
MaxEnt 2	-0.015	0.178	0.565	-0.282	0.698	0.589	0.058

Table 2: Standardized mean difference between predicted network measures and empirical data for all food webs in our abundance dataset ( $N = 19$ ). Positive (negative) values indicate that the measure is overestimated (underestimated) on average. Empirical networks include the New Zealand and Tuesday Lake food webs having abundance data. Neutral: Neutral model of relative abundances. Null 1: Type I null model based on connectance. MaxEnt 1: Type I heuristic MaxEnt model based on connectance. Null 2: Type II null model based on the joint degree sequence. MaxEnt 2: Type II heuristic MaxEnt model based on the joint degree sequence.  $\rho$ : nestedness measured by the spectral radius of the adjacency matrix. *maxtl*: maximum trophic level. *diam*: network diameter. *MxSim*: average maximum similarity between species pairs. *Cannib*: proportion of cannibal species (self loops). *Omniv*: proportion of omnivorous species. *entropy*: SVD entropy.

model	rho	maxtl	diam	MxSim	Cannib	Omniv	entropy
neutral	0.367	-0.090	0.027	0.266	6.870	0.576	-0.083
null 1	-0.134	0.950	1.919	-0.369	2.077	0.614	0.068

model	rho	maxtl	diam	MxSim	Cannib	Omniv	entropy
MaxEnt 1	-0.229	1.020	1.946	-0.355	2.215	0.801	0.121
null 2	0.128	-0.115	-0.135	0.157	1.444	0.029	-0.021
MaxEnt 2	-0.010	0.054	0.243	-0.062	-0.038	0.083	0.038

468 Despite this increase in maximum trophic level and network diameter in MaxEnt food webs, we  
 469 found that empirical food webs are close to their maximum entropy given a fixed joint degree  
 470 sequence (fig. S3). Empirical food webs in the complete dataset had an SVD entropy of  $0.89 \pm$   
 471 0.04, compared to an SVD entropy of  $0.94 \pm 0.03$  for networks generated using the type II  
 472 heuristic MaxEnt model. The relationship between the SVD entropy of empirical food webs  
 473 and their maximum entropy is plotted in the last panel of fig. 4. As expected, the SVD entropy  
 474 of maximum entropy food webs was higher than that of empirical food webs for almost all  
 475 networks, confirming that our method indeed generated more complex networks. Moreover, we  
 476 found no to a weak relationship between the increase in SVD entropy and the number of species,  
 477 the number of interactions, and connectance (fig. S4). This suggests that the slight increase in  
 478 entropy between empirical food webs and their maximum entropy configuration was sufficient  
 479 to modify some of their properties, regardless of their number of species and their number of  
 480 interactions.

481 A direct comparison of the structure of maximum entropy food webs, constrained by the joint  
 482 degree sequence, with empirical data also supports the results depicted in [tbl. 1](#). Indeed, in [fig. 4](#)  
 483 we show how well empirical measures are predicted by the type II heuristic MaxEnt model. In  
 484 accordance with our previous results, we found that nestedness was very well predicted by our  
 485 model. However, the model overestimated the maximum trophic level and network diameter,  
 486 especially when the sampled food web had intermediate values of these measures. In [fig. S5](#),  
 487 we show that the pairwise relationships between the four measures in [fig. 4](#) and species richness  
 488 in empirical food webs are similar (in magnitude and sign) to the ones found in food webs  
 489 generated using the type II heuristic MaxEnt model. This indicates that the number of species  
 490 in the network does not seem to impact the ability of the model to reproduce food-web structure.

491

[Figure 4 about here.]

492 Notwithstanding its difficulties in reproducing adequate measures of food-chain lengths, the  
493 type II heuristic MaxEnt model can predict surprisingly well the motifs profile. Motifs are the  
494 backbone of complex ecological networks from which network structure is built upon and play  
495 a crucial role in community dynamics and assembly (Stouffer and Bascompte 2011). For this  
496 reason, the motifs profile can act as an effective ecological constraint shaping species interactions  
497 networks, and thus constitute a substantial source of ecological information. In fig. 5, we show  
498 that the motifs profile of networks generated using the type II heuristic MaxEnt model accurately  
499 reproduced the one of empirical data. This model made significantly better predictions than  
500 the ones based on connectance and the type II null model based on the joint degree sequence.  
501 This is also shown in fig. 6, where we see that the relationships between motifs proportions in  
502 empirical food webs are very similar to the ones in networks generated using the type II heuristic  
503 MaxEnt model. This is in contrast with the type I null and MaxEnt models based on connectance,  
504 which produced opposite relationships than what was observed empirically. Our findings thus  
505 suggest that increasing food-web complexity within a maximum entropy framework based on  
506 the joint degree sequence does not alter the proportion of three-species motifs, but might alter  
507 their position relative to one another.

508

[Figure 5 about here.]

509

[Figure 6 about here.]

510 One of the challenges in implementing and validating a maximum entropy model is to discover  
511 where its predictions break down. The results depicted in tbl. 1 and fig. 4 show that our type II  
512 heuristic MaxEnt model can capture many high-level properties of food webs, but does a poor  
513 job of capturing others. This suggests that, although the joint degree sequence is an important  
514 driver of food-web structure, other ecological constraints might be needed to account for some  
515 emerging food-web properties, especially the ones regarding food-chain lengths. Nevertheless,  
516 fig. 5 and fig. 6 show that the model can reproduce surprisingly well the motifs profile, one of the  
517 most ecologically informative properties of food webs. This suggests that the emerging structure

518 of food webs is mainly driven by their joint degree sequence, although higher-level properties  
519 might be needed to ensure that food-chain lengths fall within realistic values.

520 **Conclusion**

521 The principle of maximum entropy is a robust mathematical method of finding least-biased  
522 probability distributions that have some specified properties given by prior knowledge about a  
523 system. We first applied this conventional MaxEnt approach on food webs to predict species  
524 level properties, namely the joint degree distribution and the degree distribution of maximum  
525 entropy given known numbers of species and interactions. We found that the joint degree dis-  
526 tributions of maximum entropy had a similar shape to the ones of empirical food webs in high-  
527 connectance systems. However, these MaxEnt distributions were more symmetric than the ones  
528 of empirical food webs when connectance was low, which suggests that other constraints might  
529 be needed to improve these predictions in low-connectance systems. Then, we used a slightly  
530 different approach that aimed at finding heuristically the network configuration with the highest  
531 SVD entropy, i.e. whose vector of relative singular values has maximum entropy. This network  
532 of maximum entropy is the most complex, or random, given the specified structure. We found  
533 that the heuristic maximum entropy model based on connectance did not predict the structure  
534 of sampled food webs very well. However, the heuristic maximum entropy model based on the  
535 entire joint degree sequence, i.e. on the number of prey and predators for each species, gave  
536 more convincing results. Indeed, this model reproduced food-web structure surprisingly well,  
537 including the highly informative motifs profile. Nevertheless, it was not able to predict realistic  
538 food-chain lengths.

539 Our results bring to the forefront the role of the joint degree distribution in shaping food-web  
540 structure. This echoes the work of Fortuna et al. (2010), who found that the degree distribution  
541 of ecological networks drive their emerging structure such as their nestedness and modularity.  
542 Network ecologists tend to focus on several measures of food webs when studying the ecological  
543 consequences of their structure (McCann 2011; Delmas et al. 2019). In fact, following Williams  
544 (2011), we believe there is a lot more ecological information in the deviation between these

545 properties in empirical systems and in their maximum entropy configuration given a fixed joint  
546 degree sequence.

## 547 Alternative MaxEnt models

548 In this contribution, we used a method based on simulated annealing to find the network config-  
549 uration with the highest SVD entropy while fixing some aspects of its structure. However, there  
550 are different ways to generate adjacency matrices using MaxEnt. Another technique, also based  
551 on simulated annealing, could begin by generating a food web randomly with fixed numbers of  
552 species and interactions and calculating its joint degree distribution. Pairs of interactions could  
553 then be swapped sequentially until we minimize the divergence between the calculated joint de-  
554 gree distribution and the one of maximum entropy obtained analytically. In that case, this is the  
555 entropy of the joint degree distribution that would be maximized, not the one of the network's  
556 topology. To a certain extent, this method would thus bridge the gap between the analytical and  
557 heuristic approaches presented in this article. More research is needed to compare the quality  
558 of different methods generating adjacency matrices of food webs using MaxEnt.

559 Maximum entropy graph models are another type of methods that predict a distribution of ad-  
560 jacency matrices under soft or hard constraints (e.g., Park and Newman 2004; Cimini et al.  
561 2019). Under hard constraints, every network with a non-zero probability exactly satisfies the  
562 constraints on its structure. This is in contrast with soft constraints, which require that networks  
563 satisfy them on average (i.e. many networks with a non-zero probability do not have the ex-  
564 act structure set by the constraints). Maximum entropy graph models are helpful because they  
565 can provide probability distributions for many network properties by measuring the structure  
566 of all adjacency matrices with a non-zero probability. However, we consider that our approach  
567 based on simulated annealing is more flexible and more computationally efficient. Indeed, many  
568 measures of food-web structure are hard to translate into mathematical constraints. Moreover,  
569 because food webs are directed networks that can have self-loops, it makes the mathematical  
570 derivation of maximum entropy graph models difficult. We believe that identifying heuristi-  
571 cally what really constrains the topology of food webs is a useful first step before attempting to  
572 derive the mathematical formulation of a maximum entropy graph model for food webs.

573 **Applications**

574 Our analytical and heuristic models can be applied for different purposes. First, they could be  
575 used to generate first-order approximations of a network's properties when state variables are  
576 known empirically. For example, knowing the number of species in an ecological community,  
577 we can predict its number of interactions using the flexible links model and then predict its joint  
578 degree distribution with minimal biases using the principle of maximum entropy. This could  
579 prove particularly useful when predicting network structure at large spatial scales, subdividing  
580 the study area into smaller communities (e.g., grid cells). Indeed, because species richness and  
581 other ecological data are increasingly abundant (e.g., Dickinson, Zuckerberg, and Bonter 2010),  
582 validated MaxEnt models can be used to respond to a wider range of macroecological questions  
583 regarding food webs.

584 Second, our analytical model can be used to generate informative priors in Bayesian analyses of  
585 the structure of ecological networks (e.g., Cirtwill et al. 2019). Indeed, the probability distri-  
586 bution of maximum entropy derived using MaxEnt can be used as a prior that can be constantly  
587 updated with novel data. For instance, if we know the number of species and the number of  
588 interactions, we can get the degree distribution of maximum entropy, as shown in this contribu-  
589 tion. The degree distribution represents the probability that a species can interact (as a predator  
590 or a prey) with a number of other species. Data on species interactions can be used to update  
591 the prior degree distribution to generate a more accurate posterior distribution, thus improving  
592 our description and understanding of the system.

593 Third, our analytical and heuristic models can be used to make better predictions of pairwise  
594 species interactions by constraining the space of feasible networks, as discussed in Strydom et  
595 al. (2021). In other words, we can use the network configuration and/or specific measures of  
596 food-web structure derived using MaxEnt to ensure that our predictions of interspecific interac-  
597 tions form feasible networks. This means that the probability that two species interact can be  
598 conditional on the network structure and on the probability of interactions of all other species  
599 pairs. For that purpose, MaxEnt can be used to predict network structure when other data is  
600 lacking.

601 Finally, our analytical and heuristic models can be used as alternative null models of ecological  
602 networks to better understand and identify the ecological processes driving food-web structure.  
603 Indeed, these mechanisms can be better described when analyzing the deviation of empirical  
604 data from MaxEnt predictions. A strong deviation would indicate that ecological mechanisms  
605 not encodes in the statistical constraints is at play for the system at hand. If deviations are sys-  
606 tematic, the maximum entropy model might need to be revised to include appropriate ecological  
607 constraints. This revision process helps us reflect on and identify what really constrains food-  
608 web structure.

609 The principle of maximum entropy can thus be applied for both the prediction and understanding  
610 of natural systems. Therefore, the model interpretation depends on how we use it. It can be used  
611 as a baseline distribution to identify the ecological constraints organizing natural systems. It can  
612 also be used as predictions of ecological systems. This distinction between understanding and  
613 predicting is essential when using and interpreting MaxEnt models.

## 614 **Final remarks**

615 One of the biggest challenges in using the principle of maximum entropy is to identify the set  
616 of state variables that best reproduce empirical data. We found that the number of species and  
617 the number of interactions are important state variables for the prediction of the joint degree  
618 distribution. Similarly, we found that the numbers of prey and predators for each species in a  
619 food web are important state variables for the prediction of the network configuration. However,  
620 our predictions overestimated the symmetry of the joint degree distribution for our analytical  
621 model and the maximum trophic level and network diameter for our heuristic model. We should  
622 thus continue to play the ecological detective to find these other topological constraints that  
623 would improve the predictions of MaxEnt models and help us understand better what really  
624 drives food-web structure.

625 **Acknowledgments**

626 We acknowledge that this study was conducted on land within the traditional unceded territory  
627 of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and Omàmiwinini-  
628 wak nations. This work was supported by the Institute for Data Valorisation (IVADO) and the  
629 Natural Sciences and Engineering Research Council of Canada (NSERC) Collaborative Re-  
630 search and Training Experience (CREATE) program, through the Computational Biodiversity  
631 Science and Services (BIOS<sup>2</sup>) program.

632 **References**

- 633 Albert, R., and A. L. Barabasi. 2002. “Statistical Mechanics of Complex Networks.” *Reviews*  
634 *of Modern Physics* 74 (1): 47–97. <https://doi.org/10.1103/RevModPhys.74.47>.
- 635 Banville, Francis, Steve Vissault, and Timothée Poisot. 2021. “Mangal.jl and EcologicalNet-  
636 works.jl: Two Complementary Packages for Analyzing Ecological Networks in Julia.” *Journal*  
637 *of Open Source Software* 6 (61): 2721. <https://doi.org/10.21105/joss.02721>.
- 638 Bartomeus, Ignasi, Dominique Gravel, Jason M. Tylianakis, Marcelo A. Aizen, Ian A. Dickie,  
639 and Maud Bernard-Verdier. 2016. “A Common Framework for Identifying Linkage Rules  
640 Across Different Types of Interactions.” *Functional Ecology* 30 (12): 1894–1903. <https://doi.org/10.1111/1365-2435.12666>.
- 642 Bascompte, J., P. Jordano, C. J. Melian, and J. M. Olesen. 2003. “The Nested Assembly of  
643 Plant-Animal Mutualistic Networks.” *Proceedings of the National Academy of Sciences*  
644 *of the United States of America* 100 (16): 9383–87. <https://doi.org/10.1073/pnas.1633576100>.
- 646 Beck, Christian. 2009. “Generalised Information and Entropy Measures in Physics.” *Contem-  
647 porary Physics* 50 (4): 495–510. <https://doi.org/10.1080/00107510902823517>.
- 648 Canard, E. F., N. Mouquet, D. Mouillot, M. Stanko, D. Miklisova, and D. Gravel. 2014. “Empir-  
649 ical Evaluation of Neutral Interactions in Host-Parasite Networks.” *The American Naturalist*  
650 183 (4): 468–79. <https://doi.org/10.1086/675363>.

- 651 Cimini, Giulio, Tiziano Squartini, Fabio Saracco, Diego Garlaschelli, Andrea Gabrielli, and  
652 Guido Caldarelli. 2019. “The Statistical Physics of Real-World Networks.” *Nature Reviews*  
653 *Physics* 1 (1): 58–71. <https://doi.org/10.1038/s42254-018-0002-6>.
- 654 Cirtwill, Alyssa R., Anna Eklof, Tomas Roslin, Kate Wootton, and Dominique Gravel. 2019.  
655 “A Quantitative Framework for Investigating the Reliability of Empirical Network Construc-  
656 tion.” *Methods in Ecology and Evolution* 10 (6): 902–11. <https://doi.org/10.1111/2041-210X.13180>.
- 658 Cohen, Joel E., Tomas Jonsson, and Stephen R. Carpenter. 2003. “Ecological Community  
659 Description Using the Food Web, Species Abundance, and Body Size.” *Proceedings of*  
660 *the National Academy of Sciences* 100 (4): 1781–86. <https://doi.org/10.1073/pnas.232715699>.
- 662 Delmas, Eva, Mathilde Besson, Marie Hélène Brice, Laura A. Burkle, Giulio V. Dalla Riva,  
663 Marie Josée Fortin, Dominique Gravel, et al. 2019. “Analysing Ecological Networks of  
664 Species Interactions.” *Biological Reviews*. <https://doi.org/10.1111/brv.12433>.
- 665 Dickinson, Janis L., Benjamin Zuckerberg, and David N. Bonter. 2010. “Citizen Science as an  
666 Ecological Research Tool: Challenges and Benefits.” In *Annual Review of Ecology, Evo-*  
667 *lution, and Systematics, Vol 41*, edited by D. J. Futuyma, H. B. Shafer, and D. Simberloff,  
668 41:149–72. Palo Alto: Annual Reviews. <https://doi.org/10.1146/annurev-ecolsys-102209-144636>.
- 669 Dunne, Jennifer A., Richard J. Williams, and Neo D. Martinez. 2002. “Network Structure and  
670 Biodiversity Loss in Food Webs: Robustness Increases with Connectance.” *Ecology Letters*  
671 5 (4): 558–67. <https://doi.org/10.1046/j.1461-0248.2002.00354.x>.
- 672 Dunning, Iain, Joey Huchette, and Miles Lubin. 2017. “JuMP: A Modeling Language for  
673 Mathematical Optimization.” *SIAM Review* 59 (2): 295–320. <https://doi.org/10.1137/15M1020575>.
- 675 Fortuna, M. A., and J. Bascompte. 2006. “Habitat Loss and the Structure of Plant-Animal  
676 Mutualistic Networks.” *Ecology Letters* 9 (3): 278–83. <https://doi.org/10.1111/j.1461-0248.2005.00868.x>.

- 678 Fortuna, M. A., Daniel B. Stouffer, Jens M. Olesen, Pedro Jordano, David Mouillot, Boris R.  
679 Krasnov, Robert Poulin, and Jordi Bascompte. 2010. “Nestedness Versus Modularity in  
680 Ecological Networks: Two Sides of the Same Coin?” *Journal of Animal Ecology* 79 (4):  
681 811–17. <https://doi.org/10.1111/j.1365-2656.2010.01688.x>.
- 682 Frank, S. A., and E. Smith. 2011. “A Simple Derivation and Classification of Common Prob-  
683 ability Distributions Based on Information Symmetry and Measurement Scale.” *Journal of*  
684 *Evolutionary Biology* 24 (3): 469–84. <https://doi.org/10.1111/j.1420-9101.2010.02204.x>.
- 686 Gómez, José M., Francisco Perfectti, and Pedro Jordano. 2011. “The Functional Consequences  
687 of Mutualistic Network Architecture.” *PLOS ONE* 6 (1): e16143. <https://doi.org/10.1371/journal.pone.0016143>.
- 689 Harremoës, Peter, and Flemming Topsøe. 2001. “Maximum Entropy Fundamentals.” *Entropy*  
690 3 (3): 191–226. <https://doi.org/10.3390/e3030191>.
- 691 Harte, J., and Erica A. Newman. 2014. “Maximum Information Entropy: A Foundation for  
692 Ecological Theory.” *Trends in Ecology & Evolution* 29 (7): 384–89. <https://doi.org/10.1016/j.tree.2014.04.009>.
- 694 Harte, J., T. Zillio, E. Conlisk, and A. B. Smith. 2008. “Maximum Entropy and the State-  
695 Variable Approach to Macroecology.” *Ecology* 89 (10): 2700–2711. <https://doi.org/10.1890/07-1369.1>.
- 697 Hoorn, Pim van der, Gabor Lippner, and Dmitri Krioukov. 2018. “Sparse Maximum-Entropy  
698 Random Graphs with a Given Power-Law Degree Distribution.” *Journal of Statistical Physics*  
699 173 (3): 806–44. <https://doi.org/10.1007/s10955-017-1887-7>.
- 700 Hortal, Joaquín, Francesco de Bello, José Alexandre F. Diniz-Filho, Thomas M. Lewinsohn,  
701 Jorge M. Lobo, and Richard J. Ladle. 2015. “Seven Shortfalls That Beset Large-Scale  
702 Knowledge of Biodiversity.” *Annual Review of Ecology, Evolution, and Systematics* 46 (1):  
703 523–49. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>.
- 704 Ings, Thomas C., José M. Montoya, Jordi Bascompte, Nico Blüthgen, Lee Brown, Carsten F.  
705 Dormann, François Edwards, et al. 2009. “Review: Ecological Networks Beyond Food

- 706 Webs.” *Journal of Animal Ecology* 78 (1): 253–69. <https://doi.org/10.1111/j.1365-2656.2008.01460.x>.
- 708 Jaynes, E. T. 1957a. “Information Theory and Statistical Mechanics.” *Physical Review* 106 (4):  
709 620–30. <https://doi.org/10.1103/PhysRev.106.620>.
- 710 ———. 1957b. “Information Theory and Statistical Mechanics. II.” *Physical Review* 108 (2):  
711 171–90. <https://doi.org/10.1103/PhysRev.108.171>.
- 712 Jordano, Pedro. 2016. “Sampling Networks of Ecological Interactions.” *Functional Ecology* 30  
713 (12): 1883–93. <https://doi.org/10.1111/1365-2435.12763>.
- 714 Khinchin, A. Ya. 2013. *Mathematical Foundations of Information Theory*. Courier Corpora-  
715 tion.
- 716 MacDonald, Arthur Andrew Meahan, Francis Banville, and Timothée Poisot. 2020. “Revisiting  
717 the Links-Species Scaling Relationship in Food Webs.” *Patterns* 0 (0). <https://doi.org/10.1016/j.patter.2020.100079>.
- 718 Martyushev, L. M., and V. D. Seleznev. 2006. “Maximum Entropy Production Principle in  
719 Physics, Chemistry and Biology.” *Physics Reports-Review Section of Physics Letters* 426  
720 (1): 1–45. <https://doi.org/10.1016/j.physrep.2005.12.001>.
- 721 McCann, Kevin S. 2011. *Food Webs (MPB-50)*. *Food Webs (MPB-50)*. Princeton University  
722 Press. <https://doi.org/10.1515/9781400840687>.
- 723 Milo, R., S. Shen-Orr, S. Itzkovitz, N. Kashtan, D. Chklovskii, and U. Alon. 2002. “Net-  
724 work Motifs: Simple Building Blocks of Complex Networks.” *Science* 298 (5594): 824–27.  
725 <https://doi.org/10.1126/science.298.5594.824>.
- 726 Park, Juyong, and M. E. J. Newman. 2004. “Statistical Mechanics of Networks.” *Physical  
727 Review E* 70 (6): 066117. <https://doi.org/10.1103/PhysRevE.70.066117>.
- 728 Pascual, Department of Ecology and Evolutionary Biology Mercedes, and Visiting Professor  
729 Jennifer A. Dunne. 2006. *Ecological Networks: Linking Structure to Dynamics in Food  
730 Webs*. Oxford University Press, USA.

- 732 Phillips, Steven J., Robert P. Anderson, and Robert E. Schapire. 2006. "Maximum Entropy  
733 Modeling of Species Geographic Distributions." *Ecological Modelling* 190 (3): 231–59.  
734 <https://doi.org/10.1016/j.ecolmodel.2005.03.026>.
- 735 Pielou, Evelyn C. 1975. "Ecological Diversity." In. 574.524018 P5.
- 736 Poisot, Timothee, Benjamin Baiser, Jennifer A. Dunne, Sonia Kefi, Francois Massol, Nico-  
737 las Mouquet, Tamara N. Romanuk, Daniel B. Stouffer, Spencer A. Wood, and Dominique  
738 Gravel. 2016. "Mangal - Making Ecological Network Analysis Simple." *Ecography* 39 (4):  
739 384–90. <https://doi.org/10.1111/ecog.00976>.
- 740 Poisot, Timothée, Gabriel Bergeron, Kevin Cazelles, Tad Dallas, Dominique Gravel, Andrew  
741 MacDonald, Benjamin Mercier, Clément Violet, and Steve Vissault. 2021. "Global Knowl-  
742 edge Gaps in Species Interaction Networks Data." *Journal of Biogeography* 48 (7): 1552–  
743 63. <https://doi.org/10.1111/jbi.14127>.
- 744 Poisot, Timothée, and Dominique Gravel. 2014. "When Is an Ecological Network Complex?  
745 Connectance Drives Degree Distribution and Emerging Network Properties." *PeerJ* 2: e251.  
746 <https://doi.org/10.7717/peerj.251>.
- 747 Poisot, Timothée, Daniel B. Stouffer, and Dominique Gravel. 2015. "Beyond Species: Why  
748 Ecological Interaction Networks Vary Through Space and Time." *Oikos* 124 (3): 243–51.  
749 <https://doi.org/10.1111/oik.01719>.
- 750 Pomeranz, Justin P. F., Ross M. Thompson, Timothée Poisot, and Jon S. Harding. 2019. "In-  
751 ferring Predatorprey Interactions in Food Webs." *Methods in Ecology and Evolution* 10 (3):  
752 356–67. <https://doi.org/10.1111/2041-210X.13125>.
- 753 Pomeranz, Justin Page, Ross M. Thompson, Timothée Poisot, Jon S. Harding, and Justin P. F.  
754 Pomeranz. 2018. "Data from: Inferring Predator-Prey Interactions in Food Webs." Dryad.  
755 <https://doi.org/10.5061/DRYAD.K59M37F>.
- 756 Shannon, C. E. 1948. "A Mathematical Theory of Communication." *The Bell System Technical  
757 Journal* 27 (3): 379–423. <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>.
- 758 Staniczenko, Phillip P. A., Jason C. Kopp, and Stefano Allesina. 2013. "The Ghost of Nested-  
759 ness in Ecological Networks." *Nature Communications* 4 (1): 1391. <https://doi.org/>

- 760 10.1038/ncomms2422.
- 761 Stock, Michiel, Timothée Poisot, and Bernard De Baets. 2021. “Optimal Transportation Theory  
762 for Species Interaction Networks.” *Ecology and Evolution* 11 (9): 3841–55. <https://doi.org/10.1002/ece3.7254>.
- 763
- 764 Stouffer, Daniel B., and Jordi Bascompte. 2011. “Compartmentalization Increases Food-Web  
765 Persistence.” *Proceedings of the National Academy of Sciences* 108 (9): 3648–52. <https://doi.org/10.1073/pnas.1014353108>.
- 766
- 767 Stouffer, Daniel B., Juan Camacho, Wenxin Jiang, and Luis A. Nunes Amaral. 2007. “Evidence  
768 for the Existence of a Robust Pattern of Prey Selection in Food Webs.” *Proceedings of the  
769 Royal Society B-Biological Sciences* 274 (1621): 1931–40. <https://doi.org/10.1098/rspb.2007.0571>.
- 770
- 771 Strydom, Tanya, Michael D. Catchesen, Francis Banville, Dominique Caron, Gabriel Dansereau,  
772 Philippe Desjardins-Proulx, Norma R. Forero-Muñoz, et al. 2021. “A Roadmap Towards  
773 Predicting Species Interaction Networks (across Space and Time).” *Philosophical Trans-  
774 actions of the Royal Society B: Biological Sciences* 376 (1837): 20210063. <https://doi.org/10.1098/rstb.2021.0063>.
- 775
- 776 Strydom, Tanya, Giulio V. Dalla Riva, and Timothee Poisot. 2021. “SVD Entropy Reveals the  
777 High Complexity of Ecological Networks.” *Frontiers in Ecology and Evolution* 9: 623141.  
778 <https://doi.org/10.3389/fevo.2021.623141>.
- 779
- 780 Vázquez, Diego P. 2005. “Degree Distribution in Plantanimal Mutualistic Networks: Forbidden  
781 Links or Random Interactions?” *Oikos* 108 (2): 421–26. <https://doi.org/10.1111/j.0030-1299.2005.13619.x>.
- 782
- 783 Williams, Richard J. 2011. “Biology, Methodology or Chance? The Degree Distributions of  
784 Bipartite Ecological Networks.” *PLOS ONE* 6 (3): e17645. <https://doi.org/10.1371/journal.pone.0017645>.
- 785
- 786 Williams, Richard J., and Neo D. Martinez. 2000. “Simple Rules Yield Complex Food Webs.”  
*Nature* 404 (6774): 180–83. <https://doi.org/10.1038/35004572>.




Figure 1: Relative number of predators ( $k_{in}$ ) as a function of their relative number of prey ( $k_{out}$ ) for each species in (a) empirical and (b) joint degree sequences obtained from the analytical MaxEnt model. Empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake datasets (our complete dataset). The predicted joint degree sequences were obtained after sampling one realization of the joint degree distribution of maximum entropy for each network, while keeping the total number of interactions constant. (c) Difference between predicted and empirical values when species are ordered according to their total degrees. In each panel, each dot corresponds to a single species in one of the networks.




Figure 2: (a) Probability density of KL divergence between in and out-degree sequences of empirical and predicted joint degree sequences. (b) Difference between the KL divergence of empirical and predicted joint degree sequences as a function of connectance. In both panels, empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake datasets (our complete dataset). The predicted joint degree sequences were obtained after sampling one realization of the joint degree distribution of maximum entropy for each network, while keeping the total number of interactions constant.




Figure 3: (a) Probability density of the mean degree of a food web obtained using different values of species richness  $S$ . The number of interactions  $L$  was simulated 1000 times using the flexible links model fitted to all empirical networks in our complete dataset (eq. 17). The mean degrees  $2L/S$  were then obtained from these simulated values. (b) Degree distributions of maximum entropy for a network of  $S = 27$  species and different numbers of interactions. The numbers of interactions correspond to the lower and upper bounds of the 67, 89, and 97 percentile intervals (PI), as well as the median (in blue), of the counterfactuals of the flexible links model. Each degree distribution of maximum entropy was obtained using eq. 14 after solving numerically eq. 15 using different values of the mean degree constraint.




Figure 4: Relationship between the structure of empirical and maximum entropy food webs. Empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake food webs (our complete dataset). Maximum entropy networks were obtained using the type II heuristic MaxEnt model based on the joint degree sequence. (a) Nestedness (estimated using the spectral radius of the adjacency matrix), (b) the maximum trophic level, (c) the network diameter, and (d) the SVD entropy were measured on these empirical and maximum entropy food webs. The identity line is plotted in each panel.




Figure 5: Motifs profile of empirical and predicted food webs. Empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake datasets (our complete dataset). Null 1: Type I null model based on connectance. MaxEnt 1: Type I heuristic MaxEnt model based on connectance. Null 2: Type II null model based on the joint degree sequence. MaxEnt 2: Type II heuristic MaxEnt model based on the joint degree sequence. Boxplots display the median proportion of each motif (middle horizontal lines), as well as the first (bottom horizontal lines) and third (top horizontal lines) quartiles. Vertical lines encompass all data points that fall within 1.5 times the interquartile range from both quartiles, and dots are data points that fall outside this range. Motifs names are from Stouffer et al. (2007).




Figure 6: Pairwise relationships between motifs proportions of empirical and predicted food webs. Empirical networks include most food webs archived on Mangal, as well as the New Zealand and Tuesday Lake datasets (our complete dataset). Null 1: Type I null model based on connectance. MaxEnt 1: Type I heuristic MaxEnt model based on connectance. Null 2: Type II null model based on the joint degree sequence. MaxEnt 2: Type II heuristic MaxEnt model based on the joint degree sequence. Regression lines are plotted in each panel. Motifs names are from Stouffer et al. (2007).