

# Supporting Information 1

## Proceedings of the Royal Society B

DOI: 10.1098/rspb.2019.2546

Article title: Extinction pulse at Eocene–Oligocene boundary drives diversification dynamics of the two Australian temperate floras

Francis J. Nge<sup>1,2,4</sup>, Ed Biffin<sup>1,2</sup>, Kevin R. Thiele<sup>3</sup>, Michelle Waycott<sup>1,2</sup>

<sup>1</sup>School of Biological Sciences, Faculty of Science, The University of Adelaide, Adelaide, South Australia, 5005, Australia

<sup>2</sup>State Herbarium of South Australia, G.P.O. Box 1047, Adelaide, South Australia 5001

<sup>3</sup>School of Biological Sciences, University of Western Australia, 35 Stirling Hwy, Crawley (Perth), WA 6009, Australia

Author for correspondence: Francis Nge

<sup>4</sup>Corresponding author Email: francis.nge@adelaide.edu.au

## Figures S10–15

**Fig. S10** BAMM estimates of diversification rate for *Acacia* (Fabaceae) under different global sampling fraction regimes: **a)** 20%; **b)** 40%; **c)** 60%; **d)** 80%; **e)** 100%. Estimated rates of net species diversification ( $\text{sp sp}^{-1} \text{My}^{-1}$ ) are colour-coded as indicated (red–blue; high–low rate).

**Fig. S11** BAMM estimates of diversification rate for Epacrids (Ericaceae) under different global sampling fraction regimes: **a)** 20%; **b)** 50%; **c)** 80%; **d)** 100%. Estimated rates of net species diversification ( $\text{sp sp}^{-1} \text{My}^{-1}$ ) are colour-coded as indicated (red–blue; high–low rate).


**Fig. S12** BAMM estimates of diversification rate for *Eucalyptus* (Myrtaceae) under different global sampling fraction regimes: **a)** 20%; **b)** 50%; **c)** 70%; **d)** 90%. Estimated rates of net species diversification ( $\text{sp sp}^{-1} \text{My}^{-1}$ ) are colour-coded as indicated (red–blue; high–low rate).

**Fig. S13** BAMM 95% credible rate shift configurations for *Acacia* (Fabaceae) under different global samplping fraction regimes: **a)** 20%; **b)** 40%; **c)** 60%; **d)** 80%; **e)** 100%. F = probability; percentage of samples in posterior assigned to shift configurations.


**Fig. S14** BAMM 95% credible rate shift configurations for Epacrids (Ericaceae) under different global samplping fraction regimes: **a)** 20%; **b)** 50%; **c)** 80%; **d)** 100%. F = probability; percentage of samples in posterior assigned to shift configurations.

**Fig. S15** BAMM 95% credible rate shift configurations for *Eucalyptus* (Myrtaceae) under different global samplping fraction regimes: **a)** 20%; **b)** 50%; **c)** 70%; **d)** 90%. F = probability; percentage of samples in posterior assigned to shift configurations.


**Fig. S10**




**Fig. S11**




**Fig. S12**




**Fig. S13a**




**Fig. S13b**




**Fig. S13c**




**Fig. S13d**




**Fig. S13e**




**Fig. S14a**




**Fig. S14b**




**Fig. S14c**




**Fig. S14d**




**Fig. S15a**



**Fig. S15b**



**Fig. S15c**



**Fig. S15d**

