

IC (i):

Calculating coefficients bn :

$$bn := \frac{\int_0^L M \cdot \sin\left(\frac{n \cdot \pi \cdot x}{L}\right) \cdot dx}{\int_0^L \sin^2\left(\frac{n \cdot \pi \cdot x}{L}\right) \cdot dx} \text{ assuming } (M > 0, n, \text{integer})$$

$$bn := - \frac{2 M ((-1)^n - 1)}{\pi n} \quad (1)$$

(a) $u(x,t)$ is equal to:

$$u := bn \cdot \sin\left(\frac{n \cdot \pi \cdot x}{L}\right) \cdot \exp\left(-\left(\frac{n \cdot \pi}{L}\right)^2 \cdot D \cdot t\right)$$

$$u := - \frac{2 M ((-1)^n - 1) \sin\left(\frac{n \pi x}{L}\right) e^{-\frac{n^2 \pi^2 D t}{L^2}}}{\pi n} \quad (2)$$

(b) Then, calculating the average temperature at $t=0$:

$$avg := \left(\frac{1}{L}\right) \cdot \int_0^L bn \cdot \sin\left(\frac{n \cdot \pi}{L} \cdot x\right) \cdot \exp\left(-\left(\frac{n \cdot \pi}{L}\right)^2 \cdot D \cdot t\right) \cdot dx$$

$$avg := \frac{2 M ((-1)^n - 1) e^{-\frac{n^2 \pi^2 D t}{L^2}} (\cos(n \pi) - 1)}{n^2 \pi^2} \quad (3)$$

$avg := \text{subs}(t=0, M=100, L=10, D=1, avg)$

$$avg := \frac{200 ((-1)^n - 1) e^0 (\cos(n \pi) - 1)}{n^2 \pi^2} \quad (4)$$

$psum := \text{sum}(avg, n = 1 .. 1000) :$
 $\text{evalf}(psum)$

$$99.95947149 \quad (5)$$

(c) Expression for calculating the average temperature at time t :

$$avg_expr := \frac{\exp\left(-\left(\frac{n \cdot \pi}{L}\right)^2 \cdot D \cdot t\right)}{L} \cdot \int_0^L \left(-\frac{2 \cdot M \cdot ((-1)^n - 1)}{\pi \cdot n}\right) \cdot \sin\left(\frac{n \cdot \pi}{L} x\right) \cdot dx$$

$$avg_expr := \frac{2 M ((-1)^n - 1) e^{-\frac{n^2 \pi^2 D t}{L^2}} (\cos(n \pi) - 1)}{n^2 \pi^2} \quad (6)$$

(d) The slowest decaying term occurs when $n = 1$. Then

$initial_value := 100 :$

$final_value := 0.1 \cdot initial_value$

$$final_value := 10.0 \quad (7)$$

$$equ := subs(M=100, n=1, L=10, D=1, avg_expr) - final_value$$

$$equ := -\frac{400 e^{-\frac{\pi^2 t}{100}} (\cos(\pi) - 1)}{\pi^2} - 10.0 \quad (8)$$

$$solve(equ, t)$$


$$21.20213514 \quad (9)$$

(e) Creating Partial Sums:

with(plots):


psum := subs(M=100, D=1, L=10, sum(u, n=1..300)):

animate(psum, x=0..10, t=0..23)



curves := [seq(subs(t=2*m, psum), m=0..11)]:

plot(curves, x=0..10)



We can see that approaches 21.20213514 time units, the amplitude of the curves (temperature) get smaller. We can also see the Gibbs phenomenon for the $t=0$ curve.

For initial condition 1.ii:

$$f := \text{piecewise}\left(x > \frac{L}{2}, M, x < \frac{L}{2}, 0\right)$$

$$f := \begin{cases} M & \frac{L}{2} < x \\ 0 & \frac{L}{2} < x \end{cases} \quad (1)$$

(a) Calculating the solution u(x,t):

We know the family of solutions is as follows:

$$u := bn \cdot \sin\left(\frac{n \cdot \text{Pi} \cdot x}{L}\right) \cdot \exp\left(-\left(\frac{n \cdot \text{Pi}}{L}\right)^2 \cdot D \cdot t\right)$$

$$u := - \frac{2 M \left(\cos\left(\frac{n \pi}{2}\right) - 1\right) \sin\left(\frac{n \pi x}{L}\right) e^{-\frac{n^2 \pi^2 D t}{L^2}}}{n \pi} \quad (2)$$

Calculating the coefficients bn for t=0:

$$bn := \frac{\text{int}\left(M \cdot \sin\left(\frac{n \cdot \text{Pi} \cdot x}{L}\right), x = 0 .. \frac{L}{2}\right)}{\text{int}\left(\sin^2\left(\frac{n \cdot \text{Pi} \cdot x}{L}\right), x = 0 .. L\right)} \text{assuming}(L > 0, n, \text{integer})$$

$$bn := - \frac{2 M \left(\cos\left(\frac{n \pi}{2}\right) - 1\right)}{n \pi} \quad (3)$$

Then, the family of solutions u(x,t) is:

$$u := bn \cdot \sin\left(\frac{n \cdot \text{Pi} \cdot x}{L}\right) \cdot \exp\left(-\left(\frac{n \cdot \text{Pi}}{L}\right)^2 \cdot D \cdot t\right)$$

$$u := - \frac{2 M \left(\cos\left(\frac{n \pi}{2}\right) - 1\right) \sin\left(\frac{n \pi x}{L}\right) e^{-\frac{n^2 \pi^2 D t}{L^2}}}{n \pi} \quad (4)$$

(b) Average temperature at time t=0:

$$\text{avg_t0} := \text{simplify}\left(\left(\frac{1}{L}\right) \cdot \text{int}(u, x = 0 .. L) \text{assuming}(L > 0, n, \text{integer})\right) :$$

$$\text{avg_t0} := \text{simplify}(\text{subs}(M = 100, t = 0, L = 10, D = 1, \text{avg_t0})) :$$

$$\text{avg_t0} := \text{evalf}(\text{sum}(\text{avg_t0}, n = 1 .. 900))$$

$$\text{avg_t0} := 49.97748417 \quad (5)$$

(c) Average temperature expression:

$$\text{avg_temp} := \left(\frac{1}{L}\right) \cdot \text{int}(u, x = 0 .. L) \text{assuming}(L > 0, D = 1, n, \text{integer}) :$$

$$\text{avg_temp} := \text{subs}(L = 10, M = 100, \text{avg_temp})$$

$$avg_temp := \frac{200 \left(\cos\left(\frac{n \pi}{2}\right) - 1 \right) e^{-\frac{n^2 \pi^2 t}{100}} (-1 + (-1)^n)}{n^2 \pi^2} \quad (6)$$

(d) Estimating how long it takes for the average temperature to decrease to 10% of its initial value (for the slowest decaying term n=1):

$$final_value := 0.1 \cdot 50; \quad final_value := 5.0 \quad (7)$$

$$equ := subs(n = 1, avg_temp - final_value)$$

$$equ := -\frac{400 \left(\cos\left(\frac{\pi}{2}\right) - 1 \right) e^{-\frac{t \pi^2}{100}}}{\pi^2} - 5.0 \quad (8)$$

$$equ := solve(equ, t)$$

$$equ := 21.20213514 \quad (9)$$

It takes 21.20213514 times units (approximately) to go from the average temperature at t=0 of 50 to the average temperature of 5.

(e) Animation, partial sums, and plots


$$u := subs(M = 100, L = 10, D = 1, u)$$

$$u := -\frac{200 \left(\cos\left(\frac{n \pi}{2}\right) - 1 \right) \sin\left(\frac{n \pi x}{10}\right) e^{-\frac{n^2 \pi^2 t}{100}}}{n \pi} \quad (10)$$


$$psum := sum(u, n = 1 .. 300) :$$

$$with(plots) :$$

$$animate(psum, x = 0 .. 10, t = 0 .. 21.5)$$



```
curves := [seq(subs(t=2*m, psum), m=0..11)]:
plot(curves, x=0..10)
```



We can see that the average temperature amplitud is lower as t increases.